

- 200 Discomyceteæ, Pezizeæ, Hyalosporæ, Sclerotinia.
hyalinis, 18-20 \approx 9, monostichis; paraphysibus filiformibus sursum
clavulatis; conidiis sphæricis.
Hab. ad baccas Vaccinii Myrtilli *Sclerotio* infectas in Ger-
mania. — Status conidicus in ramulis oritur.
18. **Sclerotinia Vaccinii** Woron. Memoir. Acad. S. Petersb. 1888. — 814
Affinis *Scl. baccarum*: ascomatibus cupulatis brunneis; stipite sæpe
prælongo; ascis cylindraceis, apice canaliculatis jodoque cœrule-
scentibus, 8-sporis; paraphysibus septulatis, clavulatis; sporidiis oc-
tonis ovoideis, 14-17 \approx 7-9, quatuor paullulo minoribus; conidiis
3-4 \approx 2-2,5, limoniformibus, hyalinis.
Hab. in baccis, *Sclerotio* correptis, Vaccinii Vitis-idææ in
Fennia. — Status conidicus, *Moniliam* sistens, in ramulis foliisque
oritur.
19. **Sclerotinia Oxycoeci** Woron. Memoir. Acad. S. Petersb. 1888. — 815
A *Scl. Vaccinii* differt ascomatibus gracilioribus; sporidiis octonis,
quatuor conspicue minoribus, 12-14 \approx 6-7; conidiis 2,5-2,8 \approx 2,2.
Hab. in baccis, *Sclerotio* correptis, Vaccinii Oxycoeci in
Fennia. — Status conidicus in ramulis foliisque oritur.
20. **Sclerotinia megalospora** Woron. Memoir. Acad. S. Petersb. 1888. 816
— A *Sclerot. Vaccinii* differt conidiis rotundioribus, 2,4-3 μ . diam.;
sporidiis multo majoribus octonis, 20-25 \approx 14-16, omnibus æquali-
bus, fertilibus.
Hab. in foliis Vaccinii uliginosi in Fennia. — Status
conidicus in foliis ramulisque oritur.
21. **Sclerotinia gracilipes** (Cooke) Sacc. *Peziza (Hymenoscypha) gra-* 817
cilipes Cooke Hedw. 1875, p. 82, Disc. U. S. I, p. 294. — Stipi-
tata, cupulata, submembranacea, delicatula, fuscescens, discoidea,
dēnum convexa 2-3 mm., e *Sclerotio* compresso, rugoso, atro ena-
ta; stipite tenuissimo 2-3 cm. long.; sporidiis oblongis 8 \approx 4.
Hab. ad petala decidua New Jersey Amer. bor. (ELLIS). — Af-
finis *Sclerot. ciborioidi* (*S. Trifoliorum*):
22. **Sclerotinia incondita** Ellis Bull. Torr. Cl. 1881, p. 74 (*Peziza*) 818
Ell. N. Am. F. n. 391. — Solitaria, stipitata, pallide castanea, di-
sco plano vel convexo, 3-6 mm. diam., absque margine distincto;
stipite centrali, erecto, brunneolo, circ. 12-13 mm. alto, fimo;
ascis cylindraceo-clavatis, sessilibus circ. 7,5 μ . longis, tetrasporis;
sporidiis ellipticis v. ovato-ellipticis 10-12,5 \approx 6-7,5 hyalinis.
Hab. inter muscos in paludibus Americæ borealis. — Stipes e
Sclerotio angusto, cupuliformi, atro exsurgens. *Sclerotium* habitu
ei *Sclerotiniæ gracilipedis* Cooke simile.

23. **Sclerotinia Tuba** Batsch Elench. p. 122 (*Pez.*) nec *Peziza Tuba* 819
Bolt. Fr. Syst. Myc. II, p. 58, Pers. Myc. Eur. p. 235, *P. perennis* Pers. Syn. p. 644, Michel. t. 86, fig. 10. — Cæspitosa, fulva,
cupula infundibuliformi, levi, radice tuberosa nigra perenni.

Hab. in silvis Castaneæ Italiæ, etiam in America boreali (SCHWEINITZ). — Stipite subflexuoso, levi, sursum in cupulam orbicularem 7-8 cm. altam, 3 cm. latam dilatato. Radix perennis est ac tribus plerumque tuberculis constat, quorum medium Pezizas profert, alterum effœtum anni præcedentis, tertium omnium minimum anno sequente fructificat. Hoc ex capite cum bulbis Orchidearum congruit. Michelius in horto coluit.

CIBORIA Fuckel Symb. Mycol. p. 311 [1869] (Etym. *ciborium* ob formam), *Rutstroëmia* Karst. ex parte. — Ascoma longe firmeque sæpe longissime stipitatum, media magnitudine, majuscum, ceraeum, extus glabrum, furfuraceum, apertum, patellæforme, marginatum; disco sæpe discolori. Asci elongati, 8-spori. Sporidia oblongo-ovata, cylindracea lanceolatave, continua, hyalina, rarius appendiculata. Paraphyses variæ. — Plerumque vernalis.

1. **Ciboria pseudotuberosa** Rehm Ascom. n. 106 (1870), Pat. Tab. 378, 820
Sclerotinia Batschiana Zopf (1880) in Zopf et Syd. myc. march. 50 c. icon. *Peziza glandicola* Doass. in Revue myc. n. 10, p. 49, Cfr. Grevillea IV, pl. 65, f. 288. — Gregaria, 2 cm. long., stipitata, stipite tenui, subflexuoso, sicco plerumque longitudinaliter striato, olivaceo, sicca infundibuliformis, subcom plicata, olivaceo-fusca, humida explanata, subpatellæformis, disco livido-fusco, 5-7 mm. diam.; ascis clavatis, 8-sporis, 120 ± 6-7; sporidiis obtuse-oblongis vel ovatis, glabris, hyalinis, 8-9 ± 4-4,5, continuis, monostichis in superiore ascii parte; paraphysibus filiformibus, superne sensim (3 µ.) incrassatis, hyalinis.

Hab. ad cotyledones putridas Quercuum in silvis prope Windsheim Franconia bav., in Gallia et Britannia.

2. **Ciboria Juglandis** Preuss Hoyersw. n. 154 (*Peziza*). — Levis, fir- 821
mula, hypocrateriformis, disco ochraceo-fusco, margine integerri-
mo, dilutiore, extus albido; stipite longo filiformi, deorsum atte-
nuato, nigrescente cum radiculis; ascis cylindricis; sporidiis ovatis
albis; paraphysibus simplicibus.

Hab. in nucleis (cotyledonibus) rancidis Juglandium, terra
tectis, Hoyerswerda.

3. **Ciboria amentacea** (Balb.) Fuck. Symb. p. 311, *Peziza amenta-* 822

cea Balb. in Act. Taur. II, t. 2, *Pez. julacea* Pers. M. E. I, p. 311, *Rutstroëmia ament*. Karst. M. F. I, p. 106. — Cupulis longe stipitatis, circ. 2-3 cm. altis, primo concavis, demum explanatis, extus fusis, disco pallidiore, 5 mm. latis; stipite firme; ascis elongatis, stipitatis, 8-sporis, 150 \pm 6; sporidiis monostichis, ovatis, continuis, hyalinis, 6 \pm 3.

Hab. ad amenta masculina *Alni glutinosæ* et *Salicu*m in Italia, Germania, Fennia, Britannia. — Sec. Phillips sporidia 12 \pm 5,5, hinc species a *C. Cauco* ægre distinguitur.

4. **Ciboria Caucus** (Rebent.) Fuck. Symb. p. 311, *P. amentalis* Schum. 823
Saell. p. 418, *P. Caucus* Rebent. Neom. p. 386, t. 4, f. 17, Fr. S. M. II, p. 126, Krombh. t. V, f. 37-39, Pat. t. 483. — Pallide fuscescens, cupula vasculiformi, margine erecto, stipite brevi crassiusculo subflexuoso; ascis elongatis, 8-sporis, 130 μ . long.; sporidiis oblique monostichis, ovatis, inæquilateralibus, 9 \pm 4, hyalinis (Fuck.), 10 \pm 6 (Phill.).

Hab. in amentis dejectis *Populi albæ* et *Tremulæ* in Dania, Germania, Britannia, Gallia. — Stipes subinde filiformis, tenuis, typice 5-10 mm. long. Cupula ceracea, 5 mm. lata, exacte vasculiformis, intus fusco-umbrina, extus pallidior, basi subuplicata.

5. **Ciboria echinophila** (Bull.) Sacc. F. Ven. Ser. IV, n. 88, F. ital. 824
fig. 1297, *P. echinophila* Bull. Champ. p. 235, t. 500, f. 1, Pers. Myc. Eur. p. 278, Pat. Tab. 84. — Cupula plano-infundibuliformi, dilute cinnamomea; stipite longo dilutiore, primo subtomentoso; ascis cylindraceis, 120-140 \pm 12-15, breve crassiuscule stipitatis, apice abrupte attenuato-truncatis, paraphysibus filiformibus obval-latis, 8-sporis; sporidiis monostichis cylindraceis, 18-20 \pm 5, vehe-menter curvatis, utrinque obtusis sed initio minute apiculatis, plu-riguttulatis dein 1-nucleato-farctis, hyalinis.

Hab. ad involucra putrescentia *Castaneæ* in Gallia, Helvetia, Britannia, Germania, Italia, Belgio. — Ad *Phialeam* accedit. Stipes longitudine varius, 12-18 mm. circiter. Cupula 2-4 mm. lata, firma.

6. **Ciboria Urnula** Weinm. Hym. Gaster. p. 459. — Solitaria, firmula; 825
ascomate oblongiusculo, obscure fusco, ore contracto, subrevoluto, extus stipiteque elongato pruinoso-villosiusculis.

Hab. semper in fructibus putrescentibus *Vaccinii Vitis-Idææ* inter cæspites humidos *Sphagnorum* in Rossia. — Ascoma 3-4 mm. altum, 2-3 mm. latum. Margo primo erectus, dein sub-revolitus, integer 1. in statu vetusto lacinulatus. Stipes $2\frac{1}{2}$ - $3\frac{1}{2}$

cm. longus, tenuis, æqualis, ut plurimum inter sphagna humida latens. An *Sclerotinia*?

7. **Ciboria rufo-fusca** (Weber.) Sacc. *Pez. rufo-fusca* Web. pag. 7, 826 t. III, fig. 4. — Solitaria vel gregaria, rubescenti-brunnea; ascocarneis ceraceis, cupulato-patellatis, prima ætate sessilibus, dein stipitatis ad 8 mm. latis; stipite tereti, exili, ascoma subæquante; ascis minutis, linearibus, octosporis; sporidiis ellipticis, subinde ovoideis, contentu granuloso fœtis, $7,5 \pm 3,7$; paraphysibus filiformibus, superne leniter ampliatis.

Hab. in acubus et squamis Abietis pectinatae et Abietis excelsæ pr. Landeck Germaniæ.

8. **Ciboria strobilina** (A. S.) Sacc. *P. strobilina* A. et S. Consp. p. 313 827 (ut var. *Scl. tuberosæ*) *Phialea strobilina* Quél. 10 Suppl. t. IX, f. 8, Gill. Disc. c. ic. (non *Pez. strobilina* Fr.). — Cyathiformis dein expansa undulata, e lacteo bisa v. fuscescens; stipite gracili longo, 1-3 cm. flexuoso, basi attenuato nigricante; sporidiis oblongis, hyalinis, $10-12 \pm 5-6$.

Hab. squamas strobilarum Abietum in Lusatia, Gallia. — Ascoma 1-2 cm. lat.

9. **Ciboria subularis** Bull. champ. pag. 236, t. 500, f. 2 (*Peziza*), *P. subulipes* Pers. Myc. Eur. p. 282, *Pez. Carpini* Batsch f. 150 (?), *Phialea subularis* Gill. Disc. c. ic. — Lateritia, fragilis, cupula hypocrateriformi, integerrima, stipite longissimo tenui, flexuoso; ascis tereti-clavatis; sporidiis oblongis, 2-guttatis, hyalinis, $17-20 \pm 4-5$; paraphysibus filiformi-clavulatis.

Hab. supra semina semiputrida Helianthi annui et Bidens tripartitæ in Gallia et Belgio, in fructibus Angelicæ in Anglia. — Ascoma 3-4 mm. lat.; stipes 12-14 mm. long.; specimina anglica extus albida. Similis *Ciboria Friesii*, loco et substantia fragili distinguenda.

10. **Ciboria firma** (Pers.) Fuck. Symb. myc. p. 312, Sacc. F. it. p. 1295, 829 *P. ochroleuca* Bolt. fung. t. 105, f. 1, Sowerb. t. 115, *P. firma* Pers. syn. p. 658, Fr. S. M. II, 117, Pat. Tab. 78, Gonn. et Rab. t. VI, f. 3, *Phialea* Gill. Disc. c. ic., *Rutströmia* Karst. M. T. I, p. 108. — Cupula infundibuliformi, dein dilatata repanda dilute fusca; stipite longo deorsum attenuato nigrescente; ascis elongatis, stipitatis, 8-sporis, 16 ± 12 ; sporidiis elongato-distichis, oblongo-cylindraceis, valde curvatis, utrinque rotundato-obtusis, hyalinis, 2-4-guttulatis, $14-16 \pm 4-5$.

Hab. ad ramos dejectos putrescentes in Suecia, Britannia, Bel-

gio, Fennia, Italia, Germania, in America boreali et Tasmania. — Stipes tenax, filiformis, glaber 12-20 mm. circ. longus: Cupula 7-12 mm. et ultra lata, integerrima, subrepanda. Colore variat rarius tota ochroleuca, saepius extus dilutior, subochroleuca, disco umbrino, olivaceo etc. *P. alutacea* et *stipitum* Schum. Saell. p. 431, colore et statura minore aliquantulum recedunt. — ***C. macrospora** Sacc. F. ital. fig. 1296: ascis majoribus nempe 150 \approx 15-16, clavatis, apice obtusatis, paraphysibus filiformibus sursum incrassatis obvallatis, octosporis; sporidiis majoribus, nempe 25-28 \approx 7, oblongo-fusoideis, saepe curvulis, 2-guttatis, hyalinis. In foliis galbulisque putrescentibus Juniperi communis, Conegliano, Ital. bor. (SPEGAZZINI).

11. **Ciboria bolaris** (Batsch) Fuck. Symb. p. 311, *P. bolaris* Batsch. 830 Cont. I, p. 221, f. 155 (f. 144, etiam prope accedit) Fr. S. M. II, p. 112. — Infundibuliformis, cupula hemisphærica, fibrilloso-venosa, subochracea, disco badio, stipite brevi nigrescente; ascis stipitatis, cylindraceis, 8-sporis, 208 \approx 16; sporidiis primo distichis, demum submonostichis, oblongo-ovatis, hyalinis, distincte scrobiculatis, 20 \approx 8, nucleo granuloso, non guttulato (Fuck.).

Hab. ad ramos sicclos deciduos Coryli, Carpinii, Ilicis, Humuli, Salicis, in Suecia, Britannia, Germania, Italia. — Sec. Phill. sporidia 14-16 \approx 4-6, 2-guttata. Ceracea, firma, 4-5 mm. alta et lata, extus e fibrillis rugoso-venosa, dilutior; margine subserrato, in siccis inflexo, junior reliquorum more passim subvillosa, mox vero glabrescens. Variat ceterum colore plus minus obscuro et stipitis longitudine. — β . **C. explanata** Holmsk. l. c. p. 35, t. 17, Pers. Eur. p. 273: subfuscata, disco subrotundo planiusculo, margine denticulato. Junior sessilis, cupulæformis, extus pilis tenuissimis obsita, adulta glabra, subtenuis, 8-10 mm. lata; stipite æquali. Forsan diversa; sed loco omnino convenit cum vulgari.

12. **Ciboria clavipes** Hazsl. Einig. Disc. p. 13, tab. III, f. 8. — Claviformis, basi incrassata rotundataque, libere inserta, lutea; stipite brunneo-lutescente, diametro ascomatis hemisphærici fere triplo longiori; disco profunde concavo; ascis angustis, clavatis, in stipitem longiusculum, tenuem desinentibus, octosporis; sporidiis primo partem superiorem dein inferiorem ascii occupantibus, cylindraceis vel clavatis, 16-20 μ . longis, utrinque rotundatis, 3-4-nucleatis.

Hab. in caulis Artemisiae vulgaris, prope Eperies Hungariæ.

13. **Ciboria fallax** Bomm. et Rouss. Fl. mycol. pag. 135. — Cupula 832
 4-5 mm. diam., parum concava dein subplana, extus fibrillis ad-
 pressis vestita; disco armeniaco, fracto (ut in Stereis quibusdam)
 rufescente; stipite 4-5 mm. longo, leniter fibrilloso apiceque am-
 pliato; sporidiis 24-6, oblongis, uno polo obtusis, altero subacutis,
 pluriguttulatis, utrinque appendicula hyalina, ægre visibili, cum
 sporidio angulum rectum formante instructis.

Hab. ad caules herbaceos et sarmenta Rubi locis uidis in Belgio.

14. **Ciboria carniolica** Rehm in Voss Mat. Krains V, p. 226, f. 1, a-d 833
 et Ascom. n. 903. — Turbinata, stipitata, firma, solitaria vel 2-3
 gregaria, disco subplano, fuscescente, pruinoso, margine tenui dilu-
 tiore cincto, orbiculari, interdum irregulariter eroso, 1-1,5 mm.
 diam., stipite flavescente, sicco rugoso, 3 mm. alt., $\frac{1}{4}$ - $\frac{1}{2}$ mm. diam.;
 ascis subcylindraceis, 8-sporis, 50-5; sporidiis 2-stichis oblongis,
 obtusis, rectis, hyalinis, 1-cellularibus, 5-7-2; paraphysibus ascos
 superantibus, filiformibus, apice sensim clavulatis, hyalinis.

Hab. ad radices (Quercus?) demersas in silvis prope Laibach
 Carnioliae. — A *Lachnella pygmaea* satis differre videtur.

15. **Ciboria calopus** (Fr.) *Peziza calopus* Fr. Observ. 2, p. 307, Syst. 834
 myc. II, p. 131, *Helotium c.* Fr. S. v. Sc. p. 355, Fuck. Symb.
 mycol. App. II, p. 62, fig. 36. — Cupulis sparsis, ceraceis, ple-
 rumque longe stipitatis sed saepè subsessilibus subturbinatisque,
 orbicularibus, in stipitem attenuatis, 3-5 mm. latis, primo conca-
 vis, demum explanatis, margine obsoleto, extus cupulis stipitibus-
 que glabris, amœne roseis, siccis plicatis, disco flavescente-ochraceo,
 sicco fusco; ascis cylindraceis oblique stipitatis, 8-sporis, 144-16;
 sporidiis in asci superiori parte distichis, late fusiforme-ovatis, con-
 tinuis, biguttulatis, hyalinis, 16-20-8; paraphysibus simplicibus,
 apice vix crassioribus.

Hab. ad stipites, culmos, folia, radices putres in Suecia, Ger-
 mania et Gallia. — Fungus conidiophorus *Dacriomycetem* exhi-
 bens, primo sub foliorum vivorum epidermide nidulans, demum tur-
 gidus liberque, versiformis, plerumque elongatus, carnosus, amœne
 roseus, hyphis niveis cinctus; intus hyphis ramosis contextus; hy-
 phis apice conidiophoris; conidia ovata, paullo breviora quam asco-
 sporæ. Fuck. l. c.

16. **Ciboria ciliatospora** Fuck. Symb. p. 311, t. IV, f. 36, Sacc. et 835
 Berl. Misc. Mycol. II, p. 29, t. VIII, f. 8. — Glabra, longissime
 stipitata, ad 2-3 cm. alta, pallide fusca; cupulis turbinatis in sti-
 pitem attenuatis, demum parum ventricosis, ad 4 mm. latis, extus

longitudinaliter striatis, disco concavo, parum obscuriore, margine vix distincto, involuto; ascis elongato-subclavatis, longe stipitatis, 8-sporis, 132 \times 12; sporidiis distichis, elongatis, basi oblique obtusis, antice acuminatis, cum ciliis sporidii diametrum æquantibus, hyalinis, 3-4-guttulatis, 24 \times 6.

Hab. in caulis *Tanaceti* et aliar. plant. in Rhenogovia et Italia bor. — Ascomata (in forma ital.) cum stipite 1 cm. alta, 2-3 mm. lata. Ascii paraphysati, deorsum longe attenuati, 100-110 \times 7-9; sporidia disticha v. oblique 1-sticha, clavato-fusoidea, 22-24 \times 5-6, deorsum acutiora, utrinque cauda acuta 4 \times 0,5 aucta. — Var. **minor** S. et B. l. c.: ascomatibus dimidio minoribus; ascis 120 \times 8-9; sporidiis similibus, sed 20-24 \times 6-7. In caulis, Fiumicello agri Patavini.

17. **Ciboria vinosa** Berl. et Sacc. *Fung. Ven. Novi* p. 14, t. XIII, 836 f. 10. — Ascomatibus majusculis, firmis, lentis, disco latente vinoso, 2-4 mm. lato, extus ruguloso, glauco-roseo; stipite subtili, 7 mm. alto, fusco; ascis clavatis, longe stipitatis, basi noduloso-incrassatulis, apice rotundatis, paraphysibus filiformibus, ramosis cinctis, 65 \times 7-8; octosporis; sporidiis irregulariter distichis, ovoido-elongatis, utrinque obtusiusculis, muticis, 7-8 \times 2,5-3, hyalinis.

Hab. ad stipitem *Angiopteridis tasmaniana* in Horto Botanico patavino.

18. **Ciboria chlorophana** (Rouss. et De Not.) Sacc. *Peziza chlorophana* Rouss. in De Not. *Rettif. Discom.* pag. 385. — Ascomate coriaceo, breviter stipitato, demum appланato subpeltato, excipulo fibroso; ascis amplis 8-sporis; sporidiis fusoideo-ellipsoideis, utrinque subacutatis, continuis, 1-nucleatis.

Hab. ad ramos *Thujæ Melun Galliæ*. — «An *Chlorosplenium*».

19. **Ciboria luteo-virescens** Rob. et Desm. XIX, Not. n. 85, *Peziza pallido-virescens* Phill. Grev. VII, p. 24. — Patellata dein convexa, firma, luteo- vel pallide virescens, margine integro, stipite tenui deorsum attenuato, flexuoso, plus minus elongato terraue immerso; ascis cylindraceo-clavatis octosporis; sporidiis ellipsoideis, 15 \times 5; paraphysibus filiformibus, obtusis granulosis.

Hab. in petiolis *Aceris* terra infossis in Gallia et Britannia. Ascoma 3-12 mm. lat.; stipes sæpe 2 cm. long., ascoma extus sericeo-tomentellum nec vere glabrum.

20. **Ciboria tabacina** Ell. et Holw. in Arthur Bull. Minn. III, p. 35. — 839 Stipe gracili, plerumque filiformi, leniter sursum ampliato, 0,5-

2,5 cm. longo, e flavescenti-tabacino brunneo, ut ascomatis parte exteriore; ascomate cupulato-patelliformi, 1,5-2,5 mm. diam. sub-tus subruguloso, margine integro, æquali, vix leniter incurvo; disco concavo, ardosiaco; ascis cylindraceis 100-115 \times 9-11; paraphysibus validis flavescentibus, sensim sed leniter superne dilatatis; sporidiis uniseriatis, ellipticis, 9-12 \times 4-5, plerumque grosse uninucleatis, e flavescenti hyalinis.

Hab. in petiolis putrescentibus ad «Vermilion lake» Amer. bor. — A *C. Sydowiana* Rehm distincta imprimis sporidiis angustioribus.

21. **Ciboria Sydowiana** Rehm in Hedwigia 1885, p. 226. — Asco- 840 matibus sparsis, primitus subglobosis, dein in patellam subcyathoidem, subtus longitudinaliter subrugulosam, pallide fuscum, margine subscabro cinctam dilatatis, 3 mm. long., stipitatis, 3-4 mm. diam. sicca margine involuto, fuscidule parenchymatico-contextis; ascis cylindraceis, apice incrassatis, 8-sporis, 110 \times 8; sporidiis 1-stichis, ovatis vel elliptico-clavatis, plerumque subcurvatis, 1-cellularibus nucleis magnis 1-2 (rarissime 3) praeditis, hyalinis, 10-12 \times 5-7; paraphysibus hyalinis, filiformibus, haud vel elliptico-clavatis, 3 μ . crass.

Hab. ad pedunculos foliorum putridorum *Quercus* pedunculatae, Thiergarten prope Berlin (SYDOW). — Teste Phill. Disc. p. 143 ab hac non differt *Phialea renispora* (Ellis).

22. **Ciboria renispora** (Ellis) Sacc. *Helotium renisporum* Ellis in 841 Cooke Disc. U. S. I, p. 299. — Ascomate stipitato, 1-2 mm. lato, convexo, dein plano (sicco concavo) pallide cinnamomeo vel brunneo-flavido; stipite æquali, gracili, deorsum atrato, 3-6 mm. longo; ascis subcylindricis; sporidiis reniformibus, 20 μ . longis.

Hab. in foliis dejectis putridis, New Jersey Amer. bor. — Species britannica in foliis quercinis (Phill. Disc. p. 143) sporidiis 10 \times 5, ascomate fimbriato videtur diversa.

23. **Ciboria nyssogena** (Ellis) Sacc. *Helotium nyssogenum* Ellis Torr. 842 Cl. 1881, p. 73. — Ascomate stipitato, brunneo, concavo, 5-6 mm. vel minus lato, margine pallido, obtuso; stipite 2,5-5 cm. longo, plerumque flexuoso, 1 mm. crasso; ascis cylindricis, 47 \times 3-4; paraphysibus linearibus, obtusiusculis, æqualibus; sporidiis monostichis, ellipsoideis, hyalinis, 10-12 \times 5.

Hab. in nucibus vetustis Nyssæ multifloræ limo infossis et in foliis putridis in uliginosis, New Jersey Amer. bor.

24. **Ciboria Friesii** Pers. M. E. I, p. 277 (*Peziza*), *Peziza ciborioi-* 843

des Fr. Obs. II, p. 307, non Auctorum. — Cupula infundibuliformi, levi obscure rufescente, stipite longissimo capillari, spadiceo.

Hab. ad culmos in uliginosis in Suecia, in foliis *Quercus* in Britannia (?) et ad ramos in Amer. bor. (?). — Gregaria, tenera, sed tenax. Stipes 2-3 cm. fere, subflexuosus; cupula 1 mm. lata, demum subcrenulata.

25. **Ciboria tenella** Karst. *Fragm. mycol. XXII*, p. 124 (ut subsp. *C. Friesii*). — Stipitata, pallescens vel lutescens, siccitate in rufum leniter vergens, plano-infundibuliformis, stipite flexuoso, usque ad 3 cm. longo; ascis subcylindraceis 60-66 \approx 6; sporidiis monostichis, eguttulatis, oblongatis, rectis, 6-8 \approx 2-3.

Hab. in calamis putrescentibus *Caricum* ad Mustiala Fenniæ.

26. **Ciboria tremellosa** Fuck. *Symb. mycol.* p. 312. — Sparsa fasci- 845 culatave, longe stipitata, ad 2-3 cm. alta, diaphana; stipite crasso, distincte ruguloso, fusco, diaphano; cupulis 4 mm. latis, applanatis, demum revolutis, margine non elevato sed paullo obscuriore, disco atro-fusco, levi; ascis sessilibus, oblongis, 8-sporis, 100 \approx 8; sporidiis distichis, oblongo-lanceolatis, inæquilateralibus, minute guttulatis, hyalinis, 16 \approx 4; paraphysibus elongato-clavatis, fuscis, apice obscurioribus.

Hab. in ligno udo putri *Quercus*, rarius pr. Eberbach Rhenogoviæ.

27. **Ciboria macilenta** Karst. *M. F. I*, p. 108 (*Rutstroemia*). — Sparsa, 846 stipitata, pallida; cupula convexa, subtus concava, sat tenui, 1,5 mm. lat., stipite filiformi, æquali, circiter 2 mm. alt.; ascis cylindraceis 110-130 \approx 7-8, obturaculo minutissimo jodo levissime cærulecente; sporidiis monostichis, ellipsoideis vel oblongato-ellipsoideis simplicibus, vulgo eguttulatis, rectis, 11-14 \approx 5-6; paraphysibus circiter 2 μ . cr. apicem versus leniter incrassatis.

Hab. ad terram vasorum floralium, Mustiala Fenniæ. — Ascomatis forma prædistincta. Serius a cl. Auctore ad *Helotium* relata.

28. **Ciboria gracilipes** Karst. *Symb. Myc. Fenn. III*, p. 56 (*Rutströmia* Rev. monogr. p. 126 (*Helot.*)). — Sparsa, stipitata, glabra; cupula concava, sicca convexa, alutacea vel rufescente, 3 mm. lat.; stipite deorsum attenuato, subflexuoso, pallidore, gracillimo, maxima parte terræ immerso, 1 cm. alt.; ascis cylindraceis, apice jodo dilute cærulecentibus, 120 \approx 11-12 (parte sporif.); sporidiis monostichis, ellipsoideis vel sphæroideo-ellipsoideis, guttulis duabus, apicalibus, exiguis præditis, hyalinis, 14-20 \approx 9-11; paraphysibus filiformibus, gracilibus.

Hab. in terra humosa prope Mustiala Fenniæ. — Cum *C. macilenta* comparanda, sed sporidiis majoribus, colore alutaceo etc. mox distincta.

PITYA Fuckel Symb. mycol. p. 317 [1869] (Etym. *pitys* pinus, hinc non *Pithya*). — Ascoma media magnitudine, ceraceum, primo teres, demum explanatum, in stipitem crassum brevem attenuatum, basi albicantè fibrillosa, ceterum subglabrum, margine acuto vix distincto, disco colorato. Asci longissimi, cylindracei, 8-spori. Sporidia monosticha, perfecte globosa, simplicia, subhyalina; paraphyses filiformes, clavatæ. — Autumnales, substantia fere *Helotii*, sed fructificatio *Barlaeæ*.

1. **Pitya vulgaris** Fuck. Symb. p. 317, *Pez. pitya* Pers. ic. et descr. 848 2, p. 43, t. 11, f. 2, Nees Syst. f. 287, Fr. S. M. II, p. 155, *Pez. Leineri* Raben. et Gonn. t. V, f. 6, Weberb. t. III, f. 2. — Cuppula planiuscula extus stipiteque brevi villosis, pallidis, disco aurantio-rubro; ascis longissimis, elongatis, 232×12 , in superiori parte 8-sporis; sporidiis monostichis, globosis, nucleo hyalino, episporio lutescente, crasso, 12μ . diam.

Hab. in caudicibus, ramis acubusque languidis emortuisve Abietis pectinatae nec alibi, in Germania, Gallia, Helvetia, in cortice Sequojae sempervirentis in Amer. bor. — Discus in adultis concaviusculus, rugosus. Gregaria, majuscula, firma, ceraceo-crassiuscula, 5-12 mm. lata et ultra, margine obtuso non prominente, subtus pallidior, demum fere glabra. Variat stipite breviore, disco demum dilute flavo subumbilicato etc.

2. **Pitya Cupressi** (Batsch) Fuck. Symb. p. 317, *P. Cupressi* Batsch 849 Elench. p. 119, *P. cupressina* Fr. Syst. Myc. II, p. 135, De Seyn. Veg. inf. III, t. III, *Fungoides aureum et omnium minimum turbinatum et vix cavum* Mich. gen. p. 207, t. 86, f. 20. — Cupulis carnosо-ceraceis integerrimis flavo-rufescensibus, primo clavæformibus, dein hemisphæricis; stipite brevi crasso; ascis longissimis, cylindraceis, 8-sporis, 68×8 (parte sporifer.); sporidiis in asci superiori parte monostichis, perfecte globosis, $8-10 \mu$. diam., nucleo hyalino, episporio levi dilutissime rubello; paraphysibus reti-clavulatis.

Hab. ad folia Cupressi et Juniperi Sabinæ et virginianæ in Italia, Germania, Belgio, Britannia, Fennia, Suecia; in foliis Sequojae sempervirentis in Amer. bor. — Gregaria, pusilla,

turbinata, aurea, substantia crassiuscula. Stipes primo villo ejusdem indolis ac in *P. vulgari* obductus.

3. **Pitya thujina** Peck 26 Rep. p. 82 (*Helot.*). — Ascomatibus gla- 850
bris, subsessilibus, concavis v. subplanis, aurantiacis, extus paullo pallidioribus; sporidiis globosis, 7-8 μ . diam.

Hab. in ramulis dejectis *Thujæ occidentalis*, Lowville Am.
bor. — *Pityæ cupressinæ* peraffinis.

4. **Pitya fascicularis** (B. et Br.) Sacc. *Helot. fasciculare* B. et Br. 851
Fung. of Ceyl. n. 958. — Fascicularis, obconico-stipitata, pallida;
hymenio convexo; ascis linearibus; sporidiis subglobosis 7-8 μ . diam.

Hab. ad cortices, Peradenya Ceylon.

HELOTIUM Fr. Summa Veg. p. 354 ex p. (1849) [Etym. *helos* clavus]. — Ascoma ceraceum, crassiusculum, disco subplano initio punctiformi dein dilatato, semper aperto, breve crasse stipitatum v. subsessile, glabrum v. pruinatum, plerumque laticolor. Asci fusoidi v. cylindracei, 8-spori, rarius oligospori, filiformi-paraphysati. Sporidia ovoidea, oblonga, fusoidea, teretiuscula, continua, subinde spurie partita, hyalina. — Genus vastum, per formas intermedias ad plura transiens, hinc ægre limitandum et sec. varios auctores varie sumptum. A typo Karstenii [Rev. monogr.] seposui *Phialeam*, *Cyathiculam*, *Pezizellam* et *Pseudohelotium*, sed plurimæ species ambiguæ, quæ sub variis generibus sunt inquirendæ.

Helotium (1-168) extra glutinosa -
A. Stipitata v. substipitata. (1-418)

I. Disco albo, pallido. (1-33)

1. **Helotium fagineum** (Pers.) Fr. Summa p. 356, *Pez. faginea* Pers. 852
M. E. I, p. 296, Fr. Syst. Myc. II, pag. 136, *Elv. fungif.* Scop.
ann. IV, p. 150, t. 2, f. 8. — Minutum; cupula planiuscula, albida;
stipite brevi crasso; ascis ovato-oblongis, subsessilibus, 8-sporis,
60-70 \approx 8; sporidiis subdistichis, ovato-oblongis, utrinque parum
attenuatis rotundatisque, hyalinis, continua, 10-12 \approx 4-5.

Hab. ad pericarpia sicca faginea, locis umbrosis dejecta, copiose in Suecia, Britannia, Germania, Belgio, Gallia. — Gregarium 1-2 mm. latum, glabrum, pallescens.

2. **Helotium Neesii** (Saut.) Sacc. *Pez. Neesii* Saut. Salzb. pag. 9. — 853
Pusillum, ceraceo-aquosum, niveum, ascomate turbinato; stipite brevissimo æquali.

Hab. in pagina inf. filiorum putrescentibus prope Mittersill
Salzburg. — Affine dicitur *H. phascoidi*.

3. **Helotium naviculisperum** Ellis in Cooke Disc. U. S. I, p. 299, et 854 Bull. Torr. 1874, p. 46. — Albidum, dein ochraceum, stipitatum, disco plano-convexo, circ. 1 mm., paullo obscuriore; ascis latis; sporidiis navicularibus, 25 μ . longis, spurie septatis, granulosis.
Hab. in foliis aqua diu immersis, New Jersey Amer. bor. — Stipes 1 mm. long.
4. **Helotium conformatum** Karst. M. F. I, p. 124. — Sparsum v. 855 garium, stipitatum, pallidum; cupula plana, latit. circiter 2 mm.; stipite solido, pallido, deorsum interdum fuscescente, breviusculo; ascis cylindraceo-clavatis, 80-125 \times 6-9, obturaculo jodo obsolete cœrulecente; sporidiis monostichis, oblongatis vel subellipsoideis eguttulatis, 6-12 \times 3,5-6; paraphysibus 1,3 μ . cr. sursum leviter incrassatulis.
Hab. ad folia Alni putrescentia in par. Tammela, Fennia et Somerset East Africæ aust. (MAC OWAN). — Extus *H. epiphyllum* sat simile, internis vero partibus magis cum *H. uliginoso* convenit.
5. **Helotium palustre** Peck 32 Rep. p. 48. — Stipitatum, planum v. 856 leviter convexum, pallidum vel albidum; stipite 6-12 mm. long., basi leniter incrassato; ascis subclavatis; sporidiis oblongis, 10-12 μ . longis.
Hab. ad folia dejecta in uliginosis, Sandlake Amer. bor. — Ascoma siccando atro-rufo-brunneum seu castaneum; stipes proportione ascomatis elongatus.
6. **Helotium chioneum** Fr. Summa pag. 356, Fuck. Symb. App. II, 857 pag. 63, *P. chionea* Fr. Syst. Myc. II, pag. 132. — Ascomatisbus exiguis ceraceo-eburneis, hyalinis albidis, primum obovatis, e basi stipite brevissimo suffulta in cupulam tenuem concaviusculam tenuiter inflexoque marginatam concolorem dilatatis; ascis elongato-clavatis, 8-sporis, 60 \times 6; sporidiis in ascis superiori parte congregatis, cylindraceis, continuis, hyalinis, 8 \times 2; paraphysibus filiformibus, multiguttulatis.
Hab. ad folia pinea et conos humi jacentes passim in Suecia, Germania. — Affine *H. herbarum* sed distincte stipitatum, ascoma regulare, concavum, tenuius.
7. **Helotium abacinum** (Fr.) Karst. M. F. I, p. 132, *P. abacina* Fr. 858 Syst. Myc. II, pag. 139, *Ombrophila abacina* Fr. Summa Veg. pag. 357. — Tremellosso-molle, minutum, aquose luteolum, cupula plano-convexa; stipite brevi crasso; ascis cylindraceo-clavatis, 50-60 \times 4-5, filiformi-paraphysatis; sporidiis fusideo-elongatis, 6-8 \times 1,5-2.

Hab. ad folia dejecta Pinuum in Suecia, Fennia et Gallia. —

Gregarium, 2-3 mm. habitu fere *Sclerotii complanati*. Stipes firmus, scabriuscus, sursum levissime attenuatus, basi saepe obscurior; capitulum pileiforme, pallidius, 2-3 mm. latum, passim irregularare et subtus excavatum. Ascoma, sec. Karst., album, 1 mm. lat. Species Queletii sporidiis 20 μ . long., 4-guttatis aliena.

8. **Helotium tenerimum** Fr. Summa p. 355, *P. tenerima* Fr. Syst. Myc. II, p. 128. — Minimum, punctiforme, albidum, cupula membranacea turbiuata irregulari, stipite brevi.

Hab. ad folia madida Pini silvestris in Suecia. — Nudo oculo puncta minima refert; cupula irregulari. Stipes tenuis, cupulæ diametro vix brevior, cum cupula glaberrimus.

9. **Helotium Huebnerianum** (Rabenh.) Sacc. *Peziza Huebneriana* Rabenb. D. C. Fl. p. 343, n. 2631, *Helotium rufescens* Ficinus Fl. Dresd. II, XXI. — Sparsum v. cæspitosum, albo-lacteum, siccum brunneo-rufum, ad 7-8 mm. altum, convexum deorsum applanatum, stipite dilute flavo, superne attenuato.

Hab. in foliis Glyceriæ fluitantis in agro dresdensi Germaniæ (HUEBNER).

10. **Helotium phormium** Cooke Grev. VIII, p. 63. — Erumpens, palidum, stipitatum; cupulis clavatis, dein plano-concavis, 0-5 mm. lat. difformibus; stipite ampio, deorsum attenuato; ascis clavatis; sporidiis subclavatis, utrinque rotundatis, nucleatis, curvulis 30-35 \times 5.

Hab. in Phormio tenace, Maungaroa Novæ Zelandiæ.

11. **Helotium album** Schum. Saell. 2, p. 412, Fr. Syst. Myc. II, p. 157. — Album, minutum, tenui, capitulo plano-convexo, stipite brevi crassiusculo; ascis elongato-clavatis, stipitatis, 8-sporis, 60 μ . long.; sporidiis ovato-clavatis, 6 \times 4 hyalinis, (FUCKEL).

Hab. in foliis subputridis graminum in Saellandia (SCHUM.), in foliis putridis Airæ cæspitosæ pr. Oestrich Rhenogoviæ. — A *P. culmigena* certe distinguitur forma et substantia firmiore alba. Gregarium, 2 mm. altum. Stipes deorsum angustatus; capitulum primo planum, dein umbonatum, demum subhemisphæricum (ex icode obtuse conicum) margine acuto subrepando.

12. **Helotium cruentatum** Karst. M. F. I, p. 139. — Sparsum, stipitatum, album v. albidum; cupula planiuscula, disco albo vel albido, trito luteo, dein sanguineo v. subaurantiaco, sicco expallente, 0,6 mm. lat.; stipite brevi æquali; ascis cylindraceo-clavatis, 50

≈ 5; sporidiis fusoideo-elongatis, rectis v. subrectis, guttulatis 8-16 ≈ 1-2.

Hab. in culmis Dactylidis glomeratæ emortuis in Mustiala Fenniæ.

13. **Helotium perpusillum** Desm. 10 Not. p. 36. — Sparsum, minutissimum. aquose album, glabrum, siccum rubescens, capitulo hemisphærico; stipite capillari elongato; ascis..... 864

Hab. ad folia arida graminum pr. Caen in Gallia (ROBERGE).

— Ascoma $\frac{1}{3}$ mm. lat; totus fungillus 1 mm. alt. An *Phialea*?

14. **Helotium Sphagnorum** Cr. Finist. p. 47, t. ult. f. 7. — Ascomate pedicellato vel turbinato, albido, glabro, convexo vel subconcavo. 1-2 mm.; ascis majusculis, subcylindricis, octosporis; sporidiis fusoideo-oblängatis, granulosis; paraphysibus filiformibus, apice piriformi-inflatilis. 865

Hab. ad Sphagna in paludibus Finistère Galliæ. — Excipulum prosthymaticum.

15. **Helotium bryogenum** Peck 30 Rep. Stat Mus. Bot. pag. 61. — 866 Minutum, substipitatum, sparsum, madore pallidum vel flavidio-albidum et expansum, siccum livido-rufum v. subviolaceum, concavum tumideque marginatum; sporidiis subfusoideis, sæpe curvulis, 15-18 μ . longis,

Hab. ad muscos e. g. *Hypnum delicatulum*, Maryland Americæ borealis.

16. **Helotium procerum** Karst. M. F. I, p. 130. — Subgregarium, stipitatum; cupula convexa, lutescenti-albida v. albida, lat. circiter 1 mm.; stipite pallido, alt. 4-5 mm.; ascis cylindraceo-clavatis, 60-70 ≈ 7, obturaculo minutissimo, jodo leviter cœrulecente; sporidiis elongatis vel fusoideo-elongatis, subrectis, eguttulatis, 8-16 μ . long.; paraphysibus sat parcis, gracillimis, sursum leviter incrassatulis. 867

Hab. supra muscos sæpe irrigatos in Fennia, ad folia emorta Sphagni acutifolii in silva Cansiglio Ital. bor.

17. **Helotium geogenum** Cooke Grev. VI pag. 65. — Candidum; cupulis obconicis in stipitem abbreviatum productis, hymenio plano-convexo, 2-3 mm. diam.; sporidiis fusiformibus, 25-35 ≈ 5-7; paraphysibus attenuatis, linearibus. 868

Hab. ad terram, fragmentis ligneis verisimiliter adfixum, in silvis pr. Paris Galliæ.

18. **Helotium vaccinum** (Schum.) Fr. Summa pag. 355, *P. vaccinea* 869 Schum. Sael. p. 430, Fr. Syst. Myc. II, p. 126, Dan. t. 1971, f. 1,

P. membranac. bovina Pers. Myc. Eur. p. 299. — Subcæspitosum, pallidum, primo planum, convexiusculum, dein umbilicatum, margine acuto repando, stipite demum elongato.

Hab. in stercore vaccino, locis umbrosis, rarius in Saellandia. — Minus, primo subsessile, dein stipite 4-6 mm. longo, cylindrico, apice dilatato. Cupula 2-4 mm. lata.

19. **Helotium politum** Phill. Disc. p. 155. — Substipitatum, planum v. ⁸⁷⁰ subconcavum, glabrum, album; margine plano obtuso; stipite brevi, sursum in cupulam incrassato; ascis cylindraceo-clavatis; sporidiis ellipsoideis, subacutis, 10-13 \times 4-5.

Hab. in radicibus plantarum adhuc fere infossis prope Shrewsbury Britanniæ. — Ascoma 0,5-2 mm. lat. Usum nitens.

20. **Helotium albumineum** C. et Peck 26 Rep. pag. 81. — Ascomate ⁸⁷¹ molli, orbiculari vel contorto, mox subplano, albido dein pallido, sicco centro depresso, margine elevato; stipite crassiusculo brevissimo, concolore; ascis cylindraceis; sporidiis allantoideis, 7-8 μ . longis.

Hab. ad ramos decorticatos, North Greenbush Amer. bor. — Albumen nucis Cocœs colore et contextu imitatur. Ascoma are-scendo fissum.

21. **Helotium insititium** Karst. M. F. I, p. 124. — Solitarium, raro ⁸⁷² cæspitosum, erumpens, obconicum, basi substipitato-constrictum, pallescens v. testaceo-pallidum, siccum sordide pallescens, margine subcrenulato, disco concaviusculo, pallido, 2 mm. lat.; ascis cylindraceo-clavatis, 95-110 \times 9-10, obturaculo jodo dilute cœrulecente; sporidiis vulgo oblique monostichis, ellipsoideo-oblongatis v. oblongatis, utroque apice aliquantulum angustatis, 1-2-guttulatis, 11-16 \times 4-6; paraphysibus filiformi-clavulatis.

Hab. in ramulis Aceris campestris siccis in Mustiala Fen-niæ. — Habitu nonnihil accedit hæc species ad *H. epiphyllum*, a quo præsertim forma, colore et magnitudine sporidiorum recedit. Omissa, ut videtur, in auctoris Revis. monogr.

22. **Helotium parile** Karst. M. F. I, p. 115. — Sparsum vel gregarium, stipitatum, albidum; cupula concava vel planiuscula, siccata dilute lutea vel subflava, 0,5-0,8 mm. lat.; stipite gracili æquali, 0,5-1 mm. alt.; ascis cylindraceo-clavatis, numerosis, 75-100 \times 5, obturaculo minutissimo vix v. obsoletissime jodo cœrulecente; sporidiis fusoideo-elongatis v. fusoideo-filiformibus, guttulis 4 majusculis fœtis v. spurie tenuiter 3-septatis, rectis v. curvulis, 6-11 \times 1,5-2; paraphysibus raris gracilibus.

Hab. ad ramulos Betulæ vetustos in par. Messuby Fenniæ. — *H. pallescenti* affine, sed cupula adhuc minore, stipite longiori et sporidiis minoribus diversum. — ****languidum*** Karst. M. F. I, p. 115: gregarium, stipitatum, album v. albidum; cupula convexiuscula v. subplana, sicca concaviuscula et lutescente 0,3-0,8 mm. lat., stipite subæquali 0,3 mm., ascis cylindraceo-clavatis, apice jodo non cœrulescentibus, 52-62 \approx 4,5, sporidiis oblique monostichis, fusoideo-oblongatis, 2-guttulatis v. spurie tenuiter uniseptatis, 4-7 \approx 2; paraphysibus gracilibus. Ad lignum et corticem ramulorum Betulæ, Alni et Rubi idæi exsiccatorum in Fennia australi, passim. Sporidia recta; margo interdum tenuissime puberulus v. subtiliter ciliatus. Variat cæspitosum, stipite brevissimo.

23. ***Helotium pallidulum*** Sacc. Fung. Ven. Ser. IV, n. 92, F. ital. 874 fig. 1327. — Cupulis ceraceo-mollibus gregariis, mediocribus, $\frac{3}{4}$ - $1\frac{1}{4}$ mm. alt., patellaribus, breve crassiuscule stipitatis, totis cereo-albidis, disco primitus concavo dein planiusculo, marginato, ex ascis paraphysibusque velutino, circulari seniove repando; ascis cylindraceis basi longiuscule attenuatis, 90-100 \approx 5; paraphysibus filiformibus obvallatis, 8-sporis; sporidiis distichis anguste cylindro-fusoideis, 12-15 \approx $2\frac{1}{2}$, curvulis, utrinque 1-guttulatis, hyalinis.

Hab. in ramis decorticatis putrescentibus fagineis in silva Cansiglio (Treviso) Ital. bor. — Peraffine *Hel. pallescenti* (Pers.) Fr. ex descript. Karst. Myc. Fenn. I, 114, non Fuckelii (sub Niptera pallescenti) Symb. myc. App. I, 46.

24. ***Helotium subsessile*** Schum. Saell. II, pag. 415, Pers. M. E. I, 875 pag. 345, *Pez. helotoides* Fr. Syst. Myc. II, pag. 135, FI. Dan. t. 1855, f. 3. — Pusillum, pallidum; cupula plano-convexa; disco dilute umbrino vel subochraceo, convexo, stipite brevissimo, basi umbrino-badio; ascis clavatis; sporidiis oblongo-fusoideis, 2-pluri-nucleatis.

Hab. in ramis dejectis subputridis in Saellandia et Britannia. — Subcæspitosum, stipite sursum dilatato in cupulam abeunte, cupula 1 mm. haud lata. Caro concolor.

25. ***Helotium nigripes*** (Pers.) Fr. Summa p. 356, *Peziza* Pers. Syn. p. 661, Fr. Syst. Myc. II, p. 132. — Dilute pallens glabrum; cupula plano-concava marginata, stipite longiusculo nigrescente; sporidiis oblongis, 5 \approx 1,2.

Hab. ad truncos et folia putrida in Suecia et Dania etiam in America boreali, Cuba (var. *alba*) et Tasmania. — β . ***cellare*** A. S. Lus. p. 132, cupula plana subdifformi sordide lutescente. — γ . ***sub-***

tile Fr. l. c., cupula minima, plana, flava, stipite æquali nigro, 1 mm. alt. In conis abiegnis in Suecia.

26. **Helotium miserum** B. et C. Cub. Fung. n. 698. — Minutum, al. 877 bidum, claviforme; stipite brevi; hymenio subgloboso; ascis clavatis; sporidiis subclavatis, endochromate demum quadripartito, 12 μ . longis.

Hab. in corticibus muscosis in ins. Cuba.

27. **Helotium imberbe** (Bull.) Fr. Summa p. 356, *P. imberbis* Bull. 878 Champ. p. 245, t. 467, fig. 2, Fr. Syst. Myc. II, p. 136, *P. nivea* Batsch. El. pag. 117, f. 59 (forma sessilis). — Carnoso-ceraceum, glabrum, album; cupulis plano-concavis, subflexuosis; stipite papillæformi; ascis anguste elongato-clavatis, longe stipitatis, 8-sporis, 76 \times 7; sporidiis cylindraceis, rectis, 8 \times 2, hyalinis.

Hab. ad ligna et ramos salicinos, alneos in Gallia, Germania, Britannia. — Habitus *H. lenticularis* sed totum album, demum planum etiam irregulare. De Candolle Fl. Fr. II, p. 81, *H. imberbis* varietatem cinereum etiam notat.

28. **Helotium pallescens** (Pers.) Fr. Summa Veg. p. 355, *P. lenticularis* Hoffm. Fl. germ. Crypt. t. 13 nec Bull., *P. pallescens* Pers. Obs. 35, Syn. 664, Myc. Eur. p. 294, Fr. Syst. Myc. II, p. 132. — Confertum, glabrum, dilute flavum v. albido-pallescens; cupula concava; stipite brevi crassiusculo pallido; ascis cylindraceo-clavatis; sporidiis ellipsoideis v. oblongis, 8-11 \times 2-3, 2-guttatis simulateque 1-septatis, hyalinis.

Hab. ad truncos vetustos, sed rarius in Suecia, Germania, Fennia, Britannia, Belgio, Amer. bor., Ceylon, Tasmania, Nova Zealandia. — Asci, sec. Fuck., 64 \times 8; sporidia 8 \times 4. Revera ab *H. citrino* differt, sed characteres difficiles. Minus, substantia tenuior, discus semper concavus, margo subtumidus regularis, stipes brevior; variat tamen et elongatum (*P. elongata* Schum. Saell. p. 429, h. 1.).

29. **Helotium hypocrita** Hazsl. Köggomb. p. 4, t. I, fig. 1. — Gregarium, majuscum, 1 cm. altum, stipitatum, album; stipite cylindrico, basi subbulboso, albo, basi griseo-fuligineo; ascomate plano-infundibuliformi, extus striatulo, margine inflexo, disco subumbilicato, griseo-fuligineo; ascis cylindraceis noduloso-substipitatis, obtusis; sporidiis octonis, ovideis, continuis, initio quaternatim plasmati obvolutis.

Hab. ad ligna pr. Eperies Hungariæ. — Cfr. Hazsl. Einig. Disc. p. 15, ubi ad *Pez. Tubam* Bolt. refertur.

30 **Helotium antarcticum** Speg. F. Fueg. n. 326. — Sparsum v. laxe 881 gregarium, turbinato-calyciforme, 2-2,5 mm. diam., firmum, tenacellum, crassiusculum, extus ex albo fulvescens, intus pallescens, margine acuto integerrimo glabrum, basi in pedicellum breviusculum crassiusculum cylindraceo-conoideum, 1-1,5 mm. long., 300-500 μ . crass. sæpe basi connato-confluens candidum productum; ascis cylindraceis, antice obtuse rotundatis, postice attenuato-pedicellatis, 180-200 \approx 8 octosporis, dense paraphysatis, iodi ope immutatis; sporidiis recte v. oblique monostichis, ellipticis v. subfusoides, utrinque acutatis, 14-15 \approx 6, non vel biguttulatis, hyalinis, continuis.

Hab. ad truncos decorticatos putrescentes *Fagi betuloidis* in silvis in Staten island Fuegiæ. — Species firmula, tenacella, habitu fere *Ciboriae firmæ*, a genere nonnihil desciscens.

31. **Helotium aciculare** (Bull.) Pers. Syn. pag. 677, Pat. Tab. anal. 882 f. 288, *Pez. acicularis* Fr. Syst. Myc. II, p. 156, *Helotium agariciforme* DC. gall. II. 75, *Leotia acicularis* Pers. obs. II, 20, t. 5, fig. 1, *Helvella agariciformis* Bolt. Sowerb., *Helvella acicularis* Bull. t. 473, f. 1. — Ascomatibus gregariis glabris candidis, dein pallentibus v. rufescens, stipite elongato æquali tenui discum primum utrinque planum dein convexum, subtus marginibus reflexis concaviusculum sustinentibus; ascis clavatis, octosporis; sporidiis distichis breve fusoides, continuis, 10-12 \approx 3-4 (ex icone Pat.); paraphysibus bacillari-clavulatis.

Hab. in truncis cavis *Quercus*, rarius *Robinia*, in Suecia, Britannia, Germania, Gallia, Belgio, Amer. bor. — Gregarium tenax diu persistens, colore candido dein pallido demum cinerescente v. rufescente. Stipes 10 mm. longus, tenuis, firmus, passim curvatus et ramosus; capitulum primo utrinque planum v. centro leviter excavatum, demum vero convexum, subtus concavum, 2-4 mm. latum. Sec. auctores nonnullos sistit formam tenellam *Cudoniellæ Queletii*. — β . **abietinum** Fr. l. c. Alb. et Schw. l. c. *Helotium elongatum* Schum., ascoin. minoribus pallidioribus.

32. **Helotium herbarum** (Pers.) Fr. Summa pag. 356, Karst. M. F. I, 883 pag. 117, *Pez. herbarum* Pers. Syn. pag. 664, Fr. Syst. Myc. II, p. 136. — Carnoso-ceraceum, glabrum, album; cupulis plano-convexus adpressis; stipite brevissimo; ascis elongato-clavatis, stipitatis, 8-sporis, 60-75 \approx 7-8; sporidiis subdistichis, oblongis, parum curvatis, hyalinis, continuis, 10-14 \approx 2, 2-3-guttatis, subinde simulate septatis.

Hab. ad caules Urticæ dioicæ, Solidaginum, Brassicæ, Scrophulariæ, Senecionum in Suecia, Fennia, Britannia, Germania, Gallia, Italia, Belgio, Amer. bor., Fuegia. — Huc forte spectat *Peziza Ebuli* Ces. in Klotzsch H. M. n. 1424. — Stipes tenuis v. subnullus; cupula 2 mm. lata et ultra, integerrima, demum subinde rufescens, saepius immarginata.

33. **Helotium herbicolum** Karst. M. F. I, p. 118, Rehm Ascom. Lojk. 884
n. 15, *Pez. Rubi* Lasch var. *herbicola* Rab. Fung. eur. ed. II, 218. — *H. herbarum* affine, sed differt cupula tenuiore, latiore; margine saepe repando-flexuoso, colore lætiore (sicco fulvescente testaceo); stipite verruciformi; sporidiis cylindraceis, guttulis 2-6 præditis v. spurie tenuiter 1-3-septatis.

Hab. ad caules herbarum variarum, præcipue Hyperici et Umbelliferarum in Fennia, Germania, Hungaria. — Formis intermediis cum *H. herbarum* conjunctum. Habitu quoque interdum refert *H. discreti* formas brevissime stipitatas, sed colore cupulæ speridiisque majoribus mox distinctum. Sec. Rehm l. c. asci filiformi-paraphysati; sporidia fusoideo-elongata, 9-12 \approx 1,5, 4-guttulata.

II. Disco rubente, roseo, incarnato, lilacino. (34-43)

34. **Helotium kermesinum** Fr. Summa p. 356, Karst. M. F. I, p. 123, 885
Pez. Girgensohni Dietr. Arch. Liv. Ehstl. 1856, p. 368, *Pez. geminella* Nyl. Obs. Pez. p. 47. — Ferrugineo-rubrum v. subcinabarinum; cupula plana, marginata, circ. 1 mm. lat.; stipite brevi crassulo v. demum brevissimo; ascis 60 \approx 7, obturaculo apicali jodo leviter cœruleacente; sporidiis binis, oblongatis v. ellipsoideis, 11-12 \approx 4.

Hab. supra folia Betulæ putrescentia ad Onegam lacum Fenniæ et in Suecia et Germania.

35. **Helotium fraternum** Peck Bull. N. Y. St. Mus. n. 2, p. 21, t. I, 886
f. 12-14. — Ascomate minuto, stipitato, 1-2 mm. lato; disco plano v. concaviusculo pallido v. flavo-rubescente, demum magis concavo et saturatus colorato (in sicco); stipite 1-2 mm. longo; ascis clavatis vel cylindraceis, 70-100 \approx 10-12; sporidiis confertis vel biseriatis, subcylindraceis, 16-20 \approx 4-5; paraphysibus numerosis, filiformi-clavulatis.

Hab. in petiolis et nervo medio foliorum dejectorum Aceris saccharini Adirondack Mount. Amer. bor.

36. **Helotium rubescens** Cr. Fl. Finist. p. 49, nec. alior. — Ascomate 887

stipitato, 2-4 mm., plano v. subtus concavo, supra convexo, rubescente v. brunneolo; ascis cylindricis, octosporis; sporidiis oblongis, utrinque granulosis; paraphysibus filiformibus.

Hab. ad radices graminum humi fusas Finistère Galliæ. — Ascomatis excipulum cellulis ovoideo-elongatis, seriatis, biformibus contextum.

37. **Helotium carneum** Fuck. Symb. App. II, p. 63 vix Fr. — Cupulis 888 sparsis, carnosis, obconicis, in stipitem brevem attenuatis, usque ad 2 mm. latis, glabris, disco concavo, carneo-flavo, marginatis extus dilute carneis, stipite albescente; ascis clavatis, 8-sporis, 48 \approx 6; sporidiis distichis, cylindraceo-fusiformibus, rectis, continuis, hyalinis, 8 \approx 2.

Hab. in foliis putridis Scirpi silvatici pr. Mappen Germaniæ.

38. **Helotium alabastrinum** Fr. Syst. Myc. II, p. 125 (*Pez.*). — Carnosum, albo-incarnatum; cupula e concavo-explanata integerrima; stipite brevissimo, papillæformi, nigrescente.

Hab. ad culmos locis humidis in Suecia. — An *Phialea?* Gregarium, pusillum, vix 3 mm. latum, carnosso-ceraceum ob hymenium crassum ascosque amplos, subtus convexum. Stipes e sordibus forsan nigricans.

39. **Helotium polytricum** Cr. Fl. Finist. pag. 48. — Ascomate sub- 890 pedicellato, roseo, glabro, initio infundibuliformi, dein expanso, 1 mm.; ascis suboblongis, octosporis; sporidiis oblongis, biguttulatis; paraphysibus filiformibus.

Hab. in foliis vivis Polytrichi communis Finistère Galliæ. — Cellulæ excipuli sphæricæ, granulis immixtæ.

40. **Helotium fimetarium** Pers. Syn. p. 678, Fr. Summa p. 355, Fuck. 891 Symb. p. 313, *Leotia fimetaria* Pers. Obs. Myc. 2, pag. 21, t. 5, f. 4-5, *P. fimetaria* Fr. Syst. Myc. II, p. 157. — Parvum, amœne rubrum, glabrum, capitulo e conico explanato subangulato, stipite æquali; ascis....; conidiis (ex Fuckel) numerosis, ovoides, 1-guttulatis, hyalinis, 6 \approx 4.

Hab. ad fimum vaccinum ovinum etc. pluviis maceratum in Suecia, Germania, Belgio. — Gregarium, perpusillum, sed sat persistens. A reliquis differt forma non juniore, sed ætate planiuscula. — Var. **americana** Schw. Carol. n. 1273, statura majore. In fimo catorum in Carolina.

41. **Helotium claviculare** Wallr. Crypt. Fr. n. 2537 (*Peziza*). — Ascomatibus gregariis, teneris fragilibus, erectis, glabris, stipite e basi

tenuata sensim incrassato pallido, discum convexum pallide roseum, margine dilutiore acuto reclinato cinctum elevantibus.

Hab. ad simum leporinum pluviis madafactum in Thuringia.

42. **Helotium Phillipsii** Sacc. *Lachnella nuda* Phill. Disc. pag. 247. 893

— Gregarium, planum v. convexum, stipitatum, ceraceum, carneum, glabrum; disco concolore undulato; stipite longo, flexuoso, in ascoma abeunte; ascis cylindraceo-clavatis, 50 \times 4; sporidiis fusoideis v. oblongo-ellipsoideis, hyalinis, 5-10 \times 2-3; paraphysibus fusoideis granulosis, ascos superantibus, 70 μ . longis.

Hab. ad terram muscosam, in pinetis, Forres Bri tanniæ. — Ascoma 2 mm. lat., 4 mm. alt. Ascoma *Helotii* sed paraphyses *Lachnella*.

43. **Helotium albo-lilacinum** Pat. Tab. an. Fung. pag. 128, fig. 289, 894

Quél. XIII Suppl. t. VIII, f. 23. — Exiguum, 1-2 mm. lat. stipitatum, violaceo-lilacinum, disco initio concavo, dein plano, tenui, membranaceo, albo v. lilacino, margine integro; stipite glabro, basi attenuato; ascis clavatis octosporis; sporidiis minutis, oblongo-ellipsoideis, hyalinis, 2-guttatis, 10-15 \times 4; paraphysibus bacillaribus.

Hab. in caulis emortuis Sambuci Eboli, quos colore rubro inficit. Stipes 0,2-0,4 mm. long.

III. Disco vidente, subolivaceo (Cfr. *Chlorosplenium*). (44-46)

44. **Helotium flavo-virens** (Pers.) Fr. Summa p. 355, *P. flavo-virens* 895

Pers. Myc. Eur. p. 323, Fr. Syst. Myc. II, pag. 139. — Ceraceo-molle, minutum, flavo-virens, primo subumbilicatum, demum convexo-planum; stipite crasso brevissimo.

Hab. in ligno putrido locis uidis prostrato in Vogesia (MOUGEOT) et Smolandia (Fr.). — Historia fere *H. conigeni*. Totum glabrum, vix marginatum, demum applanatum, 2 mm. latum. Color in siccis rufescit.

45. **Helotium agaricinum** Berk. Engl. Fl. V, p. 207 (*Peziza*), *Helotium* 896

Phillips Disc. pag. 170. — Firmum, atro-virens, convexum, subinæquale; margine revoluto; stipite crassiusculo, obconico; ascis longis flexuosis, obtusis.

Hab. in ligno emortuo, Appin Britanniæ. — Ascoma 4-5 mm. lat.

46. **Helotium subolivaceum** Karst. Ascom. add. p. 449. — Sparsum, 897

stipitatum, glabrum, olivaceo-pallidum; cupula concava, 1 mm. lat.; stipite filiformi, deorsum pedentem attenuato, diametrum cupule excedente; ascis confertis cylindraceis, infra pedentem attenuatis obturaculo jodo dilute cœrulescente, 110-140 \times 10-11; sporidiis ob-

lique monostichis, fusoideo-oblongatis, simplicibus, 1-2-guttulatis, rectis, hyalinis, 12-16 \approx 5-6; paraphysibus parcis, gracillimis.

Hab. ad petiolos (*Aceris?*) putridos in Mustiala Fenniæ.

IV. Disco flavidō, luteo, aurantio. [47-73]

47. **Helotium conigenum** (Pers.) Fr. Summa pag. 355, Fuck. Symb. 898 p. 316, *Pez. conigena* Pers. Syn. p. 634, Fr. *Pezizella conorum* Rehm in Asc. n. 415. — Gregarium, sæpe confluens, brevissime crasse stipitatum, demum sessile, primitus globulosum, siccum concavum, margine ruguloso pallidiore, disco flaveolo, humidum explanatum, orbiculare, stramineum, pellucidum, planum v. subconvexum, 1 mm. diam.; ascis clavatis, stipitatis, apice vix incrassatis, 8-sporis, 40 \approx 6; sporidiis elongato-ovatis vel ellipticis, rectis vel subcurvatis, hyalinis, 6-8 \approx 1,5-2, distichis in ascis superiori parte; paraphysibus filiformibus crassis, sæpe sensim clavulatis.

Hab. ad conos Abietum et Pinorum in Suecia, Britannia, Germania, Gallia, Belgio, Fennia. — Variat lividum incarnatum. Ad *Pezizellam* vergit.

48. **Helotium fastidiosum** Peck 27 Rep. St. Mus. N. Y. p. 107. — Asco- 899 matibus minutis, convexis v. planis, stipitatis, pallide flavis v. albidis; stipite gracili circ. diametrum ascomatis longitudine æquante, brunneo v. flavo basique brunnea; ascis anguste clavatis; sporidiis confertis v. biseriatis, elongatis, subclavatis, multinucleatis, 25 \approx 5, subinde leniter curvulis.

Hab. in petiolis costisque foliorum deciduorum Forestburgh Amer. bor.

49. **Helotium sulphurellum** Ell. et Ever. in Bull. Torr. Club. 1883, 900 p. 98. — Ascomate læte citrino vel subsulphureo, viridulo-tineto; farinoso; stipite firmo, basi brunneo, sursum sensim incrassato, 3-7 mm. longo, disco (in siccis) pallide cinnamomeo-brunneo, concavo vel subplano, subinde umbilicato, 2-4 mm. lato; margine subacuto plano; ascis cylindricis, 75 \approx 7-8; paraphysibus filiformi-clavulatis, flavidis; sporidiis monostichis, imbricatis, naviculari-ellipticis, binucleatis, subhyalinis v. flavellis, 10-12 \approx 3,5-5.

Hab. in petiolis dejectis Fraxini americanæ West-Cester Pa. Amer. bor.

50. **Helotium Buccinula** Speg. Fung. Fueg. n. 327. — Sparsum v. laxe 901 gregarium, epiphyllum, ceraceum, opacum, discoideum, 500-800 μ . diam., superne appланatum v. convexulum flavescens, inferne applato-subturbanatum, glabrum album, obtuse crasseque marginatum;

pedicello longiusculo gracili, 1-1,5 mm. long., 100-150 μ . crass., concolori glabro non v. vix striatulo; ascis clavulatis, sursum breviter cuneato-rotundatis apiceque subtruncatis, late perviis, deorsum in pedicellum mediocrem attenuatis, 90-100 \approx 8-12; paraphysibus paucis filiformibus obvallatis, octosporis; sporidiis oblique monostichis clavulatis sursum obtuse rotundatis, postice attenuato-acutatis, 25 \approx 5, hyalinis grosse 2-3-guttulatis (an demum biseptatis?).

Hab. ad folia dejecta Berberidis ilicifoliae in dumetis prope Shammacus Fuegiæ. — An *Phialea*?

51. **Helotium serotinum** (Pers.) Fr. Summa pag. 355, Sacc. F. ital. 902 f. 1345, *Helv. aurea* Bolt. fung. t. 98, sec. Nees, *P. serotina* Pers. syn. p. 661, Fr. Syst. Myc. II, p. 119, *Hel. umbelliformis* Pers. M. E. I, p. 346, *Hymenoscypha* Phill. Disc. pag. 125. — Late flavum, cupula plano-convexa, subtenui, stipite brevi firmo crassiusculo; ascis elongatis, stipitatis, 8-sporis, 128 \approx 6; sporidiis elongato-distichis, clavatis, curvatis, hyalinis, continuis, 20-22 \approx 4 (in clavula).

Hab. in umbrosissimis uidis (sæpe in ipsa pæne aqua) ad folia ramosque putridos, altero anno abundans altero parce subsparsum, in Germania, Britannia, Belgio, Italia, Amer. bor. — Gregarium, confertum. Stipes 2-4 mm. longus, concolor; cupula numquam clausa 2-4 mm. lata. — Var. **obesum** Bres. in Hedw. 1885: cupula e subgloboso inverse-conica, medio primitus convexa dein plana tandem parum impressa, 2-7 mm. lata, 2-4 mm. alta, disco orbiculari, ceterum formæ typicæ; sporidiis oblongis, subfusiformibus, 14-16 \approx 3. Ad ligna et folia quercina adhuc solida aut putrida, et juxta ea in terra gregatim in silva Crni gaj et Crkvenac Slavoniæ.

52. **Helotium confluens** Schw. Syn. p. 176 (*Peziza*). — Majuscum, 903 sæpe plagis, 2-3 cm. lat. et ultra confluens (ita ut cupulæ confluentes quasi membranam efficiunt); stipitibus crassiusculis, brevibus, albescientibus, cum cupulis ad basim ceterum vivide citrinis, subtus pallidioribus.

Hab. ad ramos, ligna, Bethlehem Amer. bor. — Affine *H. citrino* sed multo majus.

53. **Helotium infaciens** (Ces.) De-Not. Comm. I, p. 378. — Subcutaneum, diu tectum, demum per rimas longitudinales erumpens, læte flavum e stipite plus minus compresso dilutiore in ascoma obtuse marginatum dilatatum; ascis perfecte cylindricis, 100 \approx 12; sporidiis fusoideo-ovoideis obtusis, 2-guttatis, 20-22 μ . long.

Hab. ad ramos ramulosque *Cytisi alpini* emortuos in Italia boreali. — *Sparsum v. cæspitosum*, *Hel. citrino* affine.

54. **Helotium lutescens** (Hedw.) Fr. Summa p. 355, Gill. Disc. p. 157, 905
Phialea lutescens (Hedw.), *Octospora lutescens* Hedw. Musc. p. 30,
 t. 9, f. 3, *Peziza* A. S. p. 334, Fl. Dan. t. 1440, f. 1, Fr. Syst.
 Myc. II, pag. 120. — *Lutescens*, minus; cupula hypocrateriformi,
 orbiculari; stipite brevi tenui subcylindrico; ascis tereti-clavatis;
 sporidiis oblongis vel oblongo-ellipsoideis, 12-15 \approx 4; paraphysibus
 filiformibus.

Hab. ad ramos dejectos ab iegnos in Germania, Suecia, Britan-
 nia, Gallia, Sibiria asiatica, etiam in America, sed rarius, autumno.
 — Ætate juniore pro more cylindricum, dein apice turbinatum, ma-
 gis magisque ampliatum.

55. **Helotium Phiala** (Vahl) Fr. Summa p. 355, *Pez. Phiala* Vahl Fl. 906
 Dan. t. 1078, fig. 2, Schum. Saell. p. 420. — Cupula cyathiformi
 integerrima sordide flavescente, disco spadiceo, stipite brevi levi
 crassiusculo.

Hab. in ramis putridis *Alni* et *Betulæ* in Scandinavia, sed
 rarum. — Optima species, in hoc genere majuscula, solitaria, fir-
 mula. Stipes 3-4 mm. longus, sursum incrassatus, subincurvus,
 basi subinde villosa; cupula dilatata, 4-7 mm. lata, utrinque levis,
 disco obscuriore.

56. **Helotium hyalopus** Fuck. Symb. App. II, p. 63. — Cupulis cera- 907
 ceis, sparsis, distinete stipitatis, primo excavatis, marginatis, demum
 planis subconvexisque, patellæformibus, pallide-flavis, glabris, 2 mm.
 lat., stipite plerumque curvato, hyalino-diaphano, demum pallide-
 flavo; ascis subclavatis, sessilibus, 8-sporis, 126 \approx 18; sporidiis ple-
 rumque in asci superiori parte conglobatis, oblongo-fusiformibus,
 continuis, subinæquilateralibus, biguttulatis, hyalinis, 16 \approx 6; para-
 physibus numerosis.

Hab. in sarmentis emortuis *Vitis viniferæ* pr. Budenheim
 Rhenogoviæ.

57. **Helotium hongkongense** (B. et C.) Sacc. *Pez. hongkongensis* B. 908
 et C. North Pac. Exp. n. 154. — Breviter stipitatum; cupula he-
 misphærica subtus lutea, margine incurvo; hymenio convexulo vel
 plano, aurantiaco.

Hab. in ramis emortuis Hong-Kong Sinarum.

58. **Helotium crocinum** B. et C. Cub. Fung. n. 701. — Crocatum, ob- 909
 conicum; stipite brevi crasso; cupula crateriformi, margine inflexo.

Hab. ad ramos emortuos in insula Cuba. — *Phialeam crocata* Mont. in mentem revocat.

59. **Helotium citrinum** (Hedw.) Fr. Summa p. 355, *Octospora* Hedw. 910
Musc. II, t. 8, f. B, *Peziza lutea*, *parva* etc. Ray. syn. 3, p. 18,
tab. 24, f. 4, *P. cyathoides* Linn. Suec. n. 1273! With. arr. IV,
pag. 349, *P. citrina* Batsch Cont. 2, f. 218, Pers. syn. pag. 663,
Schum. Saell. 2, p. 430, Dec. fr. 6, p. 22, *P. aurea* Sowerb. t. 150.
— Confertum, citrinum; cupulis plano-concavis cum stipite brevi
crasso pallidiore obconicis; ascis clavatis 90-100 ± 8-9; sporidiis
oblongis, obtusis, hyalinis, duobus nucleis magnis instructis, 10-12 ±
4, monostichis, superne sæpe distichis; paraphysibus filiformibus
apice haud clavatis.

Hab. ad truncoſ et ramos putridos Fagi, Carpi ni, Quercus
Alni etc. in Suecia, Gallia, Britannia, Germania, Belgio, Fennia,
Italia, Tasmania, Kamtschatka, Amer. bor., Nova Zelandia, Fuegia.
— Substantia carnoſo-ceracea, crassiuscula, firma contextu prosen-
chymatico. Cupulæ margo prominens, subinde flexuosus. E situ va-
riat stipite elongato et brevissimo, tantum papillæformi, semper al-
bido; hæc forma vulgatissima in strobilis abiegnis et certe *P. flava*
Schum. Saell. p. 418, Pers. Myc. Eur. p. 294. — Var. **Phalaridis**
(Lib.) Speg. et Roum. Rel. Lib. I, n. 660, *Pez. Phalaridis* Lib.:
a typo recedit cupulis umbrinis, disco albo-rufescente. Fructificatio
perfecte congruit. Ad culmos aridos Phalaridis arundinaceæ
prope Malmedy in Arduennis.

60. **Helotium Buccina** (Pers.) Fr. Summa pag. 355, Mont. Chil. VII, 911
p. 409, *P. Buccina* Pers. syn. p. 659, Myc. Eur. p. 279, Fr. Syst.
Myc. II, p. 129, Alb. et S. p. 331. — Majusculum, infundibulifor-
me, obscure flavum; stipite incrassato, striato, subincurvo.

Hab. rarius ad ligna et ramulos pineos in Suecia, Germania,
Britannia, Chile, etiam in Carolina. — Pro diversa ætate forma
varia: ætate juniore incurvo-clavata, adulta campanulata. Color in
junioribus flavus, in adultis saturate aureus. Stipes profunde et pa-
rallele striatus, imo in speciminiibus nonnullis subundulatim sul-
cato-costatus; cupulæ etiam basis interior quandoque striato-plica-
ta, subcæspitosæ etc. A. S. l. c., *Pez. Buccina* pl. auct. est *Guepi-
nia merulina*; sed Doct. Quélet ita describit *Phialeam Buccinam*
(Pers.) Quél. 11 Suppl. p. 21: cyathiformis, pallide citrina, stipite
longo, 5-7 mm. basi fulvo; cupula tenui, 5 mm. lat.; sporidiis vir-
guliformibus, 30 μ. long., 4-5-guttulatis, hyalinis. In ramulis fagi-

neis in Vogesis (R. FERRY). Sec. Mont. l. c. asci 280 \pm 10; sporidia ellipsoidea, 20-30 \pm 6-8.

61. **Helotium aureum** Pers. syn. p. 678, *Pez. aurea* Fr. Syst. Myc. II. p. 156. — Obscure flavum, ascomate lenticulare; glabrum, stipite tenui basi tomentoso. 912

Hab. ad ligna decorticata pinea putrida et in cortice duriore in silvis, in Suecia, Germania, Sibiria. — Gregarium, tenue. Stipes teres, æqualis, subinde ramosus, basi tomentulo delicato niveo cinctus; ascoma leve, 2-3 mm. latum; asci maturi non visi (Fuck.).

62. **Helotium pineum** (Bonord.) Sacc. *Sarea pinea* Bonord. Handb. p. 198. — Ascomate breve stipitato, convexo, flavo, dein brunneo, sub lente puberulo; sporidiis ovato-oblongis curvulis, guttulatis; paraphysibus ramulosis, capitulatis. 913

Hab. in pariete interiore corticis Pinorum in Guestphalia.

63. **Helotium claro-flavum** (Grev.) Berk. Outl. p. 372, Phill. Disc. p. 165, t. 5, f. 31, *Pez. Grev. Fl. Edin.* p. 424. — Breve stipitatum v. sessile, subplanum, pallide flavum, glabrum, margine emergente, obtuso, sublobato; ascis cylindraceis, octosporis; sporidiis oblongis v. oblongo-ellipsoideis, 7-10 \pm 2-3; paraphysibus filiformibus tenuibus. 914

Hab. in ligno et ramis dejectis prope Edinburgh et Wrekin Britanniæ et in Belgio, Australia et Africa austr. — Ascoma vix 1 mm. lat.

64. **Helotium flavum** Klotzsch in Phill. Disc. p. 156. — Gregarium, stipitatum, initio subpiriforme, dein expansum, disco læte flavo, extus pallidius; stipite glabro, brevi, basi acuto; ascis tereti-clavatis; sporidiis ellipsoideis, raro subclavatis, 10 \pm 4-5; paraphysibus filiformibus. 915

Hab. in ligno decorticato in Britannia. — Ascoma 0,5-1 mm. lat.

65. **Helotium lenticulare** (Bull.) Fr. Summa pag. 356, Sacc. F. ital. fig. 1347, *P. flava* Willd. Ber. n. 1175, *P. lenticularis* Bull. Ch. t. 300, f. a, c, Fr. Syst. Myc. II p. 133, *Octosp. citrina* Fl. Dan. t. 1294, f. 1, *Helotium nigripes* Schum. Sael. p. 412, *P. citrina** *lentic.* Pers. l. c. p. 293, *P. aurea* Sow. t. 156. — Convexum, adpressum, firmum, luteum; stipite papillæformi nigrescente; ascis tereti-clavatis; sporidiis oblongis, oblongo-ellipsoideis vel clavatis, 10-15 \pm 3-4. 916

Hab. ad truncos vetustos fagineos etc. abunde in Suecia, Dania, Germania, Gallia, Italia, Britannia, Ceylon, etiam in America boreali — Subsessile, gregarium, glabrum, 2 mm. latum, constanter læte flavum, nitidum, primo cavum, dein convexum.

66. **Helotium sulphuratum** (Schum.) Phill. Disc. p. 161, *Pez. sulphurata* Schum. Fl. Dan. t. 1912. f. 2, an Friesii (Cfr. *Humaria*). — Sparsum, turbinatum, stipitatum vel subsessile, initio orbiculare, concavum, dein expansum, margine integro, e sulphureo citrinum, extus margineque pallidius; ascis tereti-clavatis, octosporis; sporidiis fusoideis rectis curvulis, 2-3-guttulatis, 12-17 \approx 3-4,5; paraphysibus filiformibus, tenuibus, subinde ramulosis.

Hab. ad lignum et folia pinea terra subinfossa in Dania et Britannia.

67. **Helotium gratum** (Berk.) Cooke Austral. Fungi pag. 51, *Peziza grata* Berk. Fl. Tasm. p. 275. — Cupula plana, hyalina, marginata, breviter stipitata subaurantiaca, 1 mm. lat.; stipite pallidiori cylindrico quandoque compresso; sporidiis fusoideis, 10 μ . long.

Hab. ad ligna emortua in Tasmania (ARCHER). — Affine dicitur *Sarcoscyphæ Colensoi* Berk., an ergo hujus generis?

68. **Helotium repandum** Phill. Disc. pag. 161. — Breve stipitatum, planum, repandum, tenue, pallide flavum, margine sublobatum; stipite brevissimo, basi attenuato; ascis cylindraceo-clavatis; sporidiis cylindraceis vel oblongo-ellipticis, 7-10 \approx 2; paraphysibus filiformibus.

Hab. in caulis emortuis Spirææ Ulmariae in uliginosis Scotiæ. — Ascoma 0,5-1 mm. lat.

69. **Helotium erythropus** (Saut.) Sacc. *Pez. erythropus* Saut. Pilz. Salzb. pag. 9. — Ascomate carnosoceraceo, concavo, albido, disco aurantiaco; stipite brevi, crasso roseo.

Hab. in caulis exsiccatis Angelicæ silvestris prope Mittersill Salzburg. — Affine *H. carneo*.

70. **Helotium? brassicæcolum** (Schw.) Sacc. *Sarea brassicæcola* Schw. Syn. pag. 178. — Late gregarium, minutum; ascomate initio convexo, rufo-aurantio, demum disco applana to marginato; stipite basi incrassato albo-pulverulento.

Hab. in caulis vetustis induratis Brassicarum inter fibras, Bethlehem. Amer bor. — Genus *Sarea* sec. Schw. sistit præterea *Helot. aureum*, *aciculare*, *fimeti*, *subtile* et *Pityam*.

71. **Helotium spicarum** Rehm Ascom. n. 158: cfr. Cooke in Grevillea IV, p. 41, t. 51, f. 258. — Plerumque sparsum, turbinatum, breviter crasse stipitatum c. 1. mm. alt., siccum fuscum, glabrum, stipite plerumque dilutiore, urceolatum, subclausum, humidum aperatum, flavescent, 0,5-7 diam., prosenchymatice contextum; ascis clavatis, apice incrassatis, 8-sporis, 40 \approx 5-6; sporidiis oblongis vel

Discomyceteæ, Pezizeæ, Hyalosporæ, Helotium. 227

subfusiformibus, obtusis, plerumque rectis, hyalinis, distichis, 6-8 = 2; paraphysibus filiformibus, apice vix incrassatis, hyalinis.

Hab. ad calamos sicclos Nardi strictæ in alpibus editioribus Tyrolensibus juxta glaciem æternam. — An *Phialea*?

72. **Helotium filicolum** Hazsl. Einig. Disc. pag. 14 et Magyar. szab. 923
Disc. t. X, f. 22. — Flavum, stipitatum; disco plano, demum convexo, margine crasso, reflexo; stipite deorsum incrassato, disco sub-duplo longiore; ascis anguste clavatis, stipitatis, octosporis; sporidiis clavatis, deorsum acutioribus, 8-guttulatis simulateque 7-septatis.

Hab. in stipibus filicis pr. Eperies Hungariæ.

73. **Helotium Mousseotii** (Pers.) Sacc. *P. Mousseotii* Pers. Myc. Eur. 924
p. 289, t. XII, f. 5, Fr. Syst. Myc. II, p. 135. — Confertum; cululis tereti-ventricosis cum stipite brevi confluentibus, disco planiusculo aurantiaco, margine tenuissime fimbriato albicante.

Hab. in Vogesia in montium summitate prope nives liquescentes ad terram. — Species pulchra, minuta, carnosa, hinc vix excavata, margine scarioso lacero. Species minus nota, forsitan *Biatula campestris*. Ex icone carneo-rubra, crusta quadam pulverulenta albicante.

V. Disco ochraceo, lutescente, fulvo, brunneo, testaceo. (93-103)

74. **Helotium epiphyllum** (Pers.) Fr. Summa p. 356, Pat. Tab. anal. 925
fig. 93, *Pez. Stizenbergerii* Rabenh. *Pez. epiphylla* Pers. disp.
p. 72, Syn. p. 669, Fr. Syst. Myc. II, p. 137. — Subsessile, glabrum, convexo-planum marginatum, ochraceo-pallidum; ascis tereti-clavatis, 100 = 9-10; sporidiis fusoideis curvatis, 2-4-guttatis, 9-12 = 4-5; paraphysibus filiformibus.

Hab. ad folia dejecta putrescentia Fagi, Carpinii, Quercus, Betulæ etc., in Suecia, Germania, Gallia, Britannia, Belgio, Italia, etiam in America bor., Fuegia, Africa austr., Ceylon. — Basi papillaris, cum qua adfixa, passim in stipitem e loco plus minus longum elongatur. Cupula carnosoceracea, subtenuis, 3 mm. lata, demum rufescit. Ad ultimas species accedunt plures formæ mediæ v. c. *P. minutissima* et *sigillatoria* Batsch Cont. I, f. 142, 143, *P. pallida* Schum. Saell. p. 428.

75. **Helotium saprophyllum** Peck 29 Rep. p. 55. — Ascomatibus minutis, stipitatis, ochraceis, planis; stipite tenui, sursum incrassato, disci diametrum subexcedente; ascis sublanceolatis; sporidiis fusoideis v. subclavatis, distichis, 1-binucleatis, 20 = 6.

Hab. in foliis dejectis, Lake Pleasant Amer. bor. — *Helotio fastidioso* affine.

76. **Helotium helodinum** Dur. et Lév. Expl. scient. Alg. Bot. t. 28, 927 fig. 3 (*Peziza*) [absque diagnosi]. — Plano-scutellatum, stipitatum, ochraceo-fulvum; disco badio, concaviusculo, pallidius marginato; stipite cylindraceo; ascis tereti-clavatis; sporidiis oblique monostichis, fusoideis, rectis, hyalinis.

Hab. in foliis dejectis Ilicis in Algeria. — Ascoma 2-3 mm. latum, stipes 2 mm. longus.

77. **Helotium leucopus** Mont. Chil. pag. 410, Syll. Crypt. n. 632. — 928 Hypophyllum, erumpens, sparsum, glabrum; cupulis obconico-turbinatis, fulvis, disco planiusculo, stipite breviusculo pallido, basi subbulbosa byssina alba; ascis majusculis, clavatis; sporidiis fusoides distichis, 4-6-guttulatis.

Hab. ad paginam inferiorem foliorum Perseæ Lingue, Chile (Cl. GAY). — Ascoma 2 mm. alt., $\frac{1}{2}$ mm. lat.

78. **Helotium Friesii** (Weinm.) Sacc. *Pez. Friesii* Weinm. Hym. Ga- 929 ster. pag. 469. — Gregarium, totum argillaceum, glabrum, asco- mate plano-convexo; margine tumidulo; stipite brevi crassiusculo, obconico.

Hab. ad folia dejecta Populi Tremulæ et Betulæ albæ in Rossia. — Substantia carnosa, crassiuscula. Ascoma $\frac{1}{2}$ -1 mm. latum, margine plano. Stipes interdum 1 mm. et paullo ultra long. et ad basim obscurior.

79. **Helotium lentum** B. et Br. Fung. of Ceyl. n. 961. — Stipite cylindrico, nigro, cupula extus rugosa; hymenio siccatione rufo, mado lutescente; sporidiis brevibus acicularibus, 6-7 μ . longis. — 930

Hab. in foliis, capsulis putridis in insula Ceylon.

80. **Helotium Ostruthii** (Saut.) Sacc., *Peziza Ostruthii* Saut. Pitzl. 931 Salzb. p. 9, Wint. Hedw. 1881, p. 132. — Ascomate minuto, 0,5 mm. lato, ceraceo, sordide luteo, plano-convexo; stipite brevissimo; ascis clavatis, octosporis, 90-100 \approx 8-9; sporidiis distichis, cylindra- ceo-fusoideis, demum (spurie?) triseptatis, hyalinis, saepè parum curvatis, 16-18 \approx 3,5; paraphysibus filiformi-clavulatis.

Hab. in caulibus putridis Ostruthii ad Mittersill Salzburg.

81. **Helotium conisteum** Dur. et Lév. Expl. sc. Alg. tab. 28, fig. 10 932 (*Pez.*) [absque diagnosi]. — Gregarium, cupulato-infundibuliforme, stipitatum, pallide alutaceum; disco concavo; stipite teretiusculo, brevi; ascis clavatis, stipitatis, filiformi-paraphysatis, 8-sporis;

sporidiis distichis cylindraceis, inæquilateralibus, latitudine 8-plo longioribus, hyalinis.

Hab. ad caules putres in Algeria. — Ascoma 1 mm. lat., stipes $\frac{1}{2}$ - $\frac{3}{4}$ mm. long.

82. **Helotium sublateritium** B. et Br. Ann. Nat. Hist. n. 1488, Phillips 933 Disc. pag. 161. — Pallide lateritium, stipitatum; ascomate plano, subitus venoso; margine elevato; stipite brevi cylindrico, initio albidio-villoso, dein glabro; ascis linearibus; sporidiis monostichis, breve fusoideis, biguttatis, 25 \approx 5.

Hab. in caulibus herbaceis, Glamis Britanniæ.

83. **Helotium vitigenum** De Not. Comm. I, pag. 377, Sacc. Fung. Fl. 934 tab. 1343. — Ascomate stipite mediocri instructo, patellato, disco pallide stramineo, subconvexo, tenuiter marginato, subitus pallecente; ascis 110 \approx 12; sporidiis ellipsoideo-fusoideis, 4-guttulatis, 18-20 \approx 6, hyalinis.

Hab. in sarmenis emortuis Vitis viniferæ, Valle Intrasca et in agro veneto Ital. bor.

84. **Helotium pullatum** Gerard in Cooke Disc. U. S. I, p. 298 et in 935 Thüm. Pilze Weinst. pag. 89. — Gregarium; ascomate 1-2 mm., vivo sordide ochraceo, sicco fuliginoso, primo clavato, dein paululum obconico, extus saturatius colorato; stipite brevi in cupulam expanso; disco plano, sicco concavo; ascis cylindraceis angustissimis; sporidiis tereti-truncatis, 2-4-guttatis simulateque septatis, 18-20 \approx 4.

Hab. in sarmenis Vitis Labruscæ ad New York Americæ borealis.

85. **Helotium sepium** (Desm.) Sacc., *Peziza sepium* Desm. 18 Not. 936 p. 5. — Erumpens minimum carnoso-ceraceum, sparsum v. cæpitosum, stipitato-turbinatum, glabrum; stipite brevi, crasso, pallido; disco plano-convexo, brunneo-ochraceo, sicco subpulverulento; margine vix prominente sinuoso, aurantiaco vel testaceo, subtuberulo; ascis amplis clavæformibus, 125-150 μ . long.; sporidiis magnis, ellipsoideis, 20-25 μ . long., hyalinis.

Hab. in ramis exsiccatis Crataegi Oxyacanthæ in Gallia bor. — Ascoma $\frac{2}{3}$ mm. diam.

86. **Helotium ochraceum** (Grev.) Phill. Berk. Outl. p. 372, *Peziza* 937 Grevill. Scott. t. V. — Breve stipitatum vel sessile, glabrum, flavido-brunneum, carnosulum; disco pallidiore, concavo, piano v. convexo, granulis nitentibus consperso; ascis cylindraceo-clavatis; sporidiis octonis.

Hab. in cortice ramorum dejectorum pr. Edinburgh et Shrewsbury Britanniæ.

87. **Helotium platypus** Hazsl. Einig. Disc. p. 14 et Magyar. szab. Disc. 938 t. 11, f. 23. — Stipitatum, flavo-brunneum; disco plano v. subconcavo; stipite longitudine vario, cylindrico, disci diametro pluries longiore, basi mycelio subcirculari stellato instructo; ascis clavatis, stipitatis; sporidiis oblongo-fusoideis, latitudine triplo-quadruplo longioribus.

Hab. in ramis Alni in paludosis Hungariæ. — An *Phialea*?

88. **Helotium? chlorosplenoides** Speg. Fung. Fueg. n. 330. — Infundibuliforme, majuscum 3-5 mm. diam., ligno-pallens, disco glabro obscuriore extus fibrilloso-striatum; stipite tereti, 3-5 mm. long., vix $\frac{1}{2}$ mm. cr., matrice late infuscata insidente; ascis cylindraceo-subclavatis, antice non v. vix attenuatis apice truncatis, postice longe attenuato-stipitatis, 100-110 \approx 6-6,5; paraphysibus filiformibus non v. vix longioribus laxè obvallatis, octosporis; sporidiis oblique monostichis, elliptico-fusoideo utrinque acutissimis, 10 \approx 4-5, rectis vel subcymbiformibus, hyalinis non vel grosse 1-guttulatis.

Hab. ad ramulos fagineos dejectos putrescentes in silvis pr. Port S. John in Staten island Fuegiæ. — Species certe a genere *Helotio* Fr. separanda, habitu *Chlorosplenio* Fr. simillima. Speg. loc. cit.

89. **Helotium indeprendum** Bizz. Fung. Ven. nov. v. crit. p. 3, t. III, 940 f. 6, a, b, c, d, e; Sacc. Fungi Ital. t. 1333. — Ascomatibus carnosulis, subsessilibus, convexiusculis; margine acutiusculo infexo, ochraceo-rufis dein obscurioribus, 0,5 mm. diam., stipite brevissimo, crassiusculo (vix manifesto); ascis subclavatis, octosporis, paraphysatis, 65-75 \approx 8; sporidiis oblique monostichis, hyalinis, oblongo-ellipsoideis, utrinque uniguttulatis, 9-11 \approx 3.

Hab. in ramulis et foliis siccis sed nondum putrescentibus Thujæ orientalis in horto botanico patavino. — Species repertu difficilis, quia cum matrice sua saepius concolor. Ab *Hel. epiphylo* colore rufescente et forma ascomatum, ascis sporidiisque minoribus facile distinguenda species.

90. **Helotium salicellum** Fr. Summ. Veg. Scand. p. 356, Karst. M. F. 941 I, p. 113, *Peziza salicella* Fr. Syst. Myc. II, p. 133, Rehm Asc. n. 208, Sacc. F. ital. 1344. — Gregarium, breviter stipitatum, firmulum, testaceum v. ochraceo-pallescens; cupula planiuscula, orbiculari, adpressa, margine vix prominulo, 1-2 mm. lat.; stipite

æquali pallidiore brevi, 0,5-1,5 mm. alt.; ascis cylindraceo-clavatis, 75-11-13, obturaculo jodo obsolete cœrulescente; sporidiis distichis fusoideo-elongatis, rectis v. subinde leniter curvulis, 2-4-guttulatis v. simulate 1-3-septatis, 25-28-5-6; paraphysibus filiformibus, 2 μ . cr.

Hab. in ramulis *Salicis* in Suecia, Fennia, Britannia, Belgio, Italia, Germania, Amer. bor. — Sec. Fuck. sporidia 12-4, an ex errore?

91. **Helotium sublenticulare** Fr. Summ. Veg. Scand. pag. 355, Karst. 942

M. F. I, p. 118, Rehm Ascom. n. 654 (forma subsessilis), *Peziza citrina* Flor. Dan. t. 1971, f. 3, nec Batsch., *Pez. subferruginea* Nyl. Obs. p. 42 (ut videtur). — Gregarium, stipitatum, firmulum, albidum vel lutescente albidum, siccum testaceo-pallidum aut interdum ferruginascens; cupula discoidea, plana aut interdum demum convexiuscula, sicca patelliformi, margine integro, disco lutescente, sicco dilute luteo, subvitellino, interdum ferrugineo, 1-5 mm. lat.; stipite plus minus elongato, sursum incrassato, deorsum leviter flocculoso, subinde flexuoso, 1-4, raro-15 mm. alt.; ascis cylindraceo-clavatis, 110-120-9-10, obturaculo apicali jodo levissime vel vix cœrulescente; sporidiis vulgo oblique monostichis, oblongato-ellipsoideis, 2-guttatis, plerumque rectis, 15-20-4-6; paraphysibus filiformibus.

Hab. ad ramulos exsiccatos Alni, Aceris, Coryli et Rubi in Fennia, Suecia, Dania, Germania.

92. **Helotium conscriptum** Karst. M. F. I, p. 118. — *H. sublenticu-* 943

lari externa facie sat simile; sporidiis autem minoribus, 10-16, (plerumque 12-14)-3,5-5, eguttulatis aut aliquando guttulis paucis per exiguis præditis; ascis 8 μ . cr.

Hab. ad ramulos *Salicis* in Fennia et Lapponia. — Var. *caulinolum* Rehm in Hedwigia 1885, p. 223: ascomatibus globosis, dein cyathoideis, stipite crasso 2-4 mm. long., disco explanato, demum 2-3 mm. diam., flavidis v. flavo-vitellinis, saepe extus striatus v. rugulosis, stipite interdum dilutiore, margine integro saepe dilutiore, sparsis prosenchymatice, dilute flavidè contextis; ascis clavatis apice incrassatis, 8-sporis, 70-6; sporidiis superne 2-stichis, oblongis, obtusis, rectis v. curvatulis, 1-cellularibus, plerumque nucleolis 2 apicalibus instructis, hyalinis, 12-3 (denique 2-cellularibus?); paraphysibus hyalinis filiformibus, haud clavatis circ. 2 μ . crass. Ad caules putridos Aconiti Napelli prope Kleines Kohlbach-Thal, Hohe Tatra Hungariæ (PROF. LINHART). — Forma

alpestre Rehm in Hedwigia 1885, p. 229: ascomatibus orbiculatis v. repando flexuosis, plerumque crasse brevissime stipitatis v. subsessilibus, 0,5-1,5 mm. diam., albo-flavidis, margine dilutiore, gregariis, hyalino-prosenchymatico-contextis.

93. **Helotium montellicum** Sacc., *Helotium macrosporum* Sacc. Mich. 944 II, p. 535 (1882), F. ital. f. 1332, nec Peck (1874). — Ascomatibus gregariis, totis lutescentibus, ceraceis subsessilibus, $\frac{1}{2}$ - $\frac{3}{4}$ mm. lat.; disco convexo-pulvinato, margine non discreto, subsinuoso; stipite brevissimo, crassiusculo; ascis late clavatis, 120-130 = 20, breve stipitatis, filiformi-paraphysatis, apice tenuato-truncatis, octosporis; sporidiis distichis oblongo-phaseoliformibus, curvulis, utrinque rotundatis, 25 = 7,5-8, varie guttatis v. granulosis, hyalinis.

Hab. in ramis corticatis Corni sanguineæ in Valle dell'Aqua silvæ Montello Ital. bor. — Ascomatibus convexis subsessilibus sporidiisque amplis, obtusis mox dignoscenda species.

94. **Helotium uliginosum** Fr. Summ. Veg. Scand. p. 355, Karst. M. 945 F. I, p. 121, *Peziza uliginosa* Fr. Syst. Myc. II, p. 138, *Mollisia uliginosa* Oudem. — Sparsum v. gregarium, breve v. longiuscule stipitatum, ceraceo-aquosum, firmulum; cupula concaviuscula vel convexiuscula, pallido-albida aut e lutescente ochracea, sicca obscurata testacea v. subferruginea, saepe flexuosa et umbilicata, 2-4 mm. lat.; stipite livido-pallescidente v. pallido, cavo, 1-8 mm. alt.; ascis cylindraceo-clavatis, 65-90 = 6-8, obturaculo jodo levissime v. vix cœrulecente; sporidiis plerumque oblique monostichis, oblongatis, plerumque 2-guttulatis, 7-14 = 3-4; paraphysibus filiformibus incrassatulis.

Hab. ad ramulos, cortices lignaque arborum frondosarum, imprimis Betulæ, Salicis et Fraxini in Suecia, Fennia, Hollandia, Britannia. — Stipes deorsum, rarius sursum incrassatus, exsiccatus longitudinaliter plicatus v. compresso-collabens, nunc pallidus nunc fuligineo-virgatus. — Var. β . **cortisatum** Karst. M. F. I, p. 122: cupula fere sessili lutescenti-pallida; sporidiis fusoideo-oblongatis aut elongato-ellipsoideis, 6-9 = 3-4; paraphysibus gracilibus, guttulis fulvescentibus præditis. Supra corticem Betulæ vetustum prope Mustiala Fenniæ.

95. **Helotium fuegianum** Speg. Fung. Fueg. n. 332. — Densiuscule 946 sparsum obconicum concaviusculum, 0,6-1 mm. diam., margine acuto integerrimo vel repandulo, intus pulchre flavidio-melleum, extus pallidius glabrum, in stipitem brevissimum ac crassum candidum, non v. vix basi puberulum desinente; ascis cylindraceis v. cylin-

draceo-subclavulatis, antice obtuse rotundatis, deorsum plus minus-ve longe attenuato-pedicellatis, 55-65 \approx 5, octosporis; paraphysibus filiformibus nonnihil longioribus densiuscule obvallatis; jodi ope ascorum apice dilute, obturaculo intense cœrulescente; sporidiis oblique monostichis, ellipticis utrinque obtusis, 5-8 \approx 2-3 rectis v. inæquilateralibus, hyalinis continuis; an demum medio 1-septatis.

Hab. vulgatissimum ad ramulos dejectos putrescentes Fagorum in Fuegia et Staten island Amer. austr.

96. **Helotium fusco-brunneum** Pat. et Gaill. in Bull. Soc. Myc. Fr. 947 IV, pag. 101. — Sparsum vel gregarium; cupula 2-4 mm. diam., concava dein appanata, glabra, in stipite brevissimum attenuata, ferrugineo-brunnea, nitida, extus levi; ascis clavatis, substipitatis, 45 \approx 6; sporidiis ovoideis, utrinque obtusis, 6-8 \approx 1,5-2, rectis vel leniter curvulis, biguttulatis; paraphysibus linearibus, hyalinis, ascos longitudine æquantibus.

Hab. ad corticem arboris, inter muscos, inter Maipures et S. Fernando Americæ austr. (GAILLARD).

97. **Helotium Fergussoni** Sacc. *Helot. melleum* B. et Br. Ann. N. A. 948 n. 1457 (1875) nec B. et Br. Fung. of Ceyl. (1873). — Pallide melleum, planum, subflexuosum, margine clavato, infexo; stipite brevi, tereti; ascis elongato-fusoideis; sporidiis distichis, fusoideis, rectis curvulisve, pluriguttulatis 30 μ . longis.

Hab. in ligno putri New Pitsligo Britanniae (FERGUSSON). — Ascoma 2 mm. lat., stipes 1 mm. long.

98. **Helotium ferrugineum** (Schum.) Fr. Summa p. 356, Sacc. F. ital. 949 f. 1331, Phill. Disc. p. 154, t. V, f. 30, *P. lenticularis* var. Bull. t. 300. — Substipitatum, obconicum, disco concavo luteo-ferrugineo, extus margineque tumido pallide luteum; ascis cylindraceo-clavatis; sporidiis elliptico-fusoideis v. clavulatis, 2-guttulatis, 10-13 \approx 3-5; paraphysibus filiformibus.

Hab. ad truncos vetustos Quercus, ad ligna putrida varia in Suecia, Belgio, Britannia, Germania, Italia, America bor., Africa austr. — Valde ad *H. lenticulare* accedit, sed habitu et loco distinctum. Minus, gregarium, primo turbinatum, dein magis dilatatum sed vix 1-2 mm. latum; stipitis rudimentum exiguum pallidum; margo integerrimus. — β . **H. nanum**, *Octosp. nana* Hedw. loc. cit. p. 29, t. IX, f. A: cinnamomeum, inferne in stipitem attenuatum. E terra pullulare dicitur in Germania.

99. **Helotium robustius** Karst. M. F. I, p. 119. — Sparsum vel gre-

garium, stipitatum, pallidum lutescens v. interdum rosellum, cupula planiuscula, sicca patelliformi, margine integerrimo, disco lutescente v. subluteo, 1-2 mm. lat.; stipite 1-4 mm. alt.; ascis cylindraceo-clavatis, 90-115 \approx 7-10 apice jodo vix v. levissime cœrulescentibus; sporidiis mono- v. subdistichis, oblongatis v. oblongato-ellipsoideis vulgo eguttulatis, 7-14 \approx 3-6; paraphysibus filiformibus, guttulatis, sursum leviter incrassatis.

Hab. ad culmos Phragmitis communis aliorumque graminum, calamos Typhæ latifoliæ nec non ad fructus Cirsii arvensis in Fennia et Ital. bor. — Forma **macrospora** Rehm Asc. n. 562: sporidiis 15-18 \approx 3-5, ascis 60 \approx 9-10. In Germania.

100. **Helotium maritimum** (Rob. et Desm.) Sacc.. *Peziza maritima* 951 Rob. et Desm. XI Not. p. 366. — Erumpens, minutum, stipitatum, carnosoceraceum, glabrum, cinnamomeum; stipite brevi, crasso in cupulam dilatato; cupula crassiuscula, juniore plano-concava, dein subconvexa; ascis magnis, flexuosis tubulosis, 150-160 μ . long.; sporidiis ovato-oblongis, utrinque acutiusculis, 2-guttulatis, 12-13 μ . long.

Hab. in foliis exsiccatis Psammæ arenariæ, Calvados Galliæ. — Stipes $1\frac{1}{4}$ - $1\frac{1}{3}$ mm. long. et cr.; ascoma $1\frac{1}{2}$ -1 mm. lat.

101. **Helotium gramineum** Phill. Disc. p. 155. — Stipitatum, planum 952 v. convexum, glabrum, dilute ochraceum; stipite brevi, subobtuso, subinde medio angustato, sursum in cupulam expanso; ascis subclavatis; sporidiis ellipsoideis, 1-2-guttulatis, 8-11 \approx 4-5; paraphysibus filiformibus.

Hab. ad culmos graminum, Kingsly Britanniæ (PLOWRIGHT). — Ascoma 1-3 mm. lat. et alt., margo obtusus.

102. **Helotium equisetinum** Quél. X Suppl. pag. 13, t. IX, fig. 13 et 953 Grevillea 1879, p. 38. — Ascomatibus planis, dein excavatis, 500-800 μ . diam., glabris tenuibus, teneris, pallide lutescentibus, breve stipitatis, hymenio plano, nitide citrino; sporidiis bacillaribus 5-6 μ . longis.

Hab. ad caulem Equiseti limosi, Alsace. — Stipes lutescens 0,5 mm. long.

103. **Helotium phascoïdes** Fr. Summa pag. 355, *Pez. phascoïdes* Fr. 954 Syst. Myc. II, p. 138, *P. dubia* Batsch Cont. I, fig. 145, vix obstat. — Ceraceo-aquosum, minutum, testaceo-pallidum; cupula turbinata plana; stipite brevi subæquali.

Hab. ad terram humidam stipulis v. Phascis forsitan innatum in Suecia, Germania et Britannia. — Solitarium, glabrum, varium.

Stipes farctus, teres 2 mm. longus ecoloratus; cupula 1 mm. lata, pallida. Asci hucusque ignoti.

VI. Disco umbrino, fusco, atro. (104-118)

104. **Helotium hysteroides** Rehm Ascom. Alp. n. 6. — Gregarium, 955 crasse stipitatum, stipite c. 1 mm. alt., siccum hysteriformiter elipticum, rima longitudinali labiis crassis marginata marginibusque involutis, fuscum vel nigrofuscum, disco rubro-fusco, humectatum subexplanatum, lentiforme, 1,5-2 mm. diam., prosenchymatico-fuscidulo-contextum; ascis clavatis, subcurvatis, 8-sporis, 50 \approx 8; sporidiis oblongis, obtusis rectis v. subcurvatis, saepe centro 1-nucleatis, hyalinis, distichis, 15 \approx 4,5; paraphysibus superne sensim incrassatis filiformibus, clava hyalina, c. 3 μ . cr.

Hab. ad caulem putrescentem Aconiti varieg. prope glaciem æternam, Kartel-Gletscher in valle Moos alpium Tyrolensium (BRITZELMAYR).

105. **Helotium nigrescens** (Cooke) Rehm Asc. n. 307, *Peziza* (Hymenoscyphe) *nigrescens* Cooke Hedw. 1875, pag. 83. — Stipitatum, atro-brunneum, nigrescens, minimum, firmulum, cupula primum clavata, dein expansa, applanata, 0,3 mm., margine vix elevato, disco pallidiore, subcinereo; stipite crassulo, superne in cupulam expanso, duplo longiore, 0,5 mm. longo; ascis subcylindraceis, 60 \approx 9; sporidiis elongatis, fusiformibus obtusis, rectis v. curvulis, binucleatis hyalinis 12 \approx 3.

Hab. ad caules Erigeronis, New Jersey Amer. bor.

106. **Helotium fumigatum** S. et Sp. Mich. II, p. 78, F. ital. f. 1337. 957 — Ascomatibus gregariis, concavis, stipitatis, $\frac{1}{2}$ - $\frac{3}{4}$ mm. altis et lati totis fuscescentibus margine in sicco connivente pallidiore, extus levibus v. obsolete striatulo; excipuli contextu prosenchymatico, fuscidulo; ascis cylindraceo-clavatis, 60 \approx 5-6, apice rotundatis, deorsum acutato-stipitatis, paraphysatis, octosporis; sporidiis distichis, cylindraceis, utrinque obtusiusculis, 8-9 \approx $1\frac{1}{4}$ - $1\frac{1}{2}$, rectis vel curvulis, 2-guttulatis granulosisve hyalinis.

Hab. in ramulis emortuis Catalpæ syringifoliæ, Conegliano Ital. bor., et Clematidis Vitalbæ in Gallia. — *Helotio Cacaliæ* sec. descript. affine videtur. *Peziza fumosa* Fr. Summa p. 353 e loco forte huic quoque affinis; sed ubi descripta?

107. **Helotium Krascheninikowii** (Weinm.) Sacc., *Pez.* *Krascheninikowii* Weinm. Hym. Gaster. p. 465. — Solitarium, ascomate carnosso-membranaceo, turbinato-globoso, dilute-brunneo, striatulo,

margine involuto, disco umbrino; stipite brevi, crassiusculo, nigrescente.

Hab. in ramulo rejecto decorticato alneo semel in Rossia. — Minuta species, sed valde memorabilis et elegans. Ascoma, vix $\frac{1}{2}$ mm. latum, initio magis turbinatum, dein turbinato-globosum, humidum obscure, siccum dilute brunneum, striis tenuibus, vagis, subelevatis ornatum, margine semper involuto; stipes $\frac{1}{2}$ mm. et ultra long. subteres, crassiusculus et sub lente subsulcatus.

108. **Helotium complanatum** Fr. Syst. Myc. II, p. 158, *Sarea complanata* Fr. Elench. II, pag. 14. — Atrum, capitulo convexo compresso, stipite inferne dilatato. 959

Hab. in ramulis Alni cordatæ in Kamtschatka. — Totum atrum, habitu *Calicii*, sed crusta nulla et discus solidus. Stipes 4 mm. circiter longus, sursum attenuatus; capitulum subtus plano-convexum adeo complanatum, ut a latere inspectum stipite paullo modo crassius. Primo obtutu facile crederes cupulam esse complicato-complanatam, sed discus omnino levigatus absque rimæ vestigio.

109. **Helotium sclerotoides** Berk. Outl. p. 371. — Convexum, umbilicatum, pallide rufo-brunneum, infra concavum et in stipitem brevem obconicum confluens; ascis tenuibus; sporidiis.... 960

Hab. in ligno putri Appin et near Bristol Britanniæ. — Primo obtuto *Sclerotium quercigenum* in mentem revocat. Ascoma 2 mm. lat.

110. **Helotium rufo-corneum** B. et Br. Fung. of Ceyl. n. 962. — Stipe brevi, sursum dilatato; cupulis subtus rugosis, castaneis; hymenio concolori; paraphysibus clavatis; sporidiis fusiformibus, apicibus auctis, hyalinis, 30μ . long. 961

Hab. in ligno putri in insula Ceylon.

111. **Helotium sphærophoroides** Phill. et Harkn. Calif. p. 1. — Ascomate minuto, stipitato, subturbanato, glabro, carneo-fuligineo, disco piano dein depresso, immarginato; ascis late clavatis, octosporis; sporidiis cymbiformibus, crasse uninucleatis, $14-19 \times 6$. 962

Hab. gregatim in pagina inferiori foliorum *Sequoiae* semper-virentis in California Amer. bor. — Habitus *Sphærophori*.

112. **Helotium spadiceo-atrum** (Mont.) Sacc., *Pez. spadiceo-atra* Mont. 963 Prodr. J. Fern. n. 26 et Fl. Chil. VII, p. 401. — Spadiceum; culpula stipitata, hypocrateriformi, tandem planiuscula, disco atro; ascis clavatis inter paraphyses numerosas nidulantibus, 75μ . long.; sporidiis 8 oblongis utroque fine attenuatis, 10μ . long.

Hab. in silvis montosis Juan Fernandez ad folia dejecta (BERTERO). — Ascoma 1 mm. lat., stipes ascomate paullo longior.

113. **Helotium coracinum** Dur. et Lév. Expl. Sc. Alg. tab. 28, fig. 4 964 (Pez.) [absque diagnosi]. — Gregarium, hypophyllum, infundibuliforme, stipitatum, fuscum, disco concaviusculo, stipite tereti brevi; ascis cylindraceis, 8-sporis; sporidiis monostichis ovoideis, hyalinis, latitudine duplo longioribus; paraphysibus filiformibus, subinde ramulosis.

Hab. in foliis emortuis *Quercus Ilicis* in Algeria. — Ascoma $\frac{1}{2}$ mm. lat., stipes paullo brevior.

114. **Helotium titubans** Mont. Fl. chil. 408, t. 8, fig. 6, Syll. Crypt. 965 n. 631. — Amphigenum, gregarium; cupula minutula, planiuscula, apice stipitis attenuati atrati inflexo nutante, disco umbrino 70-80 μ . long.; sporidiis oblongo-fusiformibus simplicibus, 10 \approx 3.

Hab. ad utramque paginam foliorum coriaceorum, Chile (GAY).

115. **Helotium vasaense** Karst. M. F. I, p. 125. — Sparsum, breviter, 966 stipitatum; cupula planiuscula, brunneo-atra (saltem in statu sicco), circiter 2 mm. lat.; ascis cylindraceo-clavatis, 8 μ . cr., apice jodo vix tinctis; sporidiis fusoideo-oblongatis, utroque apice acutiusculis, 2-guttulatis vel eguttulatis, 10-14 \approx 3; paraphysibus filiformibus, fulvo-guttulatis.

Hab. supra Sphagna emortua in regione vasaensi, Quarken Fenniae. — *Ombr. subspadiceæ*, ut videtur proximum. Stipes brevissimus, crassulus; asci et paraphyses arcte conferti.

116. **Helotium triste** Sacc. Mich. II, p. 379, F. it. f. 1338. — Ascomatibus gregariis, scutellatis, ubique nigricantibus, glabris, solidiusculis, stipitatis, disco concaviusculo $\frac{1}{2}$ mm. diam.; stipite cylindraceo $\frac{1}{3}$ mm. alto; contextu excipuli anguste prosenchymatico, castaneo, margine saturatiore; ascis tereti-fusoideis, breve stipitatis, apice obtusiusculis, 100-110 \approx 12-14; paraphysibus bacillaribus, hyalinis, exterioribus vero granulato-virescentibus obvallatis, octosporis; sporidiis distichis, crassiusculis, fusoideis, saepe leniter curvis, utrinque acutis, 20-22 \approx 4-5, crasse 1-2-guttatis, hyalinis.

Hab. in stipitibus Aspidii Filicis-maris emortuis in *M. Generoso* (PENZIG). — Affine, ut videtur, *Pezizæ pteridicolæ* Crouan Finist. 50, sed satis differt colore nigricante, nec griseovirente, sporidiis fusoideis nec subcylindricis etc.

117. **Helotium campanulæforme** Fuck. Symb. p. 315. — Sparsum, minutum; cupulis campanulatis, in stipitem cupulam æquantem, tenuem, concolorem contractis, pallide fuscis, tot. 1-2 mm. alt., $\frac{1}{2}$

- 238 Discomyceteæ, Pezizeæ, Hyalosporæ, Helotium.
diam.; ascis clavatis, 8-sporis, $60 \approx 8$; sporidiis distichis, oblongo-
ellipticis, continuis, 3-guttulatis, $12 \approx 3-4$, hyalinis.
Hab. in stipitibus Aspidii Filicis-feminæ in silva Æstrich
Germaniæ. — Affine *Pezizella chrysostigmati*.
118. **Helotium sordidum** Phill. Grevillea XV, pag. 17. — Sparsum; 969
ascomate stipitato, concavo dein plano, sordide brunneo, margiae
pallidiore, integro; stipite obtuso, pallido, basi albo, tomentoso;
ascis cylindricis octosporis; sporidiis ellipsoideis, $7-9 \approx 3-4$; para-
physibus non visis.
Hab. in ligno putri in Nova Zelandia (COLENZO). — *Hel. fibu-*
liformi affine sed sporidia diversa.
- B. Sessilia v. subsessilia.
- I. Disco albo, pallido, sordide pallido. (119-127)
119. **Helotium immarginatum** Karst. M. F. I, p. 127. — Gregarium, 97
sessile, immarginatum, saepe repando-incisum, convexum, albidum,
siccum fere concolor, planiusculum vel convexiusculum, latit. 0,4
mm.; ascis cylindraceo-clavatis, $56-75 \approx 8-10$, apice jodo non vel
obsoletissime cœrulescentibus; sporidiis distichis vel oblique mono-
stichis, oblongatis, 2-guttulatis v. spurie tenuiter uniseptatis, ple-
rumque rectis, $8-11,5 \approx 2,5-3,5$; paraphysibus sat numerosis, fili-
formibus, gracilescentibus, intus oleoso-guttulatis, $1-1,5 \mu$. cr.
Hab. supra lignum betulinum vetustum in Fennia saltem australi passim. — *H. trabinello* affine, sed omnibus partibus minus,
colore paraphysibusque numerosioribus diversum.
120. **Helotium Kunzei** (Saut.) Sacc., *Peziza Kunzei* Saut. Pilze Salzb. 971
p. 10. — Sessile; ascomatibus carnosulis, sordide albis, margine
integerrimo.
Hab. in disco putri arborum prope Mittersill Salzburg.
121. **Helotium hydrogenum** Peck 29 Rep. Bot. p. 56. — Gregarium 972
vel sparsum, sessile, leve, extus brunneolum, disco subplano, mar-
ginato, pallido vel flavescente, sicco brunneo vel viridi tincto; ascis
subclavatis; sporidiis subcylindraceis, leniter curvulis, saepe gut-
tulas minutas numerosas foventibus, $12-15$, μ . longis.
Hab. in ligno putrescente in aquis ad Sandlake Americæ borea-
lis, sociis *Mitrulâ paludosâ* et *Vibrissedâ truncorum*.
122. **Helotium Doliolum** (Saut.) Sacc., *P. Doliolum* Saut. Pilz. Salzb. 973
pag. 9. — Minimum, turbinatum, annuliforme, subsessile, niveum,
ascomate turbinato; stipite brevissimo, æquali.
Hab. in ligno putri prope Mittersill Salzburg.

123. **Helotium scoparium** Cooke in Grevillea IV, p. 112. — Ascomata 974
tibus sparsis v. subgregariis, sessilibus, pallidis, disco convexo al-
bido totaque planta, demum siccando grisea; ascis clavatis, 100
 \approx 20; sporidiis cylindraceis obtusis, rectis curvulisve binucleatis,
20-25 \approx 5 plasmate demum partito; paraphysibus filiformibus, gra-
nulosis.

Hab. in ramis emortuis Sarothamni scoparii, Dupplinkastle
Pert Britanniæ. — Ascoma 1 mm. lat.

124. **Helotium eburneum** Cr. Fl. Finist. p. 47. — Ascomate albo, tur- 975
binato, 4 mm. diam., hymenio albido; ascis subclavatis octosporis;
sporidiis fusiformibus, granulosis; paraphysibus capillaribus.

Hab. ad folia putrescentia et emortua locis udis silvarum Fi-
nistère Galliæ. — Contextus e cellulis basi ovoideis, apicem asco-
matis versus filamentosis, hyalinis, intertextis efformatus.

125. **Helotium niveum** Cr. Fl. Finist. p. 47, tab. ult. fig. 5. — Asco- 976
mate albo, sessili, glabro, disciformi, 200 μ .; ascis late clavatis
octosporis; sporidiis ovoideis, majusculis, granulosis; paraphysibus
filiformibus, ascos superantibus.

Hab. in Jungermannia albicante Finistère Galliæ.

126. **Helotium Blechni** Cr. Finist. pag. 47. — Ascomate albo sessili 977
crasso, hemisphærico subconcavo, 2-4 mm. lat. disco griseo; ascis
clavatis, octosporis; sporidiis oblongis, hyalinis; paraphysibus fili-
formibus.

Hab. in foliis emortuis Blechni Spicant in silvis Finistère
Galliæ. — Cellulæ excipuli basi ovoideæ, versus marginem prosen-
chymaticæ constipatæ.

127. **Helotium albidum** Cr. Finist. p. 47. — Ascomate albido, glabro 978
hemisphærico, sessili, 2-3 mm. lat., disco albo; ascis clavatis,
octosporis; sporidiis oblongis, granulosis; paraphysibus simplicibus
v. furcatis, sursum uncinatis.

Hab. ad terram silvaticam Finistère Galliæ. — Excipuli cellu-
læ prosenchymaticæ perangustæ, stipatæ, anastomosantes.

II. Disco rubente. (128-134)

128. **Helotium rubens** Sacc. et Roum. Reliq. Libert. IV, n. 40. — 979
Ascomatibus gregariis, sessilibus ceraceo-tenacellis prorsus super-
ficialibus, plano-scutellatis, $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. diam., ambitu sinuatis, disco
roseo persicino velutino, margine exiguo pallidiore; ascis clavatis,
longe sensim stipitatis, 85-100 \approx 8, paraphysatis octosporis; sporidiis

distichis bacillari-fusoideis, rectis v. curvulis, 18-22 \approx 2-3, hyalinis, nubilosis.

Hab. in cortice Rhamni in Arduennis. — *Mollisia rameali* Karst. subaffine videtur. An *Pezizella*?

129. **Helotium subcarneum** (Schum.) Sacc. Mich. II, p. 260, F. ital. 980 f. 1330, *Peziza subcarnea* Schum. in Flor. dan. vol. XII, t. 2084 fig. 1, *P. carnea* Fr. S. M. II, p. 135, *Helotium carneum* Fr. Summa p. 356? — Gregarium, pusillum obconico-sessile, totum sordide rubrum, concaviusculum, udum $1/2$ -1 mm. diam., in sicco valde contractum; ascis tereti-clavulatis, 60 \approx 6; sporidiis octonis subdistichis, tereti-fusoideis, curvulis, 12 \approx 3, hyalinis (rarius plene evolutis).

Hab. in fragmentis ligneis et strobilis Pinorum dejectis in Horto Patavino (BIZZOZERO), ad lignum Fagi, Carpini etc. in Suecia et Saellandia.

130. **Helotium gyalectoides** (Saut.) Sacc. *Peziza gyalectoides* Saut. 981 Pilz. Salzb. p. 14, Wint. Hedw. 1881, pag. 133. — Ascomatibus gregariis, minutis, orbicularibus, sessilibus ceraceis, concaviusculis, roseolis, margine integro pallidiore; ascis clavatis, 50-60 \approx 7, octosporis; sporidiis inordinatis, oblongo-fusoideis, saepe inaequilateribus, continuis, 8-10 \approx 2, hyalinis; paraphysibus filiformibus.

Hab. in foliis putridis Salicis Salzburg. — Habitus *Gyalectaria fareolaris*.

131. **Helotium querceti** Sacc. Mich. I, p. 68, F. ital. f. 117. — Cupulis sessilibus, centro punctiformi adfixis, applanatis, scutellaribus, vix $1/4$ mm. diam., levibus, glabris, siccis quandoque subcontortis sed apertis, ex albido roseis; ascis cylindraceo-clavatis, breve stipitatis, 50-55 \approx 7-8; paraphysibus crassiuscule filiformibus obvallatis, octosporis; sporidiis distichis cylindraceo-fusoideis subbacillaribus, 12-14 \approx 1 $1/2$ -2, bi-trinucleatis indeque spurie 1-2-septatis, hyalinis.

Hab. in foliorum pag. inf. Quercus pedunculatae in silva Montello Ital. bor. — *Helotio epiphylo* (Pers.) Fr. subaffine, at rite distinctum.

132. **Helotium nemorosum** Cr. Fl. Finist. p. 47. — Ascomate subhemisphærico, sessili, rubescente, 1-2 mm.; ascis subclavatis, octosporis; sporidiis fusoideo-elongatis, angustis, granulosis.

Hab. ad terram in nemorosis ad vias Finistère Galliæ. — Contextus anastomotico-filamentosus.

133. **Helotium roseo-rubrum** Cr. Fl. Finist. p. 48. — Ascomate hemi-

sphærico vel turbinato, sessili, roseo-rubro, 1-3 mm.; ascis subcylindricis, octosporis; sporidiis ovoideis, uninucleatis; paraphysibus filiformibus, apice clavulato-incurvis.

Hab. ad terram limosam Finistère Galliæ.

134. **Helotium Guernisaci** Cr. Fl. Finist. p. 48. — Ascomate turbi- 985
nato, albo-carneo vel brunneolo, carnosulo, glabro, leniter concavo,
dein convexo, 1-2 cm. diam.; ascis subcylindricis, octosporis; spo-
ridiis oblongis quadriguttulatis; paraphysibus filiformibus.

Hab. ad terram, fimo cuniculorum immixtam, Brest Galliæ (DE
GUERNISAC).

III. Disco flavo, vitellino, stramineo.

135. **Helotium immutabile** Fuck. Symb. App. I, p. 50. — Epiphyllum, 986
pallide immutabile flavum; cupulis sparsis, sessilibus vel in stipi-
tem brevissimum concolorem deorsum attenuatis, 2-3 mm. latis,
planis, acute marginatis, orbicularibus seu parum repandis; ascis
clavatis, 8-sporis, 75-88 \pm 8; sporidiis oblique monostichis, oblongo-
ovatis, rectis, continuis, hyalinis, 10-12 \pm 3-5; paraphysibus filifor-
mibus.

Hab. in foliis putrescentibus Populi tremulæ et nigræ in
Germania et Italia.

136. **Helotium Ilicis** Phill. Disc. p. 164. — Sessile, planum vel con- 987
cavum, ceraceum, glabrum, vitellinum, margine pallidiore obtuso
repando; ascis clavatis, apice acutis; sporidiis fusoideis biguttatis,
demum simulate uniseptatis, 6-9 \pm 2,5; paraphysibus filiformibus.

Hab. in foliis emortuis Ilicis, Shrewsbury Carlisle Britanniæ.
— Ascoma 0,5 mm. lat.

137. **Helotium phacidoides** Sacc. Misc. myc. II, p. 9. — Ascomatibus 988
sparsis, sessilibus plano-scutellatis, initio epidermide velatis (pha-
cidioideis), dein prorsus emergentibus, $\frac{1}{3}$ - $\frac{1}{2}$ mm. d., margine bre-
viter elevato, atro-fusco, levi, disco sordide cerineo, concaviusculo;
ascis tereti-clavulatis, breve crasse stipitatis, 60 \pm 10, paraphysatis,
octosporis; sporidiis oblique monostichis, bacillaribus, apice obtu-
siusculis, 15 \pm 2, leniter curvis, hyalinis.

Hab. in foliis emortuis Betulæ albæ, saepius epiphyllum, Mal-
medy (LIBERT). — Hic *Trochilis*, hinc *Phacidiis* affine, sed proprius
Helotio.

138. **Helotium callorioides** Rehm Ascom. Alp. n. 3. — Gregarium, 989
sessile, patellare, vix stipitatum, planum, aurantiaco-luteum, siccum
concavulum humectatum ad marginem pallidius, planum v. convexu-

lum, prosenchymatice flavide contextum, c. 0,8-1 mm. diam.; ascis clavatis, apice incrassatis, 8-sporis, 60 ± 7 ; sporidiis obtuse fusiformibus, rectis hyalinis, distichis 12 ± 3 ; paraphysibus filiformibus, apice hamatis, haud clavatis.

Hab. ad caules putridos Aconiti varieg. prope glaciem æternam, Kartel-Gletscher in valle Moos alpium Tyrolensium (BRITZELMAYR).

139. **Helotium citrinicolor** Cr. Fin. pag. 48. — Ascomate sessili, citrino, 2-4 mm. diam., margine pallidiore; ascis cylindraceis octosporis; sporidiis oblongis 4-guttatis; paraphysibus filiformibus. 990

Hab. in sarmenis emortuis Rubi, Finistère Galliæ.

140. **Helotium humile** Sacc. Mich. II, p. 78 (nec Desm.). — Ascomati- 991 bus gregariis, subsuperficialibus sessilibus plano-scutellatis, $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam., levibus, glabris, distincte marginatis, ex albo luteolis, disco concaviusculo, sicco aperto; ascis cylindraceis, 70-80 \pm 8, stipitatis, apice obtusiusculis, paraphysatis, octosporis; sporidiis cylindraceo-fusoideis, teniter curvis, 15 ± 2 -3, obtusiusculis, hyalinis.

Hab. in ramulis Genistæ scopariæ inque capsulis *Aesculi* in Normandia Galliæ. — *H. Humile* (Lasch.), *H. albello* (With.) et *H. separabilis* Karst. subaffinis, sed rite distinguenda species.

141. **Helotium Libertianum** Sacc. et Roum. Reliq. Libert. IV, n. 39. 992 — Ascomatibus subsparsis, sessilibus, scutellato-urceolatis, ceraeis, aurantiis, $\frac{1}{2}$ mm. diam.; ascis tereti-clavulatis, paraphysatis, octosporis, longiuscule sensim stipitatis, $40\cdot58 \pm 5$; sporidiis oblique 1-stichis, oblongo-cylindraceis, curvulis, 6 ± 2 , hyalinis.

Hab. in squamis strobilorum Pini silvestris. — A *Pez. chrysostigmate* differt ascomatibus multo majoribus, levibus.

142. **Helotium aureolum** Sacc. Mich. II, p. 79, nec Rab. — Ascomati- 993 bus gregariis, minutissimis, vix $\frac{1}{10}$ mm. d., plano-scutellaribus, distincte breve marginatis, semper apertis, brevissime stipitatis (simulate sessilibus) amœne citrinis, disco nitidiore; contextu excipuli subprosenchymatico flavo; ascis clavulatis, breviter stipitatis, apice rotundatis, $40\cdot45 \pm 8$; paraphysibus filiformibus obvallatis, octosporis; sporidiis fusoideis, curvulis, utrinque v. magis deorsum, acutiusculis, $14\cdot15 \pm 1\frac{3}{4}\cdot2$, 2-3-nucleatis, hyalinis.

Hab. in stipitibus Pteridis aquilinæ. — Cum nullâ formâ mihi notâ commutanda species, nisi cum *H. citrinulo* Karst, quod sporidiis brevioribus, cupulis prorsus sessilibus recedere videtur. A *Pez. chrysostigmate* ascomatibus glabris, sporidiis fere duplo longioribus etc. recedit.

143. **Helotium rhytidodes** B. et C. Cuban Fung. n. 700. — Sessile, 994
stramineum; cupula crateriformi, subtus rugoso-plicata; sporidiis
ellipticis levibus.

Hab. in insula Cuba.

IV. Disco subvirente.

144. **Helotium collinum** Speg. Mich. I, p. 473. — Ascomatibus laxe 995
gregariis, primo subglobosis dein explanato-patellaribus, sessilibus,
extus glabris, sordide virescentibus, disco amœne griseo-cæruleo,
subhygrophano, $1\frac{1}{4}$ - $1\frac{1}{2}$ mm. diam.; ascis cylindraceis, apophysatis, 45-50 \approx 4-5, obturaculo vel apice jodi ope dilute cœrulecente;
sporidiis distichis cylindraceo-fusoideis subclavulatisque, quandoque
utrinque uniguttulatis, hyalinis, 10 \approx 1- $1\frac{1}{2}$.

Hab. in pagina inferiore foliorum Viburni Lantanæ, Conegliano Ital. bor.

145. **Helotium conglomeratum** Cr. Fin. p. 48. — Ascomatibus brunneo-virentibus, 996 sessilibus, subhemisphæricis, sæpe glomerulatis;
ascis fusoideis, octosporis; sporidiis piriformibus, hyalinis, 5-guttulatis; paraphysibus filiformibus.

Hab. in sarmentis emortuis Rubi, Finistère Galliæ. — Contextus dense cellulosus.

146. **Helotium flavo-viride** Cr. Fl. Finist. pag. 47. — Ascomate 1-2 997
mm. diam., flavo-viridi, sessili, plano v. convexo; ascis claviformibus, octosporis; sporidiis ovoideis, granulosis; paraphysibus capillaribus.

Hab. ad truncos putrescentes pineos, Finistère Galliæ.

147. **Helotium albo-virens** Cooke Disc. U. S. I, p. 299. — Sparsum 998
v. subgregarium, viridi-album, sessile, subtus filis arachnoideis alabis
affixum; ascomate convexo demum subplano, sicco obscuriore
1 mm. diam.; ascis cylindraceis; sporidiis teretibus obtusis, rectis
v. curvulis, continuis, 18 \approx 3.

Hab. in ligno acerino, New Jersey Amer. bor. (ELLIS).

V. Disco lutescente, ochraceo, fulvo, melleo, alutaceo.

148. **Helotium Humuli** (Lasch) De Not. Comm. I, p. 379, Karst. M. 999
F. I, p. 112, Sacc. F. ital. f. 1361, *Peziza Humuli* Lasch in Rab.
Herb. Myc. ed. II, p. 221 et ed. I, p. 630, *Pez. humilis* Desm. XIV
Notic. n. 84. — Sessile aut subsessile, planiusculum, minutum, lutescens;
ascis cylindraceo-clavatis 70-80 \approx 9-10, obturaculo jodo leviter cœrulecente; sporidiis aciculari-elongatis, 2-3-guttulatis v.

spurie tenuiter uniseptatis, rectis vel leniter curvulis, 16-18 \approx 4, radius 12-15 \approx 3.

Hab. ad caules Humuli Lupuli sicos in Germania, Fennia, Italia, Gallia, Britannia. — Ab *H. virgultorum* atque formis ejus numerosis ascomatibus sessilibus vel fere sessilibus minoribusque mox distinguendum.

149. **Helotium Iridis** Cr. Fl. Finist. pag. 48. — Ascomate 2-3 mm. 1000 diam., extus griseo-brunneo, sessili, hemisphærico, concavo, dein expanso et convexo, margine undulato atro-fimbriato; disco griseo; ascis clavatis, octosporis; sporidiis longis, subcylindraceis, hyalinis, guttulatis; paraphysibus filiformibus, apice incrassatis.

Hab. ad caules emortuos et submersos Iridis Pseudo-acori, Finistère Galliæ. — Excipulum e cellulis hexagoniis, minutis efformatum.

150. **Helotium brevisporium** C. et Phil. Grev. VIII, p. 63. — Ochra-1001 ceum, sessile, sparsum 1-3 mm. lat., applanatum concavum, margine pallido elevato; ascis cylindricis; sporidiis breviter ellipticis, demum (spurie?) uniseptatis 7-10 \approx 3-4; paraphysibus linearibus.

Hab. ad ramos decorticatos, Waitaki Nov. Zelandiæ.

151. **Helotium melleum** B. et Br. Fung. of Ceyl. n. 957 (1873). — 1002 Punctiforme, gelatinosum, subsessile primo planum mox convexum, pallide melleum; sporidiis ellipticis, 20 μ . longis.

Hab. in corticibus, Peradenya Ceylon.

152. **Helotium discolor** (Hedw.) Sacc. *P. discolor* (Hedw.) Fr. S. M. 1003 II, p. 134, *Octosp. discolor* Hedw. Musc. II, p. 65, t. XXII, f. C. — Planum, adpressum, siccum ochraceum, humidum griseo-expal-lens, basi minima fusco-nigricante.

Hab. in cortice arborum Hannoveræ (EHRHART). — E basi, vix stipitiformi, mox expansum in cupulam orbicularem, tenuem, margine in siccis adparente, in humidis evanido, 1 mm. latam; ex habitu potius *Patelleæ* species.

153. **Helotium propinquum** Sacc. et Ell. Mich. II, p. 572, Fung. ital. 1004 fig. 1358. — Ascomatibus sparsis, sessilibus ceraceis patellaribus levibus, lutescentibus, $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. diam., disco plano-concaviusculo; ascis teretiusculis, 50-60 \approx 6-7; paraphysibus filiformibus obvalla-tis, breve stipitatis; sporidiis tereti-fusoideis utrinque obtusiuscu-lis, leniter curvulis, hyalinis.

Hab. in ramis corticatis Corni in Pensylvania (ELLIS). — Hinc *Helotio albello*, hinc *H. insititio* affine, sed satis diversum.

154. **Helotium simile** Sacc. et Roum. Reliq. Lib. IV, n. 41. — Asco-1005

matibus gregariis sessilibus ceraceo-tenacellis, superficialibus, plano-scutellatis, sinuatis, vix $\frac{1}{2}$ mm. diam., disco ochraceo-succineis, margine exiguo pallidiore impolito; ascis tereti-clavatis, breve crasseque stipitatis, octosporis, paraphysatis, 60-70 \approx 6; sporidiis bacillari-clavulatis, 14-15 \approx 3, deorsum tenuatis, hyalinis.

Hab. in ramis corticatis salicinis in Arduennis (LIBERT).

155. **Helotium sarmentorum** De Not. Prof. Discom. in Comm. critt. 1006 I, p. 379, Michel. I, p. 67. — Cupulis subsessilibus convexulis, rufo-aurantiacis; ascis clavatis 90-100 \approx 15, deorsum attenuatis, paraphysibus filiformibus obvallatis, octosporis; sporidiis ovato-oblongis, utrinque acutiusculis, sæpe inæquilateralibus, 20-22 \approx 6-7, 2-4-guttatis, hyalinis.

Hab. in sarmentis emortuis Vitis viniferæ, Milano et Conegliano Ital. bor.

156. **Helotium disciforme** Fr. Summa p. 356, *Peziza* Fr. Syst. Myc. 1007 II, p. 134. — Subsessile, carnosо-ceraceum, glabrum, planiusculum, margine lacerato, disco brunneo.

Hab. ad ramos putridos Fagi in Suecia. — Magnum, 4-6 mm. latum, subinde spurie stipitatum, subtus in adultis planum, pallidum et discus convexulus, margine vix prominente.

157. **Helotium Lens** Speg. Fung. Arg. Pug. IV, n. 227. — Sessile, 1008 hinc inde laxissime gregarium, udum hemisphærico-lenticulare, siccum applanato-patellare non v. vix contractulum, glaberrimum, pallide et sordide ceraceo-testaceum 1-1,5 mm. diam.; ascis clavulatis, sursum acutiuscule rotundatis crasseque tunicatis, deorsum parce attenuato-pedicellatis, 80-90 \approx 8-10, octosporis; paraphysibus filiformibus non v. vix apice incrassatulis obvallatis; sporidiis oblique monostichis v. distichis, utrinque subtruncato-rotundatis 15 \approx 3, continuis, 2 v. 4-guttulatis, hyalinis; obturaculo ascorum obsolete cœrulecente.

Hab. ad folia dejecta putrescentia Eryngii pandanifolii in uliginosis, Boca del Riachuelo Argentinæ.

158. **Helotium trabinellum** Karst. M. F. I, pag. 126. — Gregarium v. confertum, passim confluens, sessile crassiusculum, subplanum, testaceo-pallidum, siccum concaviusculum, incarnato- v. fulvescenti-testaceum, 0,5 mm. lat.; ascis clavatis, 65-80 \approx 8-10, obturaculo minuto jodo obsolete cœrulecente; sporidiis oblongato-ellipsoideis v. oblongatis, 2-guttulatis, rectis v. subrectis, 9-12 \approx 3-4; paraphysibus sat rarissimis filiformibus 2 μ . cr., apicem versus leniter incrassatis.

Hab. supra lignum arborum frondosarum (*Alni?*) in Mustiala Fenniae. — ****H. italicum*** Sacc. F. it. fig. 1329 (*H. trabinellum*): ascomatibus breve obconico-stipitellatis, disco subplano, ex incarnato succineo, marginato, extus conspicue pallidioribus, $\frac{1}{2}$ - $\frac{3}{4}$ mm. diam., excipulo prosenchymatico pallido; ascis clavatis, 80-90 \approx 7-8, apice rotundatis, stipitatis, p. s. 60 μ . long.; paraphysibus clavulatis obval-latis, octosporis; sporidiis distichis oblongis v. breve fusoideis, sub-rectis, 8-10 \approx 2-3, 2-nucleatis spurieque 1-septatis, hyalinis. In li-gno putri, Conegliano Ital. borealis.

159. ***Helotium agyrioides*** Cr. Finist. p. 47. — Ascomate flavo-ochra-ceo pallido, hemisphærico, sessili, 1 mm. diam.; ascis subcylindricis, octosporis; sporidiis ovoideis, hyalinis.

Hab. in ligno putri, spongioso *Quercus*, Finistère Galliæ.

160. ***Helotium monticolum*** Berk. North Am. Fungi n. 751. — Con- 1011 gestum pallide fulvum, obovatum; disco plano; sporidiis biseriali-bus, subfusiformibus.

Hab. in ligno putri in montibus Carolinæ sup. Amer. bor.

161. ***Helotium Marchantiæ*** (Berk.) Fr. Summa p. 356, *Peziza Mar-chantiæ* Berk. Engl. Fl. V, p. 304. — Obconicum, crassiusculum pallide flavidio-brunneum, marginatum, subflexuosum, disco planiu-scule; ascis clavatis, 100-120 \approx 8-10; sporidiis ellipsoideis, 12-16 \approx 3-5.

Hab. in frondibus *Marchantiæ polymorphæ*, *M. hemi-sphæricæ*, *Preissiaæ hemisphæricæ* in Britannia, Fennia et Fuegia (forma stipite longiore puberulo, disci margine acuto). — Ascoma 1-2 mm. lat.

VI. Disco badio, castaneo, griseo, atro-fusco.

162. ***Helotium badium*** Phill. Disc. p. 167. — Gregarium, erumpens, 1013 sessile, initio turbinatum et concaviusculum, dein planum, disco badio-brunneo, margine distincto et paullo dilutiore, extus concolor glabrum; ascis late clavatis, basi subabrupte attenuatis; sporidiis oblongo-fusoideis v. subclavatis, 3-5-guttulatis, 20 \approx 5; paraphysi-bus filiformibus.

Hab. in ramis emortuis (*Salicis?*) Oxford Britanniæ (BAXTER). — Ascoma 1-2 mm. lat. Habitus *H. ferruginei*.

163. ***Helotium castaneum*** Sacc. et Ell. Mich. II, p. 572, F. ital. fig. 1014 1362. — Hypophyllum, minutissimum, castaneum; ascomatibus ses-silibus, scutellatis, levibus, $\frac{1}{4}$ mm. diam., subceraceis; ascis tere-ti-oblongis, 45 \approx 16; paraphysibus filiformibus obvallatis, brevissi-

me stipitatis, apice obtusiusculis; sporidiis subtristichis oblongo-fusoideis, 11-14 \approx 3-4, biguttulatis spurieque 1-septatis, hyalinis.

Hab. in pag. inf. foliorum *Quercus laurifoliæ*, Florida, (Doct. GEO. MARTIN).

164. **Helotium populinum** Fuck. Symb. p. 316. — Cupulis carnosis, 1015 gregariis, punctiformibus, subsessilibus, primo globosis, umbilicatis demum applanatis, margine elevato, subfuscō, integro, crasso, extus pallide griseo, disco subconcavo, concolore, siccis subturbinatis; ascis stipitatis, oblongis, 8-sporis, 36 \approx 8; sporidiis distichis, oblongo-lanceolatis, hyalinis, 9 \approx 4.

Hab. in pag. super. foliorum *Populi pyramidalis* in Münchau pr. Hattenheim.

165. **Helotium lividum** Mont. Fl. Chil. VII, p. 407, Syll. Crypt. n. 630. 1016 — Hypophyllum, erumpens; cupula subsessili tandem explanata, marginata, extus glabra fuscescente, disco pallido tandem livido-fusco; ascis clavatis; sporidiis linearī-oblongis, hyalinis, 7-8 \approx 2-3.

Hab. in pagina inferiore foliorum *Æx toxici punctati*, in provinciis Chiles meridionalis (CL. GAY). — Ascoma $1\frac{1}{2}$ -1 mm. lat.

166. **Helotium microscopicum** (Wallr.) Sacc. *P. microscopicā* Wallr. 1017 Fl. crypt. n. 2507. — Ascomatibus omnium minutissimis conser-
tim sparsis, glabris, in basin stipitiformem attractis s. obconicis tur-
binatisve fuscis, dein vertice impressis, veluti in discum punctifor-
mem concolorem obtuse marginatum impressis.

Hab. ad frustula ligni faginei cœsi in Thuringia.

167. **Helotium aglæosporum** Berk. North Am. Fungi n. 752. — Cor- 1018
nicolor, irregulare; hymenio pulvinato; ascis linearibus; sporidiis oblongis curvulis utrinque hyalinis, 8-15 μ . long.

Hab. ad lignum putre in Amer. bor.

168. **Helotium stigmaion** Rehm Ascom. Alp. n. 767, Hedw. 1882, 1019
n. 7. — Gregarium, in foliis dealbatis sessile, conico-truncatum v.
subglobosum, apice deplanatum, orbiculare, siccum leviter urceo-
latum, nigro-fuscum, humidum planum, margine tenuissimo cinctum,
disco ferrugineo-nigro, c. 0,3 mm. diam.; prosenchymatico-fusco-
contextum; ascis clavatis, 8-sporis, 45 \approx 6, jodi ope cœrulecenti-
bus; sporidiis elongatis, obtusis, hyalinis, 6 \approx 2 (haud rite evolu-
tis visis); paraphysibus filiformibus, ascos superantibus haud distin-
cte clavatis; jodi ope apice ascorum cœrulecente.

Hab. ad folia graminea sicca prope Sölden in valle Ötz al-
pium Tyrolensis (DR. REHM). — Var. **minusculum** Rehm Ascom.
n. 767: ascomate minuto, subgloboso; ascis clavatis, 45 \approx 7-8; spo-

ridiis 2-guttatis, oblongo-ellipsoideis, 0.9×2.5 ; paraphysibus aequalibus. In vaginis siccis graminum in Tyrolia.

Subgen. *Calycella* Sacc. Ascoma extus fursfuraceum vel puberulum (Cfr. *Lachnella*).

A. Stipitata.

169. **Helotium alutaceum** B. et Br. Fung. of Ceyl. n. 956. — Sessile, 1020 subtus margineque albidum, flexuosum, pulverulentum; hymenio alutaceo; sporidiis ellipticis, 12-13 μ . long.

Hab. ad ramos, Peradenya Ceylon. — Margo initio inflexus.

170. **Helotium lætum** (Boud.) Sacc., *Pachydisca læta* Boud. Bull. Soc. Myc. IV, p. 78, t. 16, f. 3. — Stipitatum carnosum, luteo-aurantiacum, 2,5 mm. latum, 1-2 mm. altum, lentiforme, crassum, margine vix prominente leve aut sub-crenulato, extus minutissime fursfuraceum, hymenio primo piano, dein convexo, ascis prominulis granuloso, concolore; paraphysibus non aut vix incrassatis, simplicibus, cylindricis, hyalinis aut lutescentibus parce septatis, intus granulosis, 3-4 μ . crassis; ascis cylindrico-clavatis, octosporis, 170-180 \approx 10 vacuis, foramine marginato crenulato donatis; sporidiis oblongo-fusoideis, inaequilateralibus et subflexuosis, intus utrinque granulosis, centro libero, 24-30 \approx 5-7.

Hab. ad cortices madidos in paludosis Montmorency Galliæ.

171. **Helotium microspermum** Speg. Fungi Fueg. n. 334. — Cupulis 1022 in ligno denudato hinc inde dense gregariis, majusculis, firmis, culpulato-expansis, 2-3 mm. diam., sessilibus v. subsessilibus, disco concaviusculo sordide rufescente v. melleo, margine acuto integer-rimo extus sordide fuscis minutissime albo-pruinulosis; ascis cylindraceis v. subclavulatis, sursum leniter attenuatis, apice obtuse rotundatis crassiuscule tunicatis, deorsum longiuscule crasseque attenuato-pedicellatis, 40-45 \approx 2-3, octosporis; paraphysibus paucis non v. vix longioribus obvallatis; sporidiis recte v. oblique mono- v. di-stichis, ellipticis utrinque subacutato-rotundatis, exiguis, 3-3,5 \approx 1,5-1,8, minute biguttulatis, hyalinis.

Hab. in ramulis emortuis decorticatis fagineis in silvis pr. Ushuvaia Argentinæ.

172. **Helotium maculosum** Ell. et Mart. in Amer. naturalist 1883, 1023 p. 1284. — Ascomate orbiculari, sessili, $1\frac{1}{6}$ mm. diam., piano vel convexo, sicco concavo, disco sordide carneo, extus obscuriore, pilis septatis brunneis prope basin oriundis vestito; ascis oblongo-cla-

vatis, 55 ± 12 ; sporidiis distichis, late fusoideis, $16-20 \pm 4-5$, plasmate tripartito.

Hab. in foliis viventibus brunneo-maculatis *Perseæ palustris*
New Florida Amer. bor. «An *Trichopeziza?*».

173. **Helotium Stellerii** (Weinm.) Sacc. *Pez. Stellerii* Weinm. Hym. 1024
Gaster. p. 478. — Sparsum v. confertum, sessile, carnoso-tremel-
losum, ochraceum, applanatum, extus tenerrime granulosum, mar-
gine nunc integro, nunc repando.

Hab. in lignis pineis mucidis semel in Rossia. — Pusillum.
Ascoma primo globosum, margine involuto, granuloso, mox applana-
tum, margine aut integro aut repando-crispato, et in hoc statu
totius evolutionis vix 1 mm. latum.

B. Sessilia v. subsessilia.

174. **Helotium sambucinum** Cr. Fl. Finist. p. 49. — Ascomate pedi- 1025
cellato, concavo, dein convexo, sordide albo, subtus tomentello,
2-3 mm., stipite subcylindrico brunneo; ascis clavatis; sporidiis
piriformibus, 4-guttulatis; paraphysibus filiformibus, granulosis,
brunneolis.

Hab. in ramis emortuis Sambuci Finistère Galliæ. — «An
Dasyscypha?»

175. **Helotium macrosporium** Peck 26, Rep. p. 82 (1874), nec Sacc. 1026
— Ascomatibus primo subclausis, dein expansis et leniter concavis
albidis, extus furfuraceis; stipite brevi, crassiusculo; sporidiis ob-
longis, grosse uninucleatis, $25-30 \mu$. longis.

Hab. in ligno emortuis fagineo Worcester Americæ borealis.
— Fungus siccando leniter brunnescit.

176. **Helotium Laburni** B. et Br. in Phill. Disc. pag. 135 et Bresad. 1027
Trid. pag. 69, t. 74, f. 2. — Ascomate leviter stipitato v. sessili,
concavo v. plano, extus furfuraceo pallido, margine infexo, disco
ochraceo, vivo; ascis cylindraceo-clavatis; sporidiis fusoideis, sub-
acuteis v. clavatis, biquadriguttatis, $17-27 \pm 5-9$; paraphysibus fili-
formibus obtusiusculis.

Hab. in ramis demum decorticatis *Cytisi Laburni* in Britan-
nia et agro tridentino. — Ascoma $\frac{1}{2}-2$ mm. diam. Caro albo-fla-
vida. Ascii, teste Bresad. $120 \pm 10-12$, sporidia $23 \pm 7-8$. *Helot. ci-
trino* affine.

177. **Helotium pateræforme** (Berk.) Cooke Austr. Fung. p. 51, *Pe- 1028*
ziza pateræformis Berk. Fl. Tasm. p. 276. — Ochraceum; cupu-
la sessili, sublobata, concava, subtus rugosiuscula subtiliter to-

mentosa; ascis linearibus; sporidiis oblongis utrinque attenuatis subcymbæformibus, 30μ . longis.

Hab. in ligno musoso in Tasmania (ARCHER). — *Hel. citrino* affine. Ascoma 4 mm. lat.

178. **Helotium pileatum** Peck 28 Rep. pag. 67, t. I, f. 11-14 et 32 1029
Rep. p. 58. — Ascomate subhemisphærico v. pileiformi, stipitato,
glabro, albido subtus appланato, pruinuloso; stipite longiusculo albo-
pruinoso; sporidiis oblongis, hyalinis, 10-12 μ . long.

Hab. in caulis herbaceis emortuis in uliginosis Hunter Greene
county Am. bor. — Totus fungus 12 mm. altus. Agaricum minu-
tum simulans.

179. **Helotium sulphurinum** Quél. 12, Suppl. p. 13, t. VII, fig. 11 et 1030
in Grevillea 1879, pag. 116. — Ascomate cyathideo, 0,5-1 mm.
diam., substipitato, tenero, pruinoso, albo v. citrino; hymenio con-
cavo, nitide sulphureo, udo albescente; sporidiis fusiformibus, 12
 μ . longis, 4-guttulatis, rectis v. incurvatis.

Hab. gregarie in ramis exsiccatis Coryli in Jura. — Affine
H. citrino.

180. **Helotium rhizogenum** E. et E. Journ. Mycol. 1888, pag. 100. — 1031
Ascomate stipitato, pallido, dein lète flavo; stipite cylindrico, cur-
vulo, 2-4 mm. alto, 0,5-1 mm. crasso, pallido et granuloso-pube-
rulo, disco leniter concavo, 1-2 mm. lato, pallide flavo, pallidius
marginato; ascis angustis, 65-75 \approx 5, filiformi-paraphysatis; spori-
diis distichis, clavato-fusoideis, 6-8 \approx 1,3-1,5, hyalinis.

Hab. ad radices expositas Andropogonis, Newfield
N. J. Amer. bor.

181. **Helotium fibuliforme** (Bolt.) Berk. Outl. p. 371, *Helv. fibulifor-* 1032
mis Bolt. fung. t. 176, *Pez. Fibula* Pers. syn. p. 660, *Pez. fibu-*
liformis Fr. Syst. Myc. II, p. 155. — Firmum, capitulo convexo
flavo subtus stipiteque brevi crasso subvilloso nigro-fuscis; ascis
tereti-clavatis; sporidiis oblongo-fusoideis, 2-guttatis, 9-12 \approx 2-4.

Hab. in truncis ulmeis ad latera fossarum prope Bingley et
Bettws-y-Coëd Britanniæ. — Ascoma semper apertum lentiforme,
4-6 mm. fere latum.

182. **Helotium Limonium** C. et P. Peck 26 Rep. p. 83. — Citrino- 1033
flavum, extus pallidius; ascomate plano vel leniter convexo, im-
marginato, primo extus tenuiter farinaceo; stipite gracili, æquali,
diametro ascomatis breviore; ascis cylindraceis; sporidiis cylin-
draceis, utrinque obtusis, curvulis v. rectis, 7,5 μ . longis.

Hab. ad caules herbarum, Center Amer. borealis.

183. ***Helotium rhizophilum*** (Fuck.) Cooke Handb. n. 2155, Pat. Tab. 1034 anal. fig. 590, *Ciboria rhizophila* Fuck. Symb. mycol. p. 312. — Cupulis primo infundibuliformibus, dein applanatis, excavatisque, usque ad 4 mm. latis, immarginatis, extus et stipite 7-10 mm. longo albicante-furfuraceis, disco vitellino; ascis linearibus, utrinque attenuatis, stipitatis, 8-sporis, 60 = 6; sporidiis subdistichis, continuis, cylindraceis, rectis, hyalinis, 12 = 2.

Hab. in rhizomatibus Koeleriæ glaucæ aliorumque graminum in silvaticis Rhenogoviae, Galliae, Belgii et Britanniæ. — Ex iconе Pat. l. c., sporidia clavulata; furfur ascomatis e pilis brevissimis, 1-septatis, hyalinis constat.

184. ***Helotium furfuraceum*** Phill. et Harkn. Calif. p. 4. — Sparsum, 1035 minutum, breve stipitatum vel subsessile, cyathiforme, alutaceum, furfuraceum; ascis clavatis, 8-sporis, apophysatis; sporidiis oblongo-ellipsoideis, leniter curvis, 2-3-guttatis, 15 = 5.

Hab. hypophyllum in foliis Quercus agrifoliæ in California.

185. ***Helotium discretum*** Karst. M. F. I, p. 116. — Gregarium, stipe 1036 pitatum; cupula planiuscula, levissime puberula, albida v. lutescenti, disco subluteo, sicco fulvo-flavo, 0,4-0,8 lat.; stipite æquali, 0,2 mm. alt.; ascis cylindraceo-clavatis, 40-48 = 4; sporidiis cylindraceis, rectis v. subrectis, 2-5-guttulatis, 4-8 = 1,5.

Hab. ad caules Humuli Lupuli exsiccatos in Mustiala Fen-niæ.

186. ***Helotium agaricicolum*** B. et Br. Fung. of Ceyl. n. 959. — Cupulis breviter stipitatis, extus pulverulentis; hymenio vinoso; sporidiis uniseriatis cymbiformibus, pallido fuscis. 1037

Hab. in Agaricis putridis, Peradenya Ceylon.

PHIALEA Fr. Obs. II, p. 305 ut subgen. [1818], (Etym. *phiale* poculum), *Helotii* species Karst. et Auct., *Pezizæ* spec. Fuck., *Hy-menoscypheæ* spec. Phillips, *Cyathiculæ* spec. De Not. — Ascoma ceraceo-membranaceum, tenue, initio urceolato-subclausum, mox apertum concavum vel convexum, glabrum vel pruinatum, longiuscule tenuiter stipitatum, margine non dentatum. Ascii elongati v. lineares, 8-spori, filiformi-paraphysati. Sporidia ovoidea, oblonga vel teretiuscula, continua, hyalina. — Sensim hinc ad *Helotium*, illinc ad *Pezizellam* transit.

A. Eu-Phialea. Ascoma glabrum, nudum.

I. Disco albo, albido, griseo.

1. ***Phialea cyathoidea*** (Bull.) Gill. Disc. p. 106, *Peziza cyathoidea* 1038

Bull. t. 316, f. 2, Fr. Syst. Myc. II, p. 124, *Fungoides* etc. Mich. gen. p. 205, n. 14, *Elv. infundibuliformis* Scop. Carn. II, p. 480, *P. Hirudo* Batsch f. 149? *Octosp. albidula* Hedw. l. c. pag. 30, t. 9, f. B, *P. tenella* Batsch Cont. I, f. 151, Nees Syst. f. 294, *P. tenerrima* Holmsk. Ot. II, p. 33, t. 11, *Helotium cyathoideum* (Bull.) Karst. M. F. II, p. 136, *Cyathicula vulgaris* De Not. — Tenuis, albida, pallida; cupula globoso-cyathiformi, dein explanata integerrima; stipite filiformi longiusculo; ascis clavatis, 45 ± 5 , filiformi-paraphysatis; sporidiis oblongis, rectis, curvulisve, $6-10 \pm 1,5-2$, hyalinis, saepe 2-guttatis.

Hab. ad stipites et caules plantarum *Epilobii*, *Dipsaci*, *Aconiti*, *Heraclei*, *Cicutæ*, *Eupatoriæ*, *Nesææ*, *Ballotæ* etc., in Suecia, Gallia, Britannia, Italia, Belgio, Fennia, Germania, Hungaria, Nova Zelandia, Amer. bor. vulgatissima. — Stipes longus, cupula demum planiuscula et tenuis; color in humida stirpe aquose pallidus, albidus, lutescens, subincarnatus; in siccis obscurior. Bulliardus tres notat varietates, albam, luteam, ferrugineam. Sec. Karsten *Ph. albidula* (Hedw.) recedit sporidiis paullo majoribus, 11 ± 2 . — β . **convivalis** Batsch, Pers. Myc. Eur. p. 286: epiphylla, albida, cupula turbinata, stipite breviore. Cum *H. epiphyllum* non commutanda. Ad folia locis humidis. — γ . **violascens** A. et S.: epiphylla, amœne violascens. Distincta forsitan. Iisdem locis, autumno. Ex hac regione quoque Mich. t. 86, f. 13 et *P. pedicellata* Sowerb. t. 369, f. 1.

2. **Phialea Solani** (Pers.) Sacc. *Pez. Solani* Pers. Obs. 2, p. 80, Fuck. 1039
Symb. p. 307. — Cyathiformis, stipitata, firmula, extus longitudinaliter striata, pallida; ascis anguste linearibus, oblique stipitatis, 8-sporis, 52 ± 4 ; sporidiis elongato-distichis, cylindraceis, rectis, hyalinis, 8 ± 2 .

Hab. in caulis putridis Solani tuberosi, in Germania et Britannia. — Media inter *Ph. cyathoideam* et *Ph. Urticæ*.

3. **Phialea clavata** (Pers.). Gill. Disc. p. 110, *H. clavatum* (Pers.) 1040
Karst. *Peziza clavata* Pers. Myc. Eur. I, p. 283, Fr. Syst. Myc. II, p. 122. — Sparsa vel gregaria, stipitata, firmula, pallescens; cupula cyathoidea, levi, 0,6-0,8 mm. lat.; stipite æquali, crassiusculo, circiter 0,6 mm. alt.; ascis cylindraceo-clavatis, $40-50 \pm 4-5$, obturaculo jodo leviter cœrulecente; sporidiis oblongatis aut fusoideo-oblongatis, $4-8 \pm 1,5-2,5$.

Hab. ad caules Umbelliferarum e. g. *Angelicæ* et *Fi-*

licum in Fennia, Lapponia et Britannia, in caule Teucrii in Rhe-nogovia. — *Ph. caulicolæ* affinis.

4. **Phialea Urticæ** (Pers.) Sacc. *H. Urticæ* (Pers.) Karst. M. F. I, 1041 p. 133, *Peziza Urticæ* Pers. Myc. Eur. I, p. 286, *P. striata* Fr. Syst. Myc. II, p. 122, *Hymenoscypha* Phill. Disc. p. 141. — Gre-garia, stipitata, firmula; cupula turbinata, humida longitudinaliter conferte striatula, pallescenti, sicca fuscescenti aut fusca, margine connivente, leviter pruinosa, circiter 0,5 mm. lat.; stipite æquali levi, crassiusculo, firmo, gilvo-pallido, cupula breviore, subinde lon-giusculo; ascis cylindraceo-clavatis, 40–50 \approx 5–6, obturaculo jodo obsoletissime cœrulescente; sporidiis fuosoideo-oblongatis, 7–8 \approx 2.

Hab. ad caules Urticæ dioicæ exsiccatos in Suecia, Ger-mania, Fennia, Britannia. — Sec. Rehm vix differt a *Ph. Cacaliæ*.

5. **Phialea scyphiformis** (Wallr.) Sacc. *Pez. scyphiformis* Wallr. Fl. crypt. n. 2581. — Ascomatibus exiguis sparsis tenuiter membra-naceis, albo-pallescentibus, extus fibris tenerrimis adpressis parcis æqualiter adspersis, primum e basi stipitiformi firma crassa sub-clavato-turbinatis sensim osculo rotundo hiantibus, dein in discum concavum concolorem reclusis abrupte scyphiformibus, siccando margine inflexo subtili tenuissime striato inflexo conniventibus com-pressis triquetrisve.

Hab. ad caules aridos Chærophyllorum in Germania. — *Ph. Urticæ* similis, differre tamen videtur tegmine fibroso æqua-liter diffuso, nec in striae distinctas stipato, cupulis concaviuscu-lis abrupte scyphiformibus, siccando plane difformibus triquetris et stipite breviore, crassiore magisque æuali.

6. **Phialea rubicola** (Fr.). Sacc. *Hel. rubicolum* (Fr.) Fuck. Symb. 1043 p. 314, Sacc. F. it. f. 1342, *Peziza fructigena* γ. *rubicola* Fr. Syst. Myc. II, p. 119, *Hel. virgultorum rubicolum* Fr. S. v. Sc. p. 355. — Ascomate orbiculari, plano, albo-pallescente; stipite æquali; ascis elongatis (non clavatis), stipitatis, 8-sporis, 96 \approx 8; sporidiis subdistichis, oblongis, antice obtusis, basi valde attenuatis, subcurvatis, continuis, hyalinis, 20 \approx 4.

Hab. in sarmentis Rubi et Rosæ in Suecia, Germania, Italia, Gallia.

7. **Phialea Perula** (Pers.) Gill. Disc. p. 105, *P. Perula* Pers. Myc. Eur. I, p. 280, Fr. S. M. II, p. 125. — Cupula submembranacea elongata, infundibuliformi, albida, stipite brevi nigrescente.

Hab. in umbrosis uliginosis ad caules aridos Angelicæ silve-stris in Germania et Gallia. — Sparsa, forma marsupii; cupula

ore lato aperta 4 mm. alta, exsiccata nigrescens, substantia tenuis-sima.

8. **Phialea albida** (R. et D.) Gill. Disc. p. 105, *Peziza albida* Rob. 1045 et Desm. 19, Not. p. 28, *Helotium* Pat. Tab. 382. — Sparsa aut laxe gregaria, stipitata, fragilis; cupula minuta, eburnea, acetabu-liformi dein plana, margine integerrimo; stipite crassiusculo basi rufo; ascis clavatis, 100 μ . long.; sporidiis hyalinis, oblongis, ob-tutis, subcylindricis, 15-17 \times 4, 2-guttatis.

Hab. in petiolis semiputridis Fraxini, in Gallia, Britannia, Belgio. — Ascoma 1-3 mm. lat.; stipes subæquilongus. — Var. **Æsculi** Phill. Disc. p. 138: a typo differt sporidiis majoribus 20-23 \times 4-5 et saepius clavulatis. In petiolis Æsculi in Britannia.

9. **Phialea phyllophilæ** (Desm.) Gill. Disc. p. 105, *Helotium* Sacc. F. 1046 it. fig. 1341, *Pez. phyllophila* Desm. IX, Not. pag. 8, *Pezicula phyllophila* Karst. M. F. I, p. 167 (?). — Amphigena, exigua, sti-pitata, glabra, candida, subnervisequa; cupula concava, dein pla-no-convexa; ascis clavatis, 60-70 μ . long.; sporidiis oblongis cur-vatis.

Hab. in foliis semiputridis Acerum et Fagorum, in Gallia et Italia. — Stipes $\frac{1}{3}$ mm. long. In forma fagicola italica: asco-mata stipitata, $\frac{1}{4}$ mm. diam., pallide straminea, glabra; ascis cla-vulati, breve stipitati, 60 \times 12, apice rotundati, octospori; sporidia anguste fusoidea, utrinque acutiuscula, leniter curva, 12-15 \times 2 $\frac{1}{2}$ -3, 2-4-guttulata, hyalina.

10. **Phialea paludum** Quél. XIII Suppl. p. 9, tab. XII, p. 26. — Asco- 1047 matibus cyathiformibus, 2-3 mm. diam., subtilibus, glabris, albe-scentibus vel carneis, vix brunneis diaphanisque; stipite filiformi, brevissimo vel elongato; hymenio albo vel pallide lutescente; spo-ridiis ellipsoideis, 10 μ . longis.

Hab. gregaria in partibus foliorum emortuorum, Alsace. — Af-finis *Ph. capillipedii*.

11. **Phialea Berggrenii** (C. et Phil.) Sacc. *Helotium* C. et Ph. Grev. 1048 VIII, pag. 63. — Phyllogena, pallida, stipitata; cupulis sparsis, cyathiformibus, 0,5-0,7 mm.; stipite gracili, æquali circa 0,5-1 mm. long.; ascis cylindraceis; sporidiis ellipticis, nucleatis, 10-13 \times 3-5; paraphysibus filiformibus.

Hab. in foliis putridis, Melbourne Australiæ.

12. **Phialea simulata** (Ellis) Sacc. *Helotium simulatum* Ellis in Bull. 1049 Torr. Bot. Club. 1881, p. 72. — Ascomate stipitato, disco circu-lari, plano vel leniter umbilicato, sordide aquoso, 3 mm. lato, te-

nui, dein (sicco) concavo; stipite 3 mm. alto, levi, leniter deorsum contracto et saturationi; ascis sessilibus, cylindricis; sporidiis ellipsoideis, hyalinis, 12 μ . longis; paraphysibus filiformibus.

Hab. in foliis *Aceris* humi fusis putridis, New Jersey Americanæ borealis.

13. **Phialea erioloma** (Fr.) Sacc. *Pez. erioloma* Fr. Syst. Myc. II, 1050 p. 121. — Cupula acetabuliformi, pallida, marginem versus subvilosa, stipite longo tenui æquali glabro.

Hab. in ramis putrescentibus Ericæ in Suecia (ASPEGREN), ad stramina putrida in Thuringia (WALLROTH). — Sparsa, 2 mm. alta vel paullo altior; discus in siccis pallidus, humectatus lutescenti-olivaceus. Ob villum prope marginem ad *Lachnellas* accedit.

14. **Phialea hirtipes** Mout. et Sacc. (sp. nov.). — Ascomatibus stipitatis, 4-5 mm. altis; stipite elato cylindraceo, sordide albo, dein fusco, basi nigro-villoso, disco concolore applanato vel convexisculo, 2 mm. diam., immarginato; ascis teretibus, tetrasporis, 80-8; sporidiis monostichis, breve oblongis, continuis, 14-5.

Hab. in ramis dejectis locis udis *Alni glutinosæ* in Belgio. — Statura fere *Ciboriæ*.

15. **Phialea pulchella** (Fuck.) Sacc. *Peziza pulchella* Fuck. Symb. 1052 p. 299. — Sparsa; cupulis glabris, minutis, candidis, hemisphæricis applanatisve, margine pallidiore, stipite cupulam æquante, basi atro, recto, siccis sordidis; ascis oblongis, 8-sporis, 26-4; sporidiis oblongo-cylindraceis, 6-2, hyalinis.

Hab. in acubus putridis udis *Pini silvestris* in Rhenogovia.

— Sec. Fuck. status conidicus est *Sphæridium candidum* Fuck.

16. **Phialea subtilis** (Fr.) Gill. Disc. p. 161, *Hel. subtile* Fr. Summa 1053 p. 354, Karst. M. F. I, p. 131, *Pez. subtilis* Fr. Syst. Myc. II, p. 157, *Hymenoscypha* Phill. Disc. p. 132, *Fungoidaster minimus*, *discoides*, *albus* etc. Mich. gen. p. 201, t. 82, f. 3, qui *H. album* Pers. Myc. Eur. p. 347, a nostro haud essentialiter differt. Altera vero species *Fungoidastri* l. c. t. 82, f. 4 (*H. incarnatum* Pers. l. c.) ad hoc genus non pertinet. — Candida, minuta, firmula; capitulo plano-convexo; stipite tenui brevi; ascis tereti-clavatis, 66-4; sporidiis elliptico-fusoideis, 5-6-2; paraphysibus filiformibus.

Hab. in foliis dejectis præcipue *Abietis excelsæ* et *pectinatae* in Suecia, Gallia, Britannia, Germania, Italia. — Non cum *P. chionea* commutanda; stipes 2 mm. haud longus, subattenuatus. Ascoma 2 mm. latum, demum leviter convexum, subtus plano-concavum.

17. **Phialea strobilina** (Fr.) Sacc. nec Gill. (quæ *Ciboria*), *H. strobilinum* (Fr.) Fuck. Symb. p. 313, *Pez. strobilina* Fr. S. M. II, p. 125, *Ombrophila* Karst. M. F. II, p. 92. — Piriformis, firma; cupula concava pallide rufescens, margine tumido integerrimo; stipite brevi nigro; ascis subcylindricis, tenuibus, sursum leniter dilatatis; sporidiis oblongo-fusoideis, 8-12 \approx 2-2.5 hyalinis; paraphysibus filiformibus.

Hab. ad strobilos abiugnos, frequens, in Suecia, Britannia, Germania, Italia, Gallia, Blue Canon Amer. bor. — Distinctissima, eximie gregaria, valde tenax, minuta, vix 2 mm. alta: passim adest subiculum floccoso-furfuraceum (*Chalara strobilina* Sacc.). Cupula junior urceolata, dein magis dilatata, sicca nigrescens, humectata expallens. — β . **opaca** Fr. l. c.: cupula olivacea, humida spadicea. Primo subglobosa, disco punctiformi; dein lenticularis, disco concavo, margine semper obtuso integerrimo. Ad conos Pini silvestris in Suecia.

18. **Phialea proximella** (Karst.) Sacc., *Helotium proximellum* Karst. 1055
M. F. I, p. 133, Sacc. F. ital. f. 1336. — Minuta, gregaria, stipitata; cupula convexa, subtus concava, virescente v. subolivascente albida, sicca subfuscata, hymenio albo v. candido, circiter 0,7 mm. lat.; stipite tenuiusculo, subæquali, stricto, firmulo, fulvescenti v. rufescens, sicco fusco v. fusco-nigrescens, latitudine cupulae breviore; ascis cylindraceo-clavatis, 28-32 \approx 4; sporidiis vulgo oblique monostichis, eguttulatis, 4-8 \approx 1-4,5; paraphysibus parcis, gracilibus.

Hab. ad acus Pini silvestris et Abietum putrescentes Mustiala Fenniae et Susegana Ital. bor. — In forma **abietina** italica: asci fasciculati, clavato-cylindracei, sæpe e basidio unico plures orientes, 35-40 \approx 4, octospori; sporidia cylindraceo-fusoidea, 5 \approx 1 $\frac{1}{2}$, hyalina.

19. **Phialea advenula** (Phill.) Sacc., *Hymenoscypha* Phill. Disc. pag. 1056
133. — Stipitata, concava vel plana, duriuscula, alba vel flavidobalta; stipite æquali, tenui; ascis tereti-clavatis; sporidiis oblongo-ellipsoideis, 8-9 \approx 2,5-4; paraphysibus parcis, filiformibus.

Hab. in foliis dejectis Laricis, North Wales Britanniae. — Habitus omnino *Phialeæ subtilis* sed sporidia fere duplo majora.

20. **Phialea Sabinæ** (Fuck.) Sacc., *Hel. Sabinæ* Fuck. Symb. mycol. 1057
p. 313. — Sparsa, piriformis demum magis dilatata, 1 mm. alta; ascis stipitatis, elongato-clavatis, 8-sporis, 108 \approx 10; sporidiis mo-

nostichis, ovatis, hyalinis, nucleo granuloso, $10 \approx 5$; paraphysibus linearibus, ascis sabbrevioribus.

Hab. in foliis et ramulis induratis Juniperi Sabinæ et communis in silva Oestrich Rhenogoviæ.

21. **Phialea juniperinella** (Karst.) Sacc. *H. juniperinellum* Karst. M. 1058 F. I, p. 133. — Gregaria, stipitata, firmula; cupula convexa, subtus concava, alba vel candida, latit. 0,2-0,3 mm.; stipite filiformi, saepe flexuoso, fusco-nigricante vel nigrescente, apice fusco-pallido, altit. 1,5-2 mm.; ascis gracilibus, cylindraceo-clavatis, 15-20 \approx 2-3; sporidiis minutissimis.

Hab. ad acus Juniperi communis putrescentes prope oppid. Tammerfors et Mustiala Fenniæ. — E longinquo adspecta facile pro specie *Stilbi* Tode sumitur, sed vera *Phialea*, potissimum *Ph. subtili* aut potius *Ph. proximellæ* proxima; stipes crassit. 60 μ . circiter.

22. **Phialea alniella** (Nyl.) Sacc. *H. alniellum* (Nyl.) Karst. Symb. 1059 p. 239, *Peziza alniella* Nyl. Obs. p. 45, Karst. Monogr. Pez. p. 140, *Hel. Grenseri* Auersw. in Rab. Fung. eur. ed. II, 1122. — Subgregaria, stipitata, alba vel pallido-alba, plana, latit. 0,4-0,8 mm.; stipite tenello brevi vel brevissimo; ascis cylindraceo-clavatis 50-55 \approx 6-7, obturaculo jodo obsolete cœrulescente; sporidiis distichis vel oblique monostichis, oblongatis, eguttulatis, 6-12 \approx 2-4.

Hab. ad amenta Alni in Fennia et in Hungaria. — Cupula primum tenuissime furfuraceo-pulverulenta, deinde subglabra, disco saepe dilute luteolo; stipes pallido-albus, tandem fuscescens.

23. **Phialea Amenti** (Batsch) Sacc. *Peziza Amenti* Batsch Cont. I, 1060 p. 211, fig. 148 (*Pez.*) *Helotium Amenti* (Batsch) Fuck. Symb. p. 313, Gill. Disc. c. ic. — Tenuis, albida; cupula convexula cicatrice centrali notata; stipite brevi sursum incrassato; ascis stipitatis, elongatis, 8-sporis, 54-72 \approx 6; sporidiis oblique submonostichis, ovato-oblongis, 6-8 \approx 3-4, hyalinis.

Hab. in amentis fructibusque Salicum et Alnorum in Germania, Gallia, Britannia, Italia. — Sparsa, minuta, tota glabra, aquose pellucida, demum fuscescens; cupulæ margo reflexus, affinitatem cum *Helotiis* indicat.

24. **Phialea chlorella** (Saut.) Sacc. *Pez. chlorella* Sauter in Flora 1061 1845, p. 133 (*Peziza*). — Gregaria, tenuis; cupula primo plana et aquose-albida, demum convexa pileiformi viridula; stipite æquali aquose-albido.

Hab. ad radices pineas pr. Aubach Austriae (SAUTER). — P.

rhodoleucæ et *P. amenti* proxima. Stipes 1 mm. circ. longus, cupula circ. 1 mm. lata.

25. **Phialea Utriculus** (Berk.) Sacc. *Peziza Utriculus* Berk. Exot. 1062

Fung. p. 396. — Sparsa; cupulis albis, hemisphæricis, membranaceis, levibus, crispis; stipite gracili, longiusculo.

Hab. in lignis putrescentibus, Juan Fernandez (BERTERO). — *Belonid.* *Campanulæ* similis sed minor.

26. **Phialea concolor** (Phill.) Sacc. *Peziza concolor* Phill. in Stev. 1063

Myc. Scot. p. 321, *Hymenoscypha* Phill. Disc. p. 139. — Gregaria, minuta, stipitata, pallida v. albida, firma, tenuis, hemisphærica, rugulosa, margine minuto fimbriato; ascis cylindraceo-clavatis, octosporis; sporidiis oblongis, utrinque rotundatis, 8-10 \approx 2-3; paraphysibus filiformibus.

Hab. in ligno indurato, decorticato in Scotia (J. KEITH). — Ascoma $\frac{1}{4}$ -1 mm. lat., stipes $\frac{1}{2}$ mm. alt.

27. **Phialea Craginiana** (E. et E.) Sacc. *Peziza* Ell. et Ev. in Journ. 1064

Mycol. I, p. 47. — Ascomate stipitato, 2-3 mm. lat. levi, discoideo, madore albido, obscuriore in sicco; stipite filiformi, 2-3 mm. long.; ascis cylindraceis, 75 \approx 6, subsessilibus; sporidiis ovato-elipsoideis, e flavo hyalinis, binucleatis, 5-6 \approx 2,5-3, monostichis v. subdistichis.

Hab. in ligno valde putri, Topeka Amer. bor. — Affine *Ph. gracilipedi*.

28. **Phialea eburnea** (Desm.) Sacc. *Peziza* Desm. XIX, Not. p. 28, nec 1065

Cr. — Amphigena, sparsa aut vix gregaria, stipitata, minutissima eburnea; cupula hemisphærica, margine denticulato subpruinoso; stipite brevi; ascis clavatis, tenerrimis, 30 μ . long.

Hab. ad folia exsiccata graminum, in Gallia (ROBERGE). — Ascoma $\frac{1}{4}$ mm. lat., stipes $\frac{1}{6}$ mm. long.

29. **Phialea Volutella** (Fr.) Sacc. *P. Volutella* Fr. S. M. II, p. 127, 1066

Volutella nuda Tod. Meckl. fung. I, p. 29, t. 6, f. 44. — Fugax, cupula primum revoluta, demum disciformi punctato-umbilicata alba; stipite setaceo nigro.

Hab. in culmo triticeo putrido in agro Mecklenburgensi. — Diameter pilei linearis; structura fungilli tam tenera, ut vita solum ephemera gaudeat; cupula junior revoluta, dein depressa, centro nonnumquam umbonato. An a *P. culmigena* diversa?

30. **Phialea tenuissima** Quél. X Suppl. p. 13, t. IX, f. 9, Pat. t. 284. 1067

— Ascomatibus cyathiformibus, 200-400 μ . diam., ceraceis, hyalinis, margine subtili, hyalino-flocculoso; stipite capillari, glabro;

ascis cylindraceis; sporidiis bacillaribus, 5-7 μ . longis; paraphysibus filiformibus.

Hab. in culmis et foliis graminum Jura. — *Ph. caulincola* minor.

31. **Phialea petasata** (Karst.) Sacc., *H. petasatum* Karst. M. F. I, 1068 p. 131. — Sparsa, stipitata, dilute albida vel aquose pallida; cupula convexa, subitus concava, latit. 0,7-0,8 mm.; stipite subæquali, altit. circiter 0,7 mm.; ascis cylindraceo-clavatis, 48-50 \approx 4-5, obturaculo jodo vix v. obsoletissime cœrulecente; sporidiis vulgo oblique monostichis, fusoideo-elongatis, rectiusculis, eguttulatis, 10-15 \approx 2-3; paraphysibus parcis, filiformibus, apicem versus incrassatis.

Hab. ad culmos *Phragmitis communis* vetustos in par. Tammela Fenniae passim. — Externa facie a *Ph. culmigena* haud dissimilis, partibus autem internis diversa.

32. **Phialea pileata** (Karst.) Sacc., *Helotium pileatum* Karst. M. F. 1069 I, p. 130, *Pez. pileata* Karst. Monogr. Pez. pag. 150 (1869) nec Peck (1874). — Sparsa, stipitata, livido- vel cæsio-pallida; cupula convexa, margine deflexo sæpeque repando-lobato, latit. circiter 2 mm.; stipite sursum incrassatulo, altit. 2-3 mm.; ascis confertis cylindraceo-clavatis, 70-90 \approx 7-8, obturaculo jodo intense cœrulecente; sporidiis fusoideo-elongatis, rectis v. curvulis, simplicibus ac guttulatis, 16-32 \approx 2,5-4; paraphysibus gracilescensibus sat parcis.

Hab. ad culmos calamosque putridos nec non supra terram locis aquosis v. humidis et in margine rivulorum in Fennia et Britannia.

33. **Phialea confinis** (Karst.) Sacc. *H. confine* Karst. M. F. I, p. 139. 1070 — Dilute albida, stipitata; cupula scyphoidea, dein explanata, margine pruinoso-crenulato, 0,3-0,7 mm. lat.; stipite fuscescente, 0,4-0,7 mm. alt.; ascis cylindraceo-clavatis, 36-42 \approx 4,5-5,5, obturaculo jodo vix v. obsoletissime cœrulecente; sporidiis fusoideo-elongatis, distichis 10-15 \approx 2.

Hab. ad culmos *Junci conglomerati* emortuos, locis uidis dejectos, fine mensis Octobris in Mustiala Fenniae. — A *Ph. cyathoidea*, cui proxima, margine cupulæ crenulato et sporidiis majoribus nec non colore differt.

34. **Phialea culmigena** (Fr.) Sacc. *H. culmigenum* Fr. Summ. Veg. 1071 Scand. p. 355, Karst. M. F. I, p. 129, *Peziza culmigena* Fr. Syst. Myc. II, p. 127. — Subgregaria, dilute pallida vel albida, stipitata, pellucens; cupula plana, ambitu reflexa, subumbilicata, cir-

citer 0,4-0,6 mm. lat.; stipite æquali subcapillari, 0,5-1,5 mm. alt.; ascis cylindraceo-clavatis; sporidiis ellipsoideis vel oblongatis eguttulatis, 9-14 \approx 4-6.

Hab. ad culmos Tritici repentis emortuos in Fennia et Suecia. — *Var. scirpina* Fr. l. c. utrinque plana margine prominente stipite brevi firmiore. In calamo Scirpi lacustris in Suecia.

35. **Phialea culmicola** (Desm.) Gill. Disc. p. 103, *Hel. culmicolum* 1072 (Desm.) Karst. Symb. p. 237, M. F. I, p. 134, *Peziza culmicola* Desm. Espec. nouvell. 1836, p. 10, *Cyathicula culmicola* De Not. Disc. pag. 28. — Sparsa v. subgregaria, stipitata, firmula; cupula turbinata, pallescente vel albo-fuscescente sicca subfusca, pruinosa v. furfurella, margine connivente, longitudinaliter striatula, vix 0,5 mm. lat.; stipite æquali, brevi, crassiusculo, firmo, interdum subroseo; ascis cylindraceo-clavatis, 126 \approx 10, obturaculo jodo leviter cœrulescente; sporidiis fusoideo-elongatis, rectis v. subrectis, 9-12 \approx 1,5-2; paraphysibus gracilibus.

Hab. in culmis Avenæ, Secalis, Hordei, Tritici in Gallia, Fennia, Amer. bor.

36. **Phialea Constantini** (Boud.) Sacc. *Helotium Constantini* Boud. 1073 Bull. Soc. Myc. IV, p. 81, t. 17, f. 1. — Stipitata, 2-3 mm. lata et totidem alta, uda omnino hyalina, siccitate albida levis, intus gelatinosa, stipite subcrasso, etiam pellucido; hymenio primo concavo, dein expanso convexoque medio subumbilicato; paraphysibus simplicibus cylindracis, ad apicem non aut vix incrassatis, non septatis nisi ad basim, intus guttulatis, 3 μ . cr.; ascis cylindraceo-clavatis, octosporis, vacuis foramine submarginato, 170-190 \approx 7-8; sporidiis maxime arcuatis, apicibus subattenuatis, intus utrinque varie guttulatis, centro vacuis, 15-18 \approx 3.

Hab. ad folia putrida Caricum in paludosis, Montmorency Galliæ (COSTANTIN).

37. **Phialea pteridicola** (Cr.) Gill. Disc. p. 104, *Peziza pteridicola* 1074 Cr. Fl. Finist. pag. 50. — Ascomate levi, griseo, pedicellato, 1-2 mm., disco virente; ascis clavatis, octosporis; sporidiis teretiusculis, hyalinis; paraphysibus filiformibus granulosis, viridulis.

Hab. ad basim Pteridis aquilinæ, Finistère Galliæ.

II. Disco flavo, flavidо, aureo, aurantio.

38. **Phialea Hedwigii** (Phill.) Sacc. *Hymenoscypha* Phill. Disc. p. 130, 1075 *Octospora fungoidaster* Hedw. Frond. II, p. 53, t. XIX, fig. 1-4, *Pez. virgultorum* Fl. Dan. t. 1016, fig. 2. — Stipitata, concava v.

plana, flava, aurantio-tincta, margine integro, infra pallidiore; stipite longiusculo, supra medium tomentoso, albo, incrassato; ascis tereti-clavatis; sporidiis oblongis v. oblongo-ellipsoideis, 2-guttatis, 7-10 \times 3.

Hab. in ramis emortuis *Coryli* prope Shrewsbury Britanniæ.

— Ascoma 1-3 mm. lat. et alt., stipes matrici mycelio albo affixus.

39. **Phialea Tuba** (Bolt.) Gill. Disc. pag. 99, *Helotium* Fr. Summa 1076 p. 355, Hazsl. Ein. Disc. p. 15, t. III, f. 3, *Pez. Tuba* Bolt. fung. t. 106, f. 1, Pers. Myc. Eur. p. 278, Fr. Syst. Myc. II, pag. 128, *Hymenoscypha* Phill. Disc. pag. 126, *P. ochracea* Schum. Saell. p. 431. — Lutea; cupula turbinata, disco plano, margine tumido, stipite longo gracili; ascis (teste Hazsl. l. c.) cylindraceis, breve stipitatis, octosporis; sporidiis ellipsoideis, oblique monostichis, hyalinis.

Hab. ad ramos locis humidis prostratos in Britannia, Belgio, Suecia, Hungaria. — Sec. ic. Hazsl. loc. cit., stipes 10 mm. long, basi nodulosus. Stipes subfiliformis, æqualis, incurvus v. flexuosus, e loco longitudine varius. Cupula extus levis, primo clavata.

40. **Phialea salicina** (Pers.) Sacc. *H. salicinum* (Pers.) Fuck. Symb. 1077 p. 314, *Peziza salicina* Pers. Syn. p. 663, *Helotium virgullorum* $\beta.$ *salicinum* Fr. S. v. Sc. p. 355, Syst. Myc. II, p. 119, sub *Peziza fructigena* $\beta.$ *salicina*, *Pez. flavescens* Holms. Ot. t. XI. — Ascomate flavo, tenui, flexuoso, difformi, stipite tenui deorsum attenuato; ascis stipitatis, clavatis, 8-sporis, 120 \times 14; sporidiis in asci apice subdistichis, oblongis, parum curvatis, utrinque acuminatis subtilissime apiculatis, hyalinis, 24 \times 6.

Hab. in ramulis tenellis, valde humidis *Salicis triandrae* ad Rhenum pr. Æstrich, in Italia et in Suecia.

41. **Phialea Aspegrenii** (Fr.) Gill. Disc. p. 107 c. ic. *Pez. Aspegrenii* 1078 Fr. Syst. Myc. II, p. 131, *P. bicolor* Sowerb. t. 369, f. 7, Fries Obs. 2, p. 305, nec Bull. *P. Calicum* Pers. Myc. Eur. p. 280 nec Decand. — Cupula subrepanda, glabra, disco flavo extus stipiteque subadscendente albis; ascis cylindraceis, octosporis; sporidiis oblongis v. oblongo-ellipsoideis, 2-guttatis, 8-10 \times 3,5 (Phill.).

Hab. ad ligna mucida in Britannia, Gallia, Suecia. — Gregaria, 5-7 mm. alta, stipite subtenui.

42. **Phialea crocea** (Schw.) Sacc. *P. crocea* Schw. Syn. p. 116. — 1079 Subcorneo-indurata, crocea, fasciculata, 2-3 cm. alta; stipite crasso sulcato incurvo; cupula varie contorta et lobata, 4 mm. lat., margine obtuso subflexuoso; sporidiis obovatis, minutis, 5 μ . longis.

Hab. ad ramos Abietis Fraseri etc. in Carolina sup. et New York. — An *Ciboria?*, an *Guepinia?*

43. **Phialea quercicola** (Saut.) Sacc. *Peziza quercicola* Saut. Pilz. 1080 Salzb. p. 15, Wint. Hedw. 1881, pag. 133. — Ascomate tenui, albido, cyathiformi, cernuo, disco aurantiaco, stipite brevi, filiformi; ascis elongato-clavatis, octosporis, 45–53 × 5; sporidiis subdistichis, cylindraceo-fusoideis, utrinque rotundatis, continuis, hyalinis, 7–9 × 1,5–2; paraphysibus cylindraceis crassis, apice cuspidatis, ascos superantibus.

Hab. in cortice Quercino putrescente, Salzburg. — Ob paraphyses ad *Dasyscypham* nutat.

44. **Phialea verpoides** (Saut.) Sacc. *Peziza verpoides* Saut. in Hedw. 1081 1876, p. 149. — Ascomate disciformi, rotundo, flavidio, 4 mm. lato, margine integro, revoluto, disco sicco rugoso, subtus levi, pallido, stipite 2–3 cm. longo, dilatato, flexuose contorto, nudo, fusco, 2 mm. crasso.

Hab. ad ligna putrida pr. Salzburg, raro.

45. **Phialea crocata** (Mont.) Sacc. *Pez. crocata* Mont. Cent. II, n. 19, 1082 Syll. Crypt. n. 625. — Minuta, tenuis, stipitata; cupula demum applanata amœne crocea utrinque glabra; stipite albido-succineo albo-villosulo, demum glabrescente pellucido.

Hab. ad petiol. folior. deject. in silvis sinnamariensibus (LÉPRIEUR).

46. **Phialea micans** (Saut.) Sacc. *Pez. micans* Sauter in Flora 1841, 1083 p. 380. — Epiphylla, sparsa, punctiformis, sulphurea, cupulis ex urceolato deplanatis cum stipite brevi conicis.

Hab. in foliis Arabidis pumilæ in agro urbis Stiriæ pr. Mühlbach, altit. circ. 1000 m.

47. **Phialea Hymenula** (Fuck.) Sacc. *Peziza* Fuck. Symb. p. 308. — 1084 Cupulis breviter stipitatis, 1-1½ mm. lat., primo concavis, demum explanatis, extus pallidioribus, disco flavo, marginato; ascis longissime stipitatis, clavatis, 8-sporis, 76 × 4; sporidiis subdistichis, cylindraceis, parum curvatis, 3–4-guttulatis, hyalinis, 8–10 × 2, conidiis, ut in fungo conidiophoro, commixtis.

Hab. in caulibus putridis Sambuci Ebuli et Georginæ in Rhenogovia. — Sec. Fuck. l. c. status conidici hujus *Phialeæ* sunt *Hymenala vulgaris* Fr., *H. Georginæ* Wallr. et *H. Ebuli* Corda.

48. **Phialea vitellina** (Rehm) Sacc. *Helotium vitellinum* Rehm Ascom. 1085 n. 513. — Distincte et graciliter stipitata, primitus cyathoidea, cupula urceolata, pallide fusca, dein explanata, vitellina, disco auran-

tiaco, rubro, prosenchymatico-contexta, gregaria, c. 2 mm. alta, 1,5 mm. diam.; ascis clavatis, 8-sporis, 75-80 \approx 9; sporidiis elongato-cuneiformibus, uno apice obtusis, altero acutatis, rectis v. subcurvatis, nucleis 1-2 magnis præditis, hyalinis, distichis, 18 \approx 4; paraphysibus filiformibus, clavulatis, hyalinis.

Hab. ad caules putridos Spirææ Ulmariae prope Augustam Vindelicorum seu Augsburg (BRITZELMAYR).

49. **Phialea Phragmitis** (Saut.) Sacc. *Peziza Phragmitidis* Sauter 1086
in Flora 1845, pag. 133. — Tenuis, albida; cupula hemisphærica, disco flavido subexcavato, margine aquose pallido, stipite basi bulboso filiformi elongato.

Hab. in culmis putrescentibus Phragmitis in agro urbis Stiriae. — *Stamn. Personii* proxima dicitur. Stipes cupula circ. 1 mm. lata, 4-6-plo longior.

50. **Phialea piriformis** (Hedw.) Gill. Disc. pag. 107, *P. piriformis* 1087
(Hedw.) Fr. Syst. Myc. II, p. 121, *Octospora* Hedw. Musc. I, p. 32,
t. 10, f. 4. — Cupula globoso-turbanata, sordide flavescente, mar-
gine sub prominulo, albidiore, stipite filiformi albido.

Hab. ex alis Muscorum Mnii, Bryi locis udis erumpens in Suecia et Germania. — Distinctissima, solitaria, subaquosa, stipite adscendente, 6 mm. circiter longo, æquali. Variat colore magis fusco margineque crenulato.

51. **Phialea bryophila** (Fr.) Sacc. *Pez. bryophila* Fr. Syst. Myc. II, 1088
pag. 119. — Flavescens, tenuis; cupula hypocateriformi, stipite elongato tenui æquali subflexuoso.

Hab. ad muscos locis humidis in Suecia, Gallia et Belgio. — Sparsa, radicibus muscorum insidens. Stipes 12 mm. circiter lon-
gus, deorsum subfibrillosus, flacidus; cupula hemisphærica, in sti-
pitem vix attenuata, integerrima, 2-5 mm. lata. Habitus fere *Omphaliæ Fibulæ*. Huc Fl. Dan. t. 1016, f. 1 pertineret.

52. **Phialea epigæa** (Karst.) Sacc. *Helotium epigæum* Karst. Rev. mon. 1089
p. 126. — Sparsa, ceracea, stipitata, glabra, flava; cupula plano-
convexa, subtus concava, sat tenui, circiter 0,8 mm. lat., stipite
diametrum cupulæ æquante; ascis cylindraceo-clavatis, apice jodo
haud vel vix cœrulescentibus, 8-10; sporidiis in parte asci supe-
riore, distichis v. subdistichis, fusoideo-oblongatis, rectis, simplici-
bus, subeguttulatis, hyalinis, 13-14 \approx 4-5; paraphysibus filiformi-
bus, sat numerosis, apicem versus incrassatis, gracilescentibus.

Hab. prope Mustiala supra terram nudam. — Ad *Ciboriam ma-*
cilentam, quæ ad *Phialeam* facile trahitur, proxime accedit.

III. Disco rubente, roseo, incarnato.

53. **Phialea Broomei** (Phill.) Sacc. *Hymenoscypha* Phill. Disc. p. 129, 1090
t. V, f. 27, *Pez. araneosa* Herb. Kew. — Sparsa, stipitata, cyathiformis vel plana, carneo-rubra, glabra, firmula, margine integro, stipite gracili, cylindrico, flexuoso; ascis tereti-clavatis; sporidiis oblongis, utrinque rotundatis v. subfusoideis, 15 \approx 5.
Hab. in ligno putri in Britannia. — Ascoma 2 mm. lat. et alt.
54. **Phialea leucopsis** (B. et C.) Sacc. *Pez. leucopsis* B. et C. Cub. 1091
Fungi n. 691. — Carneo-pallida, irregularis, margine primum erecto demum reflexo; ascis linearibus; sporidiis allantoideis, 5 μ . long.
Hab. in ligno putri in insula Cuba. — Ascoma 3-6 mm. lat.
55. **Phialea rubescens** (Saut.) Sacc. *Peziza rubescens* Saut. Pilz. 1092
Salzb. pag. 14, Wint. Hedw. 1881, pag. 132, nec *Hel. rubescens* Crouan. — Ascomatibus gregariis, ex urceolato patellatis, extus rubescensibus, intus coccineis, margine integro, stipite brevissimo, firmo; ascis elongato-clavatis, basi attenuatis, octosporis, 100-110 \approx 6-7; sporidiis (immaturis), oblongis, hyalinis, 8 \approx 3; paraphysibus filiformi-clavulatis.
Hab. ad ligna putrida prope Salzburg. — Ascoma 1,5 mm. alt., 2,5 mm. latum, extus hyphulis patulis levibus vescitum. « An *Helotium?* ».
56. **Phialea littorea** (Fr.) Sacc. *Pez. littorea* Fr. Syst. Myc. II, p. 121, 1093
Cyathicula De Not. Comm. I, pag. 389, *Helotium* Crouan Finist. pag. 48. — Stipitata; cupula hemisphærico-depressa, disco rubescente, margine sublacerio; ascis....; sporidiis exiguis oblongo-teretiusculis, hyalinis.
Hab. ad littora lacuum supra Arundinis culmos siccios in Suecia, Germania, Belgio, Gallia. — Sec. Crouan Finist. pag. 49: asci subclavati, 8-spori; sporidia suboblonga; paraphyses filiformes, granulosæ. Parca, glabra, gregaria. Stipes brevis, sursum in cupulam dilatatus, unde tota fere obconica; margo inflexus, primo velatus. Affinis *P. alabastrinæ*, sed paullo major, nec ita pellucida.
57. **Phialea pallide-rosea** (Saut.) Sacc. *Pez. pallide-rosea* Saut. Pilz. 1094
Salzb. p. 14. — Breviter stipitata, gregaria; ascomate hemisphærico, roseolo, margine inflexo integro.
Hab. in culmis graminum in paludosis prope Salzburg. — Affinis *P. inflexæ* sed colore margineque diversa.
58. **Phialea rhodoleuca** (Fr.) Sacc. *H. rhodoleucum* Fr. Summa p. 355, 1095
Karst. M. F. I, p. 128, *Peziza* Fr. Syst. Myc. II, p. 127. — Te-

nuis, roseo-albida; capitulo utrinque plano; stipite æquali; ascis tereti-clavatis, 4-8-sporis; sporidiis oblongo-ellipsoideis, 10-14 \approx 4-5; paraphysibus filiformibus.

Hab. in stipitibus *Equiseti arvensis* aliisque quisquiliis locis udis dejectis in Suecia, Fennia, Britannia. — Substantia aquose pellucida; stipes glaber, 1 mm. longus, subinde vero ad 6 mm. elongatus; cupula orbicularis, integerrima, 2-4 mm. lata.

59. **Phialea subcarnea** (C. et P.) Sacc. *Peziza subcarnea* C. et P. 1096
Peck 27 Rep. p. 107 (*Peziza*) et Disc. of Un. St. I, p. 295. — Ascomatibus sparsis, stipitatis, tenuibus, initio clavatis, dein infundibuliformibus, omnino carneis; stipite longo, basi attenuato, sursum dilatato, margine contracto, pallido; ascis cylindraceis; sporidiis teretiusculis, obtusis, hyalinis.

Hab. ad Hepaticas in truncis vetustis in silvis Indian lake Am. bor. — Maculas suborbiculares efformat matricemque necat.

IV. Disco ochraceo, rufo, alutaceo.

60. **Phialea fructigena** (Bull.) Gill. Disc. p. 99 c. ic., *Peziza* Bull. 1097 Ch. p. 236, t. 228, Sowerb. t. 117, Fr. S. M. II, p. 118, Nees Syst. p. 267, f. 282, *Helotium* Fuck. Pat. t. 383, *Hel. flavescens* Pers. M. E. I, p. 346. — Tenacella, glabra, pallida, lutescens; cupula patellæformi; stipite longo tenui flexuoso; ascis elongato-clavatis, 8-sporis, 120 \approx 8; sporidiis oblongo-clavatis, antice obtusis, basi attenuatis, subinæquilateralibus, hyalinis, continuis, 14-17 \approx 4-5, 2-guttatis; paraphysibus filiformi-clavulatis.

Hab. in cupulis, glandibus *Quercus*, *Fagi*, capsulis *Castaneæ*, strobilis *Coniferarum* in Suecia, Gallia, Britannia, Fennia, Germania, Italia, Amer. bor. — Gregaria, quam maxime varia. Color albidus, flavus, rufescens; stipes e loco plus minus elongatus, non semper basi floccosa adhærens, sed passim nigricante subattenuata. Cupula 2-5 mm. lata, demum et convexa. (*P. convexa* Holmsk. Ot. 2, p. 34, t. 16).

61. **Phialea gracilis** (C. et P.) Sacc. *Helotium gracile* C. et P. in 1098 Peck 26 Rep. St. Mus. pag. 83. — Ochracea; cupula plana, dein convexa, immarginata, tenuiuscula, extus leniter pallidiori; stipite gracili, æquali, basim versus brunneolo, circiter diametrum cupulæ longitudine æquante; ascis cylindraceis; sporidiis cylindraceis vel subfusiformibus apice obtusis, 2-3-nucleatis, 17-20 μ . longis.

Hab. in caulis herbarum Center Americæ borealis. — Habitu

et magnitudine *Ph. cyathoideam* in mentem revocat at cupulæ nunquam clausæ.

62. **Phialea Scutula** (Pers.) Gill. Disc. pag. 108, *Pez. Scutula* Pers. 1099
M. E. I, p. 284, *Fr. Syst. Myc. II*, p. 123, *Helotium* Karst. *M. F. I*, p. 110, *Pat. Tab. f. 94*, *Hymenoscypha* Phill. Disc. p. 137, *P. virgultorum* var. *Scutula* pl. auctorum. — Firmula, hypocrateriformis, demum subplana, levis, luteo-testacea, margine integer-rimo, stipite tenui æquali; ascis tereti-clavatis, 80-90 \times 8-10; sporidiis fusoideis v. fusoideo-clavulatis, 2-3-guttulatis, subinde pseudo-septatis, 16-20 \times 3-5.

Hab. ad caules plantarum exsiccatos e. g. *Artemisiæ*, *Ballostæ*, *Teucrīi*, *Gei*, *Rumicis*, *Cirsii* in Suecia, Britannia, Germania, Italia, Gallia, Belgio, Amer. bor. — Stipes pro ratione tenuis, cupulæ diametro brevior, paullo dilutior; cupula magis magisque dilatata, leviter concava, 1 mm. lata. Colore et cupula firma a sequ. differt. — Var. **petiolicola** Sacc.: sporidiis uno apice minus acuminatis. In petiolis *Robiniæ Pseudacaciæ* in Ital. bor. — Var. **fucata** Phill. Disc. p. 137: ascomate subhemisphærico, margine incurvo, disco brunneo-flavido, extus albido, flavo-variegato. In caulibus *Polygoni aqua immersis* Ellesmere Brittaniæ. — Var. **Menthæ** Phill. Disc. p. 137: ascomate plano v. convexo, disco læte flavo, stipite tenui; sporidiis 14-20 \times 3-5, bi-triguttatis simulateque 1-septatis. In caulibus emortuis Menthæ, Shrewsbury Brittaniæ. — Var. **Rudbeckiæ** Phill. Disc. p. 138: ascomate 2 mm. lato, stipite 2-3 mm. alto, tereti, in cupulam expanso; sporidiis 18-25 \times 4-6. In caulibus emortuis Rudbeckiæ, Batheaston Brittaniæ (BROOME).

63. **Phialea virgultorum** (Vahl) Sacc. *Peziza* Vahl Fl. Dan. t. 1016, 1100 fig. 2, *Helotium virgultorum* (Vahl) Karst. *M. F. I*, p. 109, *Hymenoscypha* Phill. Disc. p. 134. — Gregaria, stipitata, rigidula, glabra, patellaris v. subinde convexa, extus pallida, disco flavidorufo; stipite tenui subflocculoso; ascis cylindraceo-clavatis; sporidiis fusoideis, sæpe curvulis, 2-guttatis, 14-23 \times 3-5, demum simulate 1-2-septatis, hyalinis; paraphysibus filiformibus.

Hab. in ramis emortuis arborum in Germania, Dania, Fennia, Gallia. — Ab affini *Ph. fructigena* differt colore disci et sporidiis.

64. **Phialea alutipes** (Phill.) Sacc. *Peziza alutipes* Phill. Grevillea VII, p. 23. — Gregaria stipitata, infundibuliformis, dein plana alutacea, margine integra; stipite firmo primo, gracili, longo et flexuoso, rarius brevi, in cupulam rugosam dilatato, concolore; ascis

crasse clavatis; sporidiis oblongo-ovoideis, multinucleatis, 18-20 \approx 5-8; paraphysibus filiformibus, clavulatis, subcoloratis.

Hab. in foliis dejectis Libocedri decurrentis in California.

65. **Phialea ceratina** (Berk.) Sacc. *Peziza ceratina* Berk. Fl. Tasm. 1102 p. 275. — Turbinata, stipitata, glabra, pallide fulvo-cornea, hymenio plano marginato; sporidiis 20-21 μ . long. oblongo-clavatis.

Hab. in foliis Eucalypti in Tasmania (ARCHER). — Asci magnifici. *Phialeæ cyathoideaæ* et *titubanti* subaffinis. Minuta 1 mm. alta.

66. **Phialea longipes** (C. et P.) Sacc. *Peziza longipes* C. et Peck Disc. 1103 Unit. St. I, p. 295. — Flavida; ascomate longe stipitato, concavo, 5-7 mm. lat., leniter concavo, demum subplano, disco rufo-flavo; stipite æquali, gracili, prælongo; 2-3 cm.; ascis cylindraceis; sporidiis anguste ellipsoideis, rectis v. curvulis, uninucleatis, 5-6 \approx 2-2,5.

Hab. in petiolis foliorum New York, Memphis in America boreali (PECK).

67. **Phialea byssogena** (Berk.) Sacc. *Peziza byssogena* Berk. Fl. 1104 Tasm. p. 275. — Ochracea; cupula concava, stipite elongato, cylindrico subtiliter pruinosa e floccis intertextis radiantibus oriundo; ascis clavatis; sporidiis elliptico-oblongis, 8-9 μ . longis.

Hab. in lignis dejectis in Tasmania (ARCHER). — *Ph. lutescenti* affinis. Ascoma 1-1,3 mm. lat., stipes 2-2,5 mm. long.

68. **Phialea aquatica** (Curr.) Sacc. *Helotium aquaticum* Curr. Linn. 1105 Trans. XXIV, p. 154, tab. XXV, fig. 19. — Solitaria, stipitata, ascomate minuto, concavo vel convexo, stipite paullo latiore; stipite filiformi tenui; ascis cylindraceis vel cylindraceo-clavatis; sporidiis ellipsoideis vel clavulatis, biguttulatis, 13-15 \approx 5-6, hyalinis.

Hab. in fragmentis ligneis aqua immersis Kent Britanniæ. — Ascoma 1-2 mm. lat.; stipes 5-7 mm. long.; color fungi non indicatus.

69. **Phialea Calyculus** (Sow.) Gill. Disc. pag. 108, *P. Calyculus* Sowerb. fung. t. 116, Pers. Syn. p. 660, Fr. Syst. Myc. II, p. 129, *Pez. Infundibulum* Batsch. f. 147, *P. Calycula* Schum. — Lutescenti-brunnea, stipitata, glabra; cupula concava elevato-marginalata; stipite brevi crasso, deorsum attenuato; ascis cylindraceo-clavatis; sporidiis oblongis vel oblongo-ellipsoideis vel subclavatis, rectis curvulis 15-18 \approx 3-5; paraphysibus filiformibus.

Hab. ad ligna et sub cortices in Britannia, Suecia, Germania, Gallia, Belgio. — Unicolor, subinde rufo-ferruginea; cupula 2-3 mm. lata, margine in siccis involuto.

70. **Phialea exarata** (Berk.) Sacc. *Peziza exarata* Berk. North Am. 1107
Fungi n. 750. — Mellea; cupula irregulari concava, margine inflexo; stipite crasso, eximie costato pruinoso; sporidiis minutis oblongis.

Hab. in ligno putri in Carolina infer. Amer. bor.

71. **Phialea emergens** (Cooke et Phill.) Sacc. *Hymenoscypha* C. et 1108
Phill. Disc. p. 139. — Sparsa vel fasciculata, e cortice emergens hemisphærica vel infundibuliformis, stipitata, glabra, dilute ochracea, disco obscuriore, margine integro; stipite flexuoso, deorsum attenuato; ascis cylindraceis vel tereti-clavatis; sporidiis fusiformibus, continuis, rectis curvulisve, 12-14 \approx 1-1,2; paraphysibus non visis.

Hab. ad ramos emortuos terra subinfossos in Scotia. — Asco-
ma 0,5-1,5 mm. lat., stipes 1-1,5 mm. longus.

V. Disco fusco, brunneo, fumoso.

72. **Phialea glanduliformis** (Rehm) Sacc. *Helotium glanduliforme* 1109
Rehm Ascom. Lojk. n. 14 et Ascom. n. 255, Helotium cyathoideum var. *multicolor* Karsten Myc. Fenn. II, pag. 136. — Ascomatibus sparsis, crasse stipitatis, stipite fuscidulo, basi subhyalino, turbinatis, c. 1-1,5 mm. altis, c. 5-7 mm. diam., siccis jodo apice cœrulecente, margine urceolato-involuto fuscis, humidis concavo-urceolatis, flavidulis, prosenchymatico-flavido-contextis, glabris, siccis extus subruguloso-striatis; ascis clavatis, 8-sporis, 45 \approx 6; sporidiis ellipticis subobtusis clavulatis, rectis, hyalinis, distichis, 9 \approx 2,5; paraphysibus filiformi-clavulatis ascos vix superantibus.

Hab. ad ramulos sicclos Ononidis spinosæ prope Windsheim Franconiæ bav. et prope Pesth. — Var. **Bartsiae** Rehm Asc. n. 559: ascis 36-42 \approx 6; sporidiis 8-10 \approx 2-2,5. In caule Bartsiae in Germania.

73. **Phialea perelegans** (Saut.) Sacc. *Peziza perelegans* Sauter in Flora 1845, p. 133. — Tenuis, albida, urceolato-hemisphærica, margine inflexo integro, disco nigrescente, stipite tenui longiusculo.

Hab. in culmis graminum vetustis in agro urbis Stiriae. — *P. litoreæ* et *cyathoideæ* affinis. Stipes 2,5 mm. longus; cupula minuta, globosa.

74. **Phialea Cacaliæ** (Pers.) Gill. Disc. p. 104, *Pez. Cacaliæ* Pers. 1111
M. E. I, p. 285 (*Pez.*) Fr. S. M. II, p. 122, *H. Urticæ* forma Rehm Asc. n. 59. — Turbinata, levis, fuscescens, ore subconnivente albidiore, stipite sublongo crassiusculo; ascis linearibus, 8-sporis, 50-

68 \approx 6; sporidiis elongato-distichis, angustissime cylindraceis, sub-curvatis, 8 \approx 6-2.

Hab. ad caules plantarum, Compositarum, e. g. *Cardui*, *Senecionis*, *Cacaliæ*, *Carlinæ*, *Lappæ*, *Daturæ*, in *Helvetia*, *Germania*, *Belgio*, *Britannia*. — Glaberrima, subnitens, firmula, junior albida adulta et sicca fusca, humida infundibuliformis, immo explanata, 0,5 mm. lat.; stipes quam *P. cyathoideæ* brevior et firmior. — ***pallidior** Sacc. Mich. II, p. 328: ascomatibus typo minoribus atque pallidioribus; ascis 50-60 \approx 5; sporidiis tereti-fusoides, 7-9 \approx 1-1 $\frac{1}{4}$, hyalinis. In caule putri *Amaranthi* in Gallia.

75. **Phialea sordida** (Fuck.) Sacc. *Helotium sordidum* (Fuck.) Rehm 1112
Ascom. n. 414 nec Phill. (= *Helotium*), *Pezizella sordida* Fuck. Symb.
pag. 414. — Gregaria, sordida; cupulis glabris, 1-2 mm. latis, pa-
telliformibus, marginatis, margine demum criso-lacerato, involu-
to, stipite firmo, cupulæ diametrum vix æquante, disco demum ru-
bello; ascis linearibus, stipitatis, 8-sporis, 30-40 \approx 6-7; sporidiis
distichis, cylindraceis, curvatis, hyalinis, 8-9 \approx 2.

Hab. in ramis putridis plerumque decorticatis *Rosæ caninæ*
in Germania.

76. **Phialea fumosella** (C. et E.) Sacc. *Peziza fumosella* C. et E. 1113
Grevillea VI, p. 91. — Stipitata, minuta, fumosa; cupulis clavatis,
demum apertis, cyathiformibus, margine discoque pallido; stipite
obscurore sursum in cupulam expanso; ascis cylindraceo-clavatis;
sporidiis linearibus, rectis vel curvulis, 10 μ . longis; paraphysibus
filiformibus.

Hab. in foliis Pinorum in New Jersey Amer. bor. — Asco-
ma $\frac{1}{4}$ - $\frac{1}{3}$ mm. lat., $\frac{1}{2}$ mm. alt.

77. **Phialea uveata** (Schum.) Sacc. *Pez. uveata* Schum. Saell. p. 430, 1114
Fries Syst. Myc. II, pag. 126, Fl. Dan. t. 2034, fig. 1. — Pallide
fusco-umbrina; cupula subhemisphærica concava, margine tumido,
stipite brevissimo crassiusculo.

Hab. ad stipulas, radices exsiccatas graminum raro in Sue-
cia. — Affinis *Cib. Cauco*; an ex hoc genere?. Gregaria, subcæspi-
tosa, ceracea, 1 mm. lata, glabra, levis; stipite sursum dilatato.

78. **Phialea nitidula** (B. et Br.) Sacc. *Peziza nitidula* B. et Br. Ann. 1115
Nat. Hist. n. 570, *Hymenoscypha* Phill. Disc. p. 142. — Asco-
mate minuto, subhemisphærico, irregulari, extus aquose brunneo,
minute furfuraceo, nitidulo, sat firmo, stipite brevi æquali; ascis
cylindraceo-clavatis; sporidiis fusoideis, 7-10 \approx 3; paraphysibus fi-
liformibus.

Hab. in foliis emortuis Airæ cæspitosæ prope Bath in Britannia. — *Phialea Cacalice* affinis.

79. **Phialea palustris** (Saut.) Sacc. *Pez. palustris* Saut. Salzb. p. 9. 1116
— Albido-nigrescens; stipite cylindrico; ascomate hemisphærico, urceolato, margine inflexo integro.

Hab. in radicibus Equiseti palustris in Pinzgau. — $\frac{1}{2}$ mm. crassa, gregaria, levis; stipite brevi, in sicco nigrescente. Affinis *P. littoreæ*.

B. Hymenoscypha Sacc. Ascoma pruinatum v. puberulum.

I. Disco albo, albido, cinereo, dein saepe diverse colorato.

80. **Phialea tigillaris** (Karst.) Sacc. *Hel. tigillare* Karst. M. F. I, 1117 p. 145. — Subsparsa, substipitata, obconica, villosulo- vel puberulo-furfurella, albida; cupula sicca subsphæroideo-contracta, pallida, albo-puberulo-furfurella, 0,3-0,4 mm. lat.; stipite crassulo, brevissimo, in cupulam abeunte vel fere nullo; ascis clavatis, 63-73 \approx 8-10, obturaculo minutissimo jodo vix vel obsolete cœrulecente; sporidiis plerumque distichis, oblongatis, curvulis vel rectis, 2-raro 4-guttulatis vel spurie uniseptatis, 10-16 \approx 3-4-5; paraphysibus haud bene discretis, 1 μ . cr.

Hab. ad lignum fabrefactum pineum putridum in par. Tam-mela Fenniæ.

81. **Phialea sauciella** (Karst.) Sacc. *Hel. sauciellum* Karst. M. F. I, 1118 p. 143. — Gregaria, primum obconica, pallida vel lutescens, albo- vel albido-furfurella-villosula, deinde distincte stipitata, pallida vel lutescens vel sanguinea vel rubella vel fuscescens vel atrata, glabrescens; cupula e sphæroideo-hemisphærica, margine obtuso, repando-lacero, 0,3-0,4 mm. lat.; ascis cylindraceo-clavatis, 30-60 \approx 5-7; sporidiis fusoideo- vel ovoideo-oblongatis aut ovoides aut subellipsoideis, 4-10 \approx 2-3.

Hab. ad lignum ramorum Alni incanæ putrescens, prope Mustiala Fenniæ. — Facillime dignoscitur colore sanguineo vel sanguineo-purpureo, quo matricem inficit.

82. **Phialea pertenuis** Sacc. *Peziza pertenuis* Sacc. Fung. Ven. Ser. IV, n. 84, *Helotium* Sacc. F. ital. fig. 1334. — Cupulis ceraceo-mollibus, microscopicis, cum stipite brevi crassiusculo infundibuliformibus, niveis, furfurellis, totis 120-130 μ . altis, stipite 80 \approx 20, disco concaviusculo concolori, margine minutissime crenulato; ascis tenellis clavulatis, 30 \approx 4, subsessilibus; paraphysibus filifor-

mibus clavulatis, obvallatis, 8-sporis; sporidiis oblique monostichis v. subdistichis, ovoideis, $3\frac{1}{2} \times 2$, hyalinis.

Hab. in ramis decorticatis fag in eis in silva Cansiglio, Ital. bor.

— Affinis videtur *Trichopezizæ tenuissimæ* (KARST.).

83. **Phialea gibba** (A. S.) Sacc. *Hel. gibbum* A. S. Nisk. p. 350, t. 1120 IV, f. 1, *Pez. tubæformis* Wall. Crypt. n. 2583. — Ascomatibus umbonato-gibbis, margine inflexo cupulatis, sparsis, tenuissime membranaceis pallidis, extus fibrillis tenerrimis delicatis floccoso-furfuraceis gryseo-cinerescentibus undique æqualiterque superfusis, siccando margine inflexo fissis angulatisve; stipite gracili delicatulo.

Hab. ad strobilos Pini in Lusatia et ad ramulos Rosæ humi jacentes agri heringens. Thuring. (WALLR.). — An potius *Cyphella?* Stipes 6-7 mm. long., ascoma vix 1 mm. lat.

84. **Phialea tenera** (Saut.) Sacc. *Peziza tenera* Saut. Pilz. Salzb. 1121 p. 9. — Albida, cupulata, tenuis, concava; margine niveo, flocculoso, stipite brevi crassiusculo.

Hab. in foliis putrescentibus Berberidis prope Mittersill. — Affinis *P. chionæ*; $\frac{1}{4}$ mm. crassa.

85. **Phialea sordidata** (K. et Starb.) Sacc. *Helotium sordidatum* Karst. 1122 et Starb. Hedw. 87, p. 124 et Symb. XXIV, pag. 15. — Subgregaria, sessilis vel breviter stipitata, firmula, glabra; cupula primus sphæroidea, dein explanata, subpuberula, fuscescente-pallida, disco pallido, circiter 0,3 mm. lata; stipite æquali brevi crassiusculo, firmulo, saepe nullo; ascis clavatis, 60×9 ; sporidiis distichis, fusoideo-oblongatis, 1-2-guttulatis, $15 \times 3-4$.

Hab. in foliis putrescentibus Spirææ Ulmariae ad Mustiala Fenniæ (STARBACK).

86. **Phialea Eschscholtziæ** (P. et H.) Sacc. *Peziza Eschscholtziae* 1123 Phill. et Harkn. Calif. p. 2. — Sparsa, breviter stipitata, cyathiformis, cinerea, pruinosa; margine subinciso, disco concolori; ascis cylindraceis; sporidiis fusoideis, biguttulatis, 10×2 ; paraphysibus non visis.

Hab. in caulis emortuis Eschscholtziæ californicæ in California. — Ascoma 0,5-0,7 mm. latum.

87. **Phialea gemmarum** (Boud.) Sacc. *Helotium gemmarum* Boud. 1124 Bull. Soc. Myc. IV, pag. 81, tab. 17, fig. 2. — Minuta, alba, stipitata 0,3-0,7 mm. lata, 1-1,5 mm. alta, tenuis, crateriformis, marginata, nivea, extus pube tenui vix conspicua conspersa; hymenio levigato concolore et stipite gracili etiam puberulo ad basim lutescente; paraphysibus parcis, simplicibus aut divisis, non septatis,

intus subgranulosis, ad apicem non incrassatis, 2-3 μ . crassis; ascis cylindrico-clavatis, octosporis, 50 \approx 6-7, vacuis submarginatis; sporidiis minutis, oblongo- aut clavato-fusoideis, intus vix v. tenuiter granulosis, 6-8 \approx 3-4.

Hab. ad gemmas putridas Populi nigræ prope Paris Galliae.

88. **Phialea lateritio-alba** (Karst.) Sacc. *Helotium lateritio-album* ¹¹²⁵ Karst. Rev. Myc. 1887, p. 159 et Symb. XXIII, p. 3. — Sparsa, breviter stipitata, cyathoidea, alba vel lateritia, tomento pertenui niveo obtecta, 0,2-03 mm. alta; ascis cylindraceo-clavatis, 33-40 \approx 6; sporidiis distichis, fusoideo- vel clavulato-bacillaribus, rectis, vulgo guttulatis, 6-10 \approx 1,5.

Hab. in foliis siccis Scirpi silvatici in prato mustialensi, Myllypera Fenniæ.

89. **Phialea Morieri** Gill. Disc. p. 104. — Cyathiformis, madore aper- ¹¹²⁶ta, siccitate clausa, rufo-pallida, glabra, excepto margine albo-pulveraceo, 2-3 mm. lat.; disco albido; stipite brevi, deorsum le- niter incrassato, albido vel pallide rufo.

Hab. ad gramina et herbas varias in Gallia. — Ascoma sic- cando fuscescit.

90. **Phialea versicolor** Quél. V Suppl. p. 329, tab. VI, f. 10, *Caly- ¹¹²⁷cella mira* Quél. Enchir. p. 305. — Ascomatibus cyathiformibus, 500-800 μ . diam., breviter stipitatis, pruinosis, margine involuto, sub lente puberulis niveo-albis sed mox flavo-citrinis v. roseis; sporidiis ellipsoideis, fusiformibus, 6 μ . longis, binucleatis.

Hab. in Filicibus e. g. Scolopendrio putrescentibus, Jura. — Stipes teretiusculus ascomate paullo brevior.

II. Disco flavo, flavidо, lutescente.

91. **Phialea affinissima** (Peck) Sacc. *Helotium affinissimum* Peck 33 ¹¹²⁸ Rep. p. 32. — Ascomatibus subcæspitosis, stipitatis, planis vel con- vexitatis, 2-4 mm. lat., flavis, margine extusque, pruinulosis, stipite subcylindrico, flavidо 4-9 mm. long.; ascis cylindraceis; sporidiis oblongis obtusis, 8-10 μ . longis.

Hab. in ramis putridis terra infossis, Albany Amer. bor. — A *Ph. lutescente* differt habitu cæspitoso, sporidiis minoribus etc.

92. **Phialea fugitiva** Quél. X Suppl. p. 13, tab. IX, f. 7. — Ascoma- ¹¹²⁹ tibus ceraceis, urceolatis, 500-800 μ . diam., breviter stipitatis, te- nueriter puberulis, margine crenulato, isabellinis diaphanis; hymenio concavo, primo flavescente dein croceo; sporidiis ellipsoideo-lanceo- latis, 11 μ , longis, binucleatis.

Hab. in cortice emortuo Tiliæ in Jura.

93. **Phialea subgranulosa** (Rehm) Sacc. *Helotium subgranulosum* 1130

Rehm Ascom. Illustr. et Exsicc. n. 607. — Gregaria, minutissima cyathoideo-substipitata, albido-flavidula, scabra, sicca cyathoideo-inoluta, humida subexplanata, disco flavidulo c. 0,4 mm. diam., prosenchymatico-hyalino-contexta, fragilissima; sporidiis evoideis, hyalinis, $5 \approx 2,5$, 1-2-stichis; ascis clavatis, c. 30-36 ≈ 5 ; paraphysibus perpaucis, filiformibus.

Hab. ad corticem putrescentem Abietis excelsæ in fauibus montis Hochgrad alpium Allgoviæ, c. 1500 m. alt. (BRITZELMAYR). — A *Pseudohelot. hyalino* ascomate stipitato, sporidiis rectis etc. dignoscitur.

94. **Phialea straminella** (Karst.) Sacc. *Helotium straminellum* Karst. 1131

Symb. M. F. XXIII, pag. 27. — Subsparsa, obconica, stipitata, straminea vel subaurea, marginem versus tenuiter albido-furfurilla, 0,1-0,2 mm. alt., disco concaviusculo, stipite glabro, cupulam æquante; ascis cylindraceo-subclavatis, 38-40 ≈ 5 ; sporidiis cylindraceis, rectis, eguttulatis, 4-4,5 ≈ 1 ; paraphysibus discretis.

Hab. ad lignum Betulæ prope Mustiala Fenniæ. — *Ph. lectissimæ* affinis.

95. **Phialea Galbula** (Karst.) Sacc. *Helotium Galbula* Karst. M. F. I, 1132

p. 140. — Sparsa, stipitata, aureo-lutea, puberula, circiter 0,3 mm. alt.; cupula concava, sicca sphæroidea, 0,2-0,3 mm. lat.; stipite in cupulam dilatato, brevissimo; ascis cylindraceo-subclavatis, 28-30 ≈ 4 , jodo non cœrulecentibus; sporidiis subellipsoideis vel ovoideo-oblongatis, sæpe 2-guttulatis, 3-4 ≈ 2 ; paraphysibus parcis gracilibus.

Hab. ad corticem et lignum Alni incanæ pr. Mustiala Fenniæ.

96. **Phialea sulphurella** Sacc. Fung. Ven. Ser. IV, n. 83, f. ital. f. 1133

1328. — Cupulis gregariis, tenuissimis, 200 μ . altis, 120 latis, flavis, subdiaphanis, breve sensim stipitatis, glabris sed undique minutissime verruculoso-fibrillosis, disco excavato obtusiusculo marginato, excipuli contextu prosenchymatico tenerrimo; ascis oblongo-cylindraceis sessilibus apice rotundatis, brevibus, 48 $\approx 5-6$; paraphysibus filiformibus obvallatis, 8-sporis; sporidiis in ascii parte superiore distichis, oblongo-ovoideis, 4-5 ≈ 2 , eguttulatis, hyalinis.

Hab. in ramis decorticatis putrescentibus Quercus pedunculatae in silva Montello (Treviso). — A *Ph. Galbula* Karst. vix differt cupulis minoribus non vere puberulis, ascis sporidiisque paullulo majoribus.

97. **Phialea improvisa** (Karst.) Sacc. *Hel. improvisum* Karst. M. F. 1134
I, pag. 141. — Gregaria, stipitata; cupula sphæroideo-cyathoidea, extus præcipue marginem versus sulphureo-farinoso-puberula, 0,3-0,5 mm. lat.; stipite glabro; cupula breviore; ascis cylindraceo-clavatis, 35 \approx 4; sporidiis distichis, fusoideo-oblongatis, 4-6 \approx 1,5-2,5; paraphysibus gracilibus.

Hab. ad truncum abietinum mucidum loco uliginoso silvatico prostratum prope prædium Bukkila parœciæ Lemo Fenniæ.

98. **Phialea lectissima** (Karst.) Sacc. *H. lectissimum* Karst. M. F. 1135
I, p. 141. — Sparsa, substipitata, obconica, flava, marginem versus tenuiter albido-furfurella, disco planiusculo, sicco concavo, circiter 0,2 mm. lat.; ascis cylindraceo-clavatis, 40-56 \approx 6-7,5, obturaculo minutissimo jodo obsoletissime cœrulescente; sporidiis oblongatis vel fusoideo-oblongatis, rectis vel curvulis, 8-11 \approx 2-3; paraphysibus gracilibus.

Hab. ad corticem ramulorum Salicis phylicifoliæ exsicatorum in regione mustialaensi Fenniæ. — Stipes sursum sensim in cupulam dilatatus, *Ph. Galbulæ* et *Ph. albohyalino* affinis, externa facile subliciformem in memoriam aliquantulum referens.

99. **Phialea phyllogena** (Rehm) Sacc. *Helotium phyllogenum* Rehm 1136
Ascomyc. fasc. XVI, p. 7, n. 768. — Ascomatibus parenchymatico-hyalino-contextis, ex cyathoideo patellaribus, tenuiter stipitatis, stipite 1 mm. longo, extus pallide farinoso-flavidulæ, disco flavo, 1-1,2 mm. diam., margine dilutiore cinctis, planis, gregariis; ascis clavatis, subcurvatis, 8-sporis, 75 \approx 8; sporidiis distichis, ellipticis vel elongato-clavatis, continuis, crasse binucleatis, plerumque rectis, hyalinis, 12-15 \approx 3,5; paraphysibus filiformibus, apice vix incrassatis.

Hab. ad folia Populorum putrida prope Ungarisch Altenburg Hungariæ (LINHART).

III. Disco rubente, testaceo.

100. **Phialea calathicola** (Rehm) Sacc. *Helotium calathicolum* Rehm 1137
Ascom. n. 305. — Gregaria, crasse stipitata, cyathiformis, stipite c. 0,5 mm. crasso, 1 mm. long., recens extus fusco-rubella, sicca complicata, flavidocarnea, extus pilosula, humectata explanata, disco rubro vel rubello, 2-3 mm. diam., contextu prosenchymatico flavidulo; ascis clavatis, apice incrassatis, 8-sporis, 45 \approx 6; sporidiis ellipticis, hyalinis, fere monostichis, 7-8 \approx 3; paraphysibus filiformibus, superne subcurvatis, clavulatis, hyalinis.

Hab. ad calathides putridas *Cirsii spinosissimi* in alpibus Tyrolensis juxta glaciem æternam.

101. **Phialea dolosella** (Karst.) Sacc. *Helot. dolosellum* Karst. M. F. 1138

I, p. 137. — Albida vel testaceo-albida; cupula e cyathideo explanata, primo furfurella vel leviter puberula, disco sicco testaceo, circiter 1 mm. lat.; stipite tenui, æquali, circiter, 0,8 mm. alt.; ascis 50-60 = 5,5; sporidiis bacillaribus vel fusoideo-bacillaribus plerumque curvulis, tenuiter spuriæ uniseptatis, 10-16 = 1,5-2,5; paraphysibus gracilibus.

Hab. ad partem inferiorem culmorum Tritici repentis emortuorum, ad radices Rubi idæi, in caule Angelicæ in Mustiala et Abo Fenniæ, in Germania et Hungaria. — In forma graminum sec. Rehm asci 60-66 = 7-8; sporidia 15-17 = 2,5-3; in forma Angelicæ asci 45-50 = 5-6; sporid. 9-12 = 2. — * **Ph. albinea** Karst. l. c.: priori similis, sed sporidiis longioribus 14-25 = 1-2. Ad caules Galii veri aridos in Mustiala. — * **Ph. nivella** Karst. l. c., *Ph. albinea* similes sed minor, plantula 0,3 mm. alt., cupula 0,3 mm. lat., stipite altitudine cupulæ; sporidiis fusoideo-bacillaribus guttulatis vel tenuiter uniseptatis 16-18 = 2. Ad folia Caricis ampullaceaæ putrida in Mustiala.

IV. Disco brunneo, fusco, atro.

102. **Phialea concolor** (Phill.) Sacc. *Hymenoscypha* Phill. Disc. p. 139. 1139

— Hemisphærica, tenuis, firma, pallida vel sordide alba, extus pruinosa, disco pallide brunneo; margine minute fimbriato; stipite brevi, subobtuso; ascis tereti-clavatis; sporidiis oblongis vel sub-fusoideis, 8-10 = 2-3; paraphysibus filiformibus.

Hab. in ligno duro decorticato Forres Britanniæ (KEITH). —

Ascoma circ. $\frac{1}{4}$ -1 mm., stipite 0,5 mm. long.

103. **Phialea viridi-fusca** (Fuck.) Sacc. *Pez. viridi-fusca* Fuck. Symb. 1140

p. 309. — Cupulis stipitatis, plerumque obliquis, 2-4 mm. altis, stipite in cupulam sensim dilatato, turbinatis, 2 mm. latis, margine crasso, infexo, integerrimo, extus stipiteque amœne flavo-viridi furfuraceis, disco concavo, atro-fusco; ascis elongatis, stipitatis, 8-sporis, 64 = 8; sporidiis elongato-distichis, oblongo-ovatis, 8 = 4, hyalinis.

Hab. in amentis femineis putridis Alni glutinosæ pr. Oestrich Rhenogoviæ.

PEZIZELLA Fuck. Symb. p. 299, ex max. p. [1869] (Etym. demin. a

Peziza), *Mollisia* sect. *Dilutella* Phill. Br. Disc. p. 193, *Helotium* et *Mollisia* Auctor. p. p. — Ascoma ceraceo-membranaceum, tenue, urceolatum cupulatum v. discoideum, sessile, glabrum, leve, pallidum vel læte coloratum, excipulo quoque pallido et molli. Ascitereti-clavati, filiformi-paraphysati, octospori. Sporidia ellipsoidea oblonga, teretiuscula, hyalina, continua. — Est *Phialea* sessilis v. *Mollisia* læticolor, excipulo quoque pallido.

I. Disco albo, albido, pallente-flavido. (1-29)

1. **Pezizella albo-puncta** (Peck) Sacc. *Helotium albo-punctum* Peck 31 1141

Rep. p. 47. — Ascomate peregrino, albo, disco dein plano v. concavusculo, margine plerumque distincto; ascis cylindraceis; sporidiis biseriatis, oblongis, angustis, saepe binucleatis, 15-18 μ . long.

Hab. in foliis dejectis fagineis in silvis Adirondak Am. bor.

— Siccando flavescit et vix visible. Stipes tam brevis ut ascoma subsessile videatur.

2. **Pezizella glagosa** (E. et C.) Sacc. *Mollisia glagosa* E. et C. Journ. 1142

Myc. 1888, p. 56. — Minuta, orbicularis, sessilis, levis, lacteo-alba, sicca rufo-succinea; ascis clavulatis, 70-75 \approx 7-8; sporidiis subdistichis, oblongis vel clavato-oblongis, saepe curvulis, hyalinis, nucleatis simulateque 1-2-septatis, 8-10 \approx 2,5-3; paraphysibus filiformibus copiosis.

Hab. in foliis nondum putridis in paludosis New Field N. J. Am. bor. — Ascoma 250-275 μ . lat.

3. **Pezizella tumidula** (R. et D.) Sacc. *Peziza tumidula* Rob. et Desm. 1143

19 Not. p. 30. — Amphigena, erumpens, minutissima, sparsa aut laxe gregaria, sessilis, ceracea, glabra, albida, dein subplana, sicca concava ochroleuca v. argillacea, margine lacerato subdenticulata; ascis tubulosis; sporidiis hyalinis ellipsoideis subcylindricis.

Hab. in foliis semiputridis Betulæ in Gallia bor. — «An Cyathicula?»

4. **Pezizella leucella** (Karst.) Sacc. *Hel. leucellum* Karst. M. F. I. 1144

p. 151. — Gregaria vel sparsa, sessilis, planiuscula, subnuda, subimmarginata, hyalino-albida, sicca albida 0,1-0,2 mm. lat.; ascis clavatis, 45-65 \approx 6,5-8,5 obturaculo minutissimo jodo leviter vel obsolete cœrulecente; sporidiis distichis vel oblique monostichis, fusoideo-bacillaribus, rectis vel curvulis, guttulatis, 13-18 \approx 1,5-2,5; paraphysibus 1,5-2 μ . cr. apicem versus leniter incrassatis.

Hab. supra folia Betulæ, Salicis et Alni putrescentia vel putrida locis uidis in Fennia saltem australi.

5. **Pezizella minutissima** (Karst.) Sacc. *Helotium minutissimum* Karst. 1145
et Starb. Hedw. 87, pag. 124. — Sparsa vel subgregaria, sessilis;
cupula plana v. concaviuscula, nuda, pallescente, sicca pallida, di-
sco dilutiore, circiter 0,1 mm. lata; ascis clavatis; sporidiis.....
non descriptis.
Hab. in foliis putrescentibus Comari palustris prope Mu-
stiala Fenniæ (K. STARBACK).
6. **Pezizella pulvinata** (Karst.) Sacc. *Helotium pulvinalum* Karst. M. 1146
F. I, p. 140. — Gregaria, sessilis, convexa, fulvescens, disco albo,
sicco lutescente, circiter 0,5 mm. lat.; ascis cylindraceo-clavatis,
50-62 \approx 5-6, apice jodo non tinctis; sporidiis fusoideo-elongatis,
rectis v. leniter curvulis, eguttulatis, 6-10 \approx 1-2.
Hab. ad acus Pini silvestris putrescentes Mustiala Fenniæ.
7. **Pezizella dilutella** (Fr.) Fuck. Symb. pag. 300, *Peziza* Fr. Syst. 1147
Myc. II, p. 147, *Mollisia* Gill. et Phill. *Helotium xeroplasticum*
Rehm Ascom. n. 655. — Sessilis, gregaria, primitus subglobosa,
subclausa, sicca urceolata, flavo-ceracea v. albo-grisella, margine
integro pallidiore, humectata explanata v. subinfundibuliformis, fla-
vido-diaphana, $\frac{1}{2}$ -3 mm. diam., plerumque saltem 1 mm., prosen-
chymatico-contexta; ascis clavatis, apice incrassatis, 8-sporis, 36-
45 \approx 6 (Rehm), 24 \approx 6 (Fuck.); sporidiis oblongis, obtusis, rectis v.
subcurvatis, hyalinis, distichis, 8-10 \approx 1-1,5 (Phill.), 6-7 \approx 2-2,5
(Fuck. Oudem.); paraphysibus filiformibus superne subincrassatis,
2,5 μ . cr.
Hab. ad caules putridos Epilobii, Dahliæ, Rubi, Angelicæ, Adenostyolidis in Suecia, Germania, Britannia, Hollandia.
— Var. **Smyrnii** Phill. et Plow. Disc. p. 192: gregaria, sessilis,
hemisphærica dein expansa, carnosula, glabra, pallide argillacea,
margine minute fimbriato; ascis tereti-clavatis; sporidiis fusoideis
v. tereti-fusoideis, 3-guttatis, 10-13 \approx 3; paraphysibus filiformibus.
In caule emortuo Smyrnii, King's Lynn Britanniæ (PLOWRIGHT).
8. **Pezizella leucostigmoides** (Sacc.) Rehm Ascom. n. 855, *Calloria* 1148
leucostigmoides Sacc. Mich. II, p. 77, F, ital. f. 119. — Cupulis
superficialibus, sparsis, scutellaribus, minutissimis $\frac{1}{3}$ - $\frac{1}{4}$ mm., di-
lute chlorino-flavis, diaphanis, levissimis, glabris, jugiter (et in sicco)
apertis; ascis clavulatis, 30-45 \approx $3\frac{1}{2}$ - $4\frac{1}{2}$, octosporis; sporidiis cy-
lindraceis, curvulis, minutissimis, 4-5 \approx $\frac{3}{4}$ -1, hyalinis; paraphysi-
bus filiformibus, tenellis.
Hab. in caulibus herbarum majorum in silva Montello Ital. bor.,
in Gallia et Germania. — Maculis latis æqualiter atratis innasci-

tur. Ascoma, sec. Rehm, basi minute parenchymaticum, margine pseudoprosenchymaticum.

9. **Pezizella pulviscula** (Cooke) Sacc. *Peziza (Mollisia) pulviscula* 1149
Cooke Hedw. 1875, p. 84, Disc. U. S. I, p. 297. — Gregaria, mi-
nutissima, saccharina, subtremelloidea; cupula globosa, applanata,
glabra, pallida, aquoso-lutescente, 0,4 mm. lata; ascis cylindraceis
30 = 5; sporidiis minutissimis spermatoideis.

Hab. ad caules Phytolaccæ, New York Am. bor. (GERARD).

10. **Pezizella Oenotheræ** (C. et E.) Sacc. *Peziza Oenotheræ* C. et E: 1150
Grev. VI, pag. 90. — Sparsa, ochraceo-albida; cupulis demum
expansis, margine elevato, flexuoso; ascis cylindraceis; sporidiis
linearibus, rectis, obtusis 8 μ . long.; paraphysibus filiformibus.

Hab. in caulibus Oenotheræ, New Jersey Americæ bor. —
Ascoma 1 mm. lat.

11. **Pezizella albo-viridis** (Saut.) Sacc. *Peziza albo-viridis* Saut. Pilz. 1151
Salzb. p. 10. — Sessilis, minuta, submembranacea, glabra, concava,
albida, disco viridulo.

Hab. in sarmentis putrescentibus Rubi idæi, Salzburg. — Af-
finis *P. xanthostigmatis*. $\frac{1}{2}$ cm. lata. Sec. Wint. vix diversa a *M. dilutella*.

12. **Pezizella chlorinella** (Ces.) *Peziza chlorinella* Ces. Bot. Zeit. 1152
1854, pag. 186 et Kl. Herb. myc. 1812. — Minutissima, viridula,
gregaria, semipellucens.

Hab. ad herbarum caules, Vercelli Ital. bor. — *Pezizellæ leu-*
costigmoidi affinis videtur.

13. **Pezizella Carolinæ** (De-Guer.) Sacc. *Peziza Carolinæ* De-Guer. 1153
in Cr. Fin. p. 49. — Ascomate 2-3 mm. diam., albo, dein brun-
neo, hemisphærico, subpedicellato, primo concavo, dein plano, mar-
gine undulato; ascis clavatis, 8-sporis; sporidiis ovoideis; paraphy-
sibus filiformibus, sursum granulosis et brunneo-coloratis.

Hab. in ramulis emortuis Aceris, Morlaix Finistère Galliæ. 1154

14. **Pezizella vulgaris** (Fr.) Sacc. *Peziza vulgaris* Fr. Syst. Myc. II,
pag. 146, *P. albella* With. arr. IV, p. 350. — Sessilis, subcæspit-
tosa, membranacea, mollis, glabra, utrinque concolor, albido-palle-
scens v. roseo-expallens, orbicularis.

Hab. ad cortices ramorum exsiccatorum, v. c. Coryli, Rubi
in Suecia, Germania, Belgio, America bor., Cuba, Ceylon, Vera-
quaz. — Conferta, subcæspitosa, concaviuscula, orbicularis, sed ob-
mutuam pressionem flexuosa et quasi contorta, præcedente major,
sed vix lineam superans. Hymenium discretum nullum. *P. cera-*

cella pilis tantum paginæ inferioris differt. — Forte eadem est **Pezizella** (*Niptera*) **vulgaris** Fuck. Symb. App. II, p. 59: cupulis ascigeris sparsis seu gregariis, carnosso-diaphanis, mollibus, albidis, glabris, 1-3 mm. latis, sessilibus, primo urceolatis, demum concavo-subexplanatis, margine distincto, integro, regulari seu pressione repando, disco concolori; ascis elongato-clavatis, 8-sporis, $54 \pm 8-10$; sporidiis subdistichis, cylindraceo-subclavatis, rectis seu parum curvatis, continuis, hyalinis, $10-12 \pm 3-4$; paraphysibus copiosis, linearibus, simplicibus. In fragmentis (Zapfen) putridis Pini silvestris in Rhenogovia. Sec. Fuck. l. c., hujus status conidiophorus est superficialis, acervulos gelatinose molles, orbiculares, 1-2 mm. latos, convexos, demum planos, albido-cinereos formans; conidiis cylindraceis, rectis, continuis, $8-9 \pm 2$. — Var. **myceticola** Berk. N. A. Fung. p. 137: magis concava. In *Polyporis* vetustis ad *Quercus* in Carol. inf., Santee River et Britannia. — Var. **sanguinella** Berk. l. c.: extus alba, disco pallide aurantio. In *Liquidambare*, Carolina inf. et New England Amer. bor.

15. **Pezizella leucostigma** (Fuck.) Sacc. *Nipt. leucostigma* Fuck. Symb. 1155
App. II, p. 59 nec *Peziza leucostigma* Fries. — Cupulis sparsis, aquoso-diaphanis, 1 mm. lat., albis, sessilibus, orbicularibus, primo concavis, demum planis, glaberrimis, margine obscuro, disco concolori, siccis lutescentibus; ascis substipitatis, cylindraceis, 8-sporis, 48 ± 9 ; sporidiis oblique monostichis, cylindraceo-oblongis, parum curvatis, biguttulatis, continuis, hyalinis, $10-12 \pm 4$; paraphysibus filiformibus.

Hab. ad basim putridam truncorum *Fagi* v. *Betulae* supra Hallgarten Rhenogoviæ.

16. **Pezizella diaphanula** (Cooke) Sacc. *Peziza* (*Mollisia*) *diaphanula* 1156
Cooke Hedw. 1875, p. 84, Disc. U. S. I, p. 295. — Gregaria, minutissima, diaphana albida, pallescens, hemisphærica, demum applanata $\frac{1}{10}$ mm. lata; ascis clavatis; sporidiis elongato-ellipticis, uninucleatis, $17-20 \times 8$; paraphysibus linearibus vix incrassatis.

Hab. ad ligna vetusta, New Jersey Amer. bor. (ELLIS).

17. **Pezizella uvidula** (Karst.) Sacc. *Hel. uvidulum* Karst. M. F. I, 1157
p. 127. — Gregaria, sessilis, sphæroideo-pulvinata, immarginata, hyalina, glabra, 0,1-0,2 mm. lat.; ascis oblongatis, longe pedicellatis, $75-100 \times 18-20$, parte sporif. 40μ . long.; sporidiis distichis ellipsoideis v. ovoideo-ellipsoideis sœpe irregularibus, intus subgrumosis, $12-13 \times 6-7,5$.

Hab. ad lignum mucidum (*Betulae*?) loco udo prope Mustiala

Fenniæ. — Facillime prætervidenda ob summam minutiem (in sicco statu fere invisibilis); ab aliis *Pezizellæ* speciebus ascis et sporidiis pro ratione maximis optime distincta.

18. *Pezizella sordidula* (Speg.) Sacc. *Helotium sordidulum* Speg. 1158

F. Fueg. n. 336. — Densuscule sparsa, turbinato-calyciformis, minutissima, 300-400 μ . diam. subpellucida, disco infundibulari, dilutissime alutacea, margine acutissimo integro v. irregulariter erosodenticulato, extus sordide albo non v. vix tenuissime pruinuloso, sessilis v. pedicello brevissimo ac crassiusculo concolore suffulta; ascis cylindraceo-subclavulatis, sursum leniter attenuatis, apice subtruncato-rotundatis, crassuscule tunicatis, minute 1-foveolatis, deorsum attenuato-pedicellatis, 45-50 \times 3-3,5, octosporis; paraphysibus parcis, filiformibus, apice non v. vix subclavulatis, obtusis, obvallatis; sporidiis oblique monostichis, elliptico-cylindraceis, utrinque obtususcule rotundatis 4-4,5 \times 1,5, rectis v. inæquilateralibus, hyalinis, continuis.

Hab. ad corticem fagineum putrescentem peridermio denudatum in silvis prope Port Cook in Staten island Fuegiæ. — Iodi ope tunica ascorum dilute, obturaculus intensius cœrulescunt. Prope Ushuviaia specimina nonnulla notis externis perfecte cum typo congruentia, internis autem nonnihil recedentia (ascis 145 \times 3; paraphysibus apice non clavulatis, sporidiis minoribus 2,5-3,2 \times 1,5 μ .) legiit; cl. SPEGAZZINI an, ex ipso, species distincta?

19. *Pezizella soleniiformis* (B. et C.) Sacc. *Pez. soleniiformis* B. et 1159

C. North Am. Fung. n. 749. — Minuta, candida, primum hemisphærica, margine tumido, dein cylindrica, demum ore expanso flexuoso.

Hab. in ligno putri Alabama Amer. bor. (PETERS).

20. *Pezizella lactea* (Cooke) Sacc. *Helotium lacteum* Cooke Grev. 1160 VIII, p. 63. — Sparsa, lactea, sessilis; cupulis concavis, 2-3 mm. lat., margine flexuoso; ascis cylindraceis; sporidiis uniseriatis, lancelatis, nucleatis, 20-30 \times 4-6; paraphysibus filiformibus.

Hab. in ligno valde putrido, Maungaroa Novæ Zelandiæ. — Diaphana, sicca subochracea.

21. *Pezizella albella* (With.) Sacc. *Peziza albella* With. Arr. vol. 1161 IV, pag. 350, sec. Fr. *Pez. vulgaris* Fr. Syst. Myc. II, pag. 146, *Pezizella Avellaneæ* (Lasch) Fuck. Symb. p. 299, *Peziza Avellaneæ* Lasch in F. E. n. 28, *Helotium albellum* (With.) Karst. M. F. I, pag. 116. — Conferta, subcæspitosa, membranacea, sessilis, concaviuscula, albida v. albido-pallens, 1 mm. lat.; ascis cylindra-

ceo-clavatis, 35-45 \approx 5-6, filiformi-paraphysatis; sporidiis oblongo-teretiusculis, 6-9 \approx 1,5-2, rectis v. curvulis, hyalinis.

Hab. in ligno putri et cortice Coryli etc. in Suecia, Germania, Gallia, Fennia, Sibiria asiatica. — Var. **incarnata** Fr. l. c. In Rubo in Suecia et Germania.

22. **Pezizella Cyathus** (Nees) Sacc. *P. Cyathus* Nees in Mart. Erl. 1162 p. 463, Pers. Mycol. Eur. p. 287, Fr. Syst. Myc. II, p. 114. — Sessilis, subcylindrica, fusca, ore denticulato, connivente, disco albido.

Hab. in ramulis Betulæ albæ, Erlangen in Germania. — An *Mollisia*? Cupula membranacea, 1 mm. lata, extus rimuloso-substriata.

23. **Pezizella turgidella** (Karst.) Sacc. *Helot. turgidellum* Karst. M. 1163 F. I, p. 158. — Gregaria v. subsparsa, sessilis, convexa, immarginata, subnuda, albido-hyalina, sicca convexiuscula, subluteola, 0,2-0,4 mm. lat.; ascis fusoideo-clavatis, 25-34 \approx 5-7; sporidiis distichis vel oblique monostichis, fusoideo-elongatis, guttulatis, vulgo rectis, 7-13, plerumque 9-12 \approx 1,5-2; paraphysibus gracilescentibus.

Hab. ad culmos graminis putridos locis uidis in Fennia.

24. **Pezizella planodisca** P. et C. in Peck 31 Rep. Bot. Mus. p. 46. 1164 — Cupulis minutis, sessilibus, albidis, disco plano v. leniter convexo, margine oblitterato; ascis brevibus; sporidiis consertis v. distichis, subfusiformibus, 7,5-8,5 μ . longis.

Hab. in foliis emortuis Andropogonis scoparii, Buffalo Americæ borealis (CLIFTON).

25. **Pezizella microspis** (Karst.) Sacc. *Helotium microspis* Karst. M. 1165 F. I, p. 152. — Gregaria, sessilis, subnuda, convexa, subimmarginata, hyalino-albida, sicca planiuscula v. concaviuscula, albida, 0,1-0,2 mm. lat.; ascis clavatis, 25-35 \approx 4-5, apicè jodo non tinctis; sporidiis elongatis vel fusoideo-elongatis, plerumque 2-guttulatis, 6-8 \approx 1-1,5; paraphysibus vix ullis propriis.

Hab. supra folia Phragmitis communis putrescentia in Fennia usque ad oppid. Vasa.

26. **Pezizella Filicum** (Phill.) Sacc. *Mollisia* Phill. Disc. p. 191. — 1166 Gregaria, minuta, initio globosa, dein cupulata, demum expansa, alba, glabra; ascis tereti-clavatis; sporidiis oblongo-ellipsoideis, rectis v. curvulis bi-triguttatis, demum simulate 1-2-septatis, 8-12 \approx 3; paraphysibus tenuissimis.

Hab. in caule emortuo Aspidii Filicis-maris, Sohbden in

Britannia. — Ascoma 100-300 μ . lat. In Grevillea tamen (XIII, p. 74), ascoma indicatur 1-3 mm. lat., quod probabilius.

27. **Pezizella epithallina** (Ph. et Pl.) Sacc. *Mollisia epithallina* Ph. 1167 et Plow. Disc. p. 173. — Congregata, sessilis, disciformis, immarginata, albo-pallens v. flavidо-alba; ascis cylindraceo-clavatis, octosporis; sporidiis oblongo-ellipsoideis, 8 \approx 2; paraphysibus tenuibus, sursum incrassatulis.

Hab. in thallo Peltigeræ caninæ, King's Lynn Britanniæ.

28. **Pezizella Glæocapsæ** (Boud.) Sacc. *Epiglia Glæocapsæ* Boud. 1168 Disc. charn. pag. 25. — Ascomatibus minutis, $\frac{1}{4}$ - $\frac{1}{2}$ mm. dian., rotundatis, subturbanatis v. lenticularibus, albescensibus v. griseis, glabris; hymenio leviter papillato; ascis 110 \approx 10-12, paraphysatis; paraphysibus gracilibus, dichotomis et apice hamatis; sporidiis obtuse fusiformibus, granulosis, continuis, 12-16 \approx 5-6, hyalinis.

Hab. ad humum inter muscos, *Glaeocapsa* invasos, Montmorency in Gallia.

29. **Pezizella albula** (Phill.) Sacc. *Mollisia* Phill. Disc. pag. 192. — 1169 Gregaria, minuta, sessilis, concava, demum plana v. convexa, glabra, nitidula, albida; ascis clavatis; sporidiis subcylindricis v. fusoides, hyalinis, 6-10 \approx 2; paraphysibus filiformibus tenuibus.

Hab. in fimo cunicolorum Forres Britanniæ (KEITH). — Ascoma 300-400 μ . lat., carnosulum. Excipulum prosenchymaticum.

II. Disco roseo, rubro-aureo, lilacino.

30. **Pezizella Rhemii** Sacc. *Pezizella Aconiti* Rehm Ascom. n. 163, 1170 Lojk. n. 34, nec Saut. — Gregaria, sessilis, orbicularis, primitus subglobosa urceolata, dein plus minusve expansa, sicca albido-rosea, extus scabra, disco roseo, humida dilatata plana, margine crasso, albidio, denticulato cincta, roseo-cinerea, tenerrime, contextu subhyalino, prosenchymatico, c. 3-5 mm. diam. (humida); ascis clavatis, apice incrassatis, 8-sporis, 45-50 \approx 7, jodo cœrulescentibus; sporidiis subellipticis, obtusis, 2-plerumque 4 nucleis præditis, rectis vel subcurvatis, hyalinis, distichis, 12-14 \approx 3; paraphysibus filiformibus.

Hab. ad caules putridos Aconiti Napelli in alpe Dzurowa prope Teplicska, Liptau, Hungariæ (LOJKA).

31. **Pezizella Nectriella** (Karst.) Sacc. *Helotium Nectriella* Karst. 1171 Fragm. mycol. XIX, p. 5, Revisio Ascom. Fenn. p. 127. — Gregaria vel sparsa, subinde confluens, sessilis, convexa, subimmarginata, ceracea, incarnata, glabra, 0,3 mm. lat.; ascis clavatis, 33-

45 \approx 7; sporidiis distichis, elongatis, 6-7 \approx 1,5-2; paraphysibus filiformibus, apice incrassatis.

Hab. ad caules aridos Solani tuberosi in Mustiala Fenniæ.

32. **Pezizella exigua** (Cooke) Sacc. *Peziza exigua* Cooke Hedw. 1875, 1172 p. 83, Disc. U. S. I, p. 296. — Sparsa, minutissima, sessilis, sub-tremellosa, miniata, cupula hemisphærica, dein applanata vel convexa, margine vix elevato; ascis minutis lanceolatis; sporidiis linearibus hyalinis, vix 0,5 mm. long.

Hab. ad caules Erigeronis, New Jersey Amer. bor. (ELLIS).

— Oculo nudo inconspicua.

33. **Pezizella Rozei** (Quél.) Sacc. *Orbilia Rozei* Quél. Grev. X Suppl. 1173 p. 14, t. IX, f. 15. — Punctiformis 0,3 mm. lat., gibbosa, lobulata, granulosa, glabra, mollis, ex aurantio-incarnato roseola; sporidiis ellipsoideis 5 μ . long., deinde fusiformibus et guttulatis.

Hab. in caulibus emortuis Hellebori fœtidi, Jura Galliæ.

34. **Pezizella elæina** (Pass. et Beltr.) Sacc. *Niptera elæina* Pass. 1174 et Beltr. — Cupulis sparsis, depressis, pallide miniatis, margine (in sicco) undulato, truncato, nudo; ascis brevibus cylindricis, 8-sporis, jodo immutatis, 27-30 μ . long.; sporidiis tenuibus, bacillaribus, continuis, 12 \approx 4; paraphysibus ascos æquantibus.

Hab. in ligno indurato Oleæ europææ, Monostalla Siciliæ.

35. **Pezizella lilacina** Sacc. *Niptera lilacina* Sacc. Fung. Ven. Ser. 1175 IV, n. 70, *Ombrophila lilacina* (Wulf.) Karst. Myc. Fenn. I, 90?? — Cupulis gregariis, ceraceo-mollibus, plano-patellaribus, adpresso sessilibus, marginatis, $\frac{1}{2}$ -1 mm. diam., vivis amoëne lilacinis, siccis pallide lilacinis, disco pallidiore et ob ascos minutissime velutino; ascis copiosissimis, cylindraceo-clavatis, basi attenuato-stipitatis, 120 \approx 6-7, p. s. 50-60 longa; paraphysibus filiformibus obvallatis, 8-sporis; sporidiis in asci parte superiore subdistichis, ovoideis, 7-8 \approx 5, crassiuscule 1-2-guttulatis, hyalinis.

Hab. in ligno fagineo putrescenti in silva Cansiglio (Treviso)

Ital. bor. — Species certe pezizacea nec bulgariacea.

36. **Pezizella brassicæcola** (Berk.) Sacc. *Peziza brassicæcola* Berk. 1176 North Am. Fung. n. 736, Cooke Disc. U. S. I, p. 297. — Tenuis, expansa, flexuosa, extus intusque rufa; sporidiis ellipticis concatenatis («?»), uninucleatis, 10 μ . longis.

Hab. in caulibus Brassicæ putridis, New England Amer. bor.

(SPRAGUE). — Habitus *Pezizella vulgaris*.

37. **Pezizella rufula** (Schw.) Sacc. *Peziza rufula* Schw. Syn. p. 177. 1177 — Basi affixa, margine crasso subalbescente, ceterum e rubro rufa,

disco impresso, quasi umbilicato, limbo interdum undulato, plerumque regulariter orbicularis.

Hab. in lignis ad Bethlehem Amer. bor. — *P. rubellæ* affinis sed major et minus tremellosa. An *Calloria*?

38. **Pezizella elegans** (Saut.) Sacc. *P. elegans* Saut. Pilz. Salzb. p. 9. 1178

— Sessilis, glabra, gregaria, ceraceo-mollis, carnea, ascomatibus urceolatis.

Hab. in ligno putri prope Mittersill Salzburg. — Affinis *Pez. mycophilæ*. Ascoma $\frac{1}{2}$ -1 mm. lat.

39. **Pezizella hæmatoidea** (C. et Ph.) Sacc. *Peziza hæmatoidea* C. 1179 et Phil. Grevillea VIII, p. 62. — Sparsa, sanguineo-rubra, ceraceo-mollis, sessilis; cupulis demum applanatis, 1- $1\frac{1}{2}$ mm., margine elevato; ascis clavatis; sporidiis cylindricis, utrinque attenuatis, guttulatis $16 \approx 3$ -4; paraphysibus filiformibus.

Hab. in ligno decorticato Waitaki Novæ Zelandiæ. — *P. rubellæ* subaffinis.

40. **Pezizella carneo-rosea** (C. et H.) Sacc. *Peziza carneo-rosea* 1180 Cooke et Hark. Grev. IX, pag. 130. — Discoidea, sessilis, carneo-rosea, subcarnosa, $\frac{1}{3}$ mm. diam.; ascis cylindraceis; sporidiis ellipticis $5 \approx 3$; paraphysibus linearibus.

Hab. ad ramos Eucalypti in California.

41. **Pezizella rosea** (Mont.) Sacc. *Niptera rosea* Mont. Fl. Chil. VII, 1181 p. 415, t. 10, f. 7, Syll. Crypt. n. 636. — Cupula rosella, variaeformis, turbinata vel obovata, supra plano-concava, submarginata, margine tenui vel crassiusculo.

Hab. ad ramos decorticatos, Valdivia (CL. GAY).

42. **Pezizella regalis** (C. et E.) Sacc. *Peziza regalis* C. et E. Grev. 1182 VI, pag. 91. — Sparsa, miniata; cupulis sessilibus, subhemisphaericis, demum expansis; margine elevato, albo-crenato; ascis cylindraceo-clavatis; sporidiis linearibus, rectis 7μ . longis; paraphysibus superne globoso-clavatis.

Hab. in cortice pirino, New Jersey Amer. bor. — Ascoma 1 mm. lat. *P. lasiæ* affinis.

43. **Pezizella cruenta** (Schw.) Sacc. *Peziza cruenta* Schw. Syn. 177. 1183 — Minuta, sparsa, sessilis, applanata, subpellucida, margine elevato albidulo aut pruina adsperso; disco concavo, forma totius fungi orbiculata, vix 1 mm. excedens.

Hab. in cortice ramorum, Bethlehem Amer. bor. — An *Calloria*?

44. **Pezizella conchella** (Schw.) Sacc. *Peziza conchella* Schw. Syn. 1184

Amer. bor. p. 177, — Minutissima, conchiformis, sessilis, subpellucida, rubra, sicca, connivens, margine acuto.

Hab. ad ramos dejectos Mori albæ, Bethlehem in Pennsylvania Amer. bor.

45. **Pezizella carneo-pallida** (R. et D.) Sacc. *Peziza carneo-pallida* 1185

Rob. et Desm. 19, Not. pag. 31. — Amphigena, minutissima, plerumque laxe gregaria, sicca epidermide tecta, humida, erumpens, sessilis, subimmersa, rotundata, crassa, pallide carneæ; disco convexo fere tremelloso, margine subdenticulato; ascis mediocribus, clavatis, 50 μ . long.; sporidiis hyalinis, continuis, ellipsoideis, subcylindricis, obtusis, 15 μ . long.

Hab. in foliis semiputridis Ulmi in Gallia bor. — Ascoma $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam.

46. **Pezizella incarnata** (Cooke) Sacc. *Pez. incarnata* Cooke Grev. I, 1186

p. 131. — Sparsa, minuta incarnata, sessilis, mox expansa, fere explanata, extus pallida; ascis cylindraceis; sporidiis linearibus, obtusis, hyalinis, 10 \approx 1,5.

Hab. in foliis Pinorum in Scotia (A. JERDON).

47. **Pezizella Cucurbitæ** (Ger.) Sacc. *Peziza Cucurbitæ* W. K. Ger. 1187

Torr. Bot. Club. Jun. 1874. — Ascomatibus sessilibus, udis ceraeis, scutatis, margine integro, disco pallide brunneo siccis contortis subcarneis, disco atro-purpureo; sporidiis ovoideis continuis, 10 μ . longis.

Hab. in epicarpio putri Cucurbitæ, sociâ *Phoma Cucurbitacearum*, Poughkeepsie Amer. bor.

48. **Pezizella phymatodes** (Phill.) Sacc. *Mollisia phymatodes* Phill. 1188

in Grevillea V, p. 517, t. 88, fig. 9. — Sessilis, sparsa, sphærica, rubescenti-carnea, glabra, subtiliter striata, margine tenui, serrato, pallidiori; ascis subclavatis; sporidiis oblongo-fusoideis, continuis, 13-16 \approx 3-4; paraphysibus filiformibus.

Hab. in culmis, Blue Canon Sierra, Nevada Californiæ.

III. Disco flavo, aureo, flavido.

49. **Pezizella aureola** (Rabenh.) Sacc. *Peziza Rabenh.* F. E. n. 1622, 1189

Hedw. 1873, pag. 139. — Sessilis sparsa v. subsparsa humiditate expanso-patelliformis, carnosò-mollis, fusco-ferruginea, distincte plicato-striata; margine leviter inflexo, crenulato subdenticulata, siccitate involuta, subclausa, disco subaureo; ascis linearibus, 6 μ . crassis, octosporis; sporidiis subcylindraceis, monostichis, rectis, utrinque obtusis; paraphysibus nullis.

Hab. ad caules sicclos Eupatorii cannabini Dresden Germaniæ.

50. **Pezizella hungarica** Rehm Ascom. Lojk. n. 35, cfr. Winter in ¹¹⁹⁰ Flora 1872, p. 526. — Minutissima, c. 0,3 mm. diam., gregaria, sessilis, primitus globosa, dein urceolata, sicca flavidula, humectata hyalinella, prosenchymatico-contexta; ascis clavatis, 8-sporis, 30-4; sporidiis elongato-cuneiformibus, rectis, hyalinis, obtusis, 5-7-1,5, distichis; paraphysibus teneris, filiformibus; jodii ope apice ascorum cœrulecente.

Hab. ad caules putridos Umbelliferarum in horto Esterházyano Kaposvár (Somogy) Hungariæ (LOJKA). — Var. **Heraclei**: differt sporidiis subcurvatis, 6-8-1, paraphysibus apice globosa clava c. 3 μ. crassa instructis. Ad caulem putridum Heraclei palmati infra montem Retyezát (Hunyad) Transylvaniæ (LOJKA).

51. **Pezizella tyrolensis** Rehm Ascom. Lojk. n. 36. — Plerumque ¹¹⁹¹ sparsa, rarius gregaria, sessilis, primitus globosula, dein substipitata, serius explanata, sicca junior aureo-vel aurantiaco-flavida, scabra, concava, plicata, margine, subfimbriato et rugoso, dein explanata, disciformis, haud distincte marginata, margine albido-fimbriata, humida flava, expansa, prosenchymatico-flavidulo-contexta, 1-1,2 mm. diam.; ascis clavatis, apice incrassatis, 8-sporis, 40-60-6-8 jodo cœrulecentibus; sporidiis oblongis vel obtuse oblongatis, ellipticis, hyalinis, rectis, saepe 1-2 nucleolis præeditis 6-9-2,5; paraphysibus filiformibus, apice haud clavatis, hyalinis, p. p. crassis ascosque superantibus, c. 2 μ. diam.

Hab. ad caules putridos Angelicæ circa lacum Zenoga infra alpem Retyezát (Hunyad) Transylvaniæ (LOJKA), et Cirsii spinosissimi in Alpibus tyrolensis. — *Pezizella hungarica* Rehm huc forte pertinet, ex ipso auctore.

52. **Pezizella Sedi** (Karst.) Sacc. *Helotium Sedi* Karst. Rev. monogr. p. 128. — Sparsa vel subgregaria, sessilis aut distincte stipitata; cupula planiuscula immarginata, luteo-rubella, sicca subaurantiaca mollis, sicca cornea, glabra, 0,3-0,5 mm. lat.; ascis crassule clavatis, deorsum in pedicellum pedetentim attenuatis, apice obtusissimis, jodo haud cœrulecentibus, 100-110-18-20; sporidiis distichis, ellipsoideis, massa minute granulosa fœtis, eguttulatis, hyalinis, 17-21-9-10; paraphysibus filiformibus, ramosis, sat confertis gracilibus.

Hab. in Sedo acri putrido prope Mustiala Fenniæ.

53. **Pezizella crocitincta** (B. et C.) Sacc. *Peziza crocitincta* B. et ¹¹⁹³

C. North Am. Fungi n. 749, Grev. t. I, fig. 5. — Sessilis, mollis, primum globosa dein expansa, extus glabra, rivulosa, vivide croci-color; hymenio concavo citrino; ascis cylindraceis; sporidiis allantoideis exiguis, luteolis.

Hab. ad lignum quercinum putre in Georgia, Pennsylvania, Poughkeepsie, North Greenbush Amer. bor. — Conferenda cum *Chlorosplenio Chlora* Schw.

54. **Pezizella translucens** (Gill.) Sacc. *Mollisia translucens* Gill. Disc. 1194 p. 212 cum icono Pat. Tab. pag. 36, n. 88. — Sessilis, expansa, tenuissima, margine undulato, flavo-straminea, translucida, basi hyphulis albidis, septatis prædita; ascis cylindraceis; sporidiis filiformibus brevibus curvulis, $4 \approx 0,5$.

Hab. in ligno putri Quercus et Fagi, Alençon et Poligny Galliæ.

55. **Pezizella floriformis** (Peck) Sacc. *Mollisia floriformis* Peck 33 1195 Rep. 31. — Minuta, tenuis, glabra, cyathiformis, regularis, dein ob marginem undulato-inflexum floriformis, sordide flava; ascis brevibus cylindricis, $18-20 \mu$. longis; sporidiis spermatoideis.

Hab. in ligno dejecto Aceris Verona Amer. bor. — Ascoma 1-2 mm. lat.

56. **Pezizella infixa** (Wallr.) Sacc. *Pez. infixa* Wallr. Crypt. Germ. 1196 n. 2504. — Ascomatibus sessilibus glaberrimis exiguis sparsis, basi punctiformi insculptis, primum urceolatis, dein discum concavum centro saturatiore notatum, margine elevato conformi patulo cinctum recludentibus, vivis pallide cerinis subtremellosis, siccis aurantio-flavis.

Hab. ad ligna salicina putredine exesa in Thuringia. — Adfigendi modo peculiaris.

57. **Pezizella excavata** (Saut.) Sacc. *P. excavata* Saut. Pilz. Salzb. 1197 p. 60. — Exigua, sessilis, gregaria, submembranacea glabra, lutea, ex cavata, margine repando, undulato.

Hab. in ligno putri prope Mittersill Salzburg. — Affinis *M. leucostigmati* $\frac{1}{2} \mu$. lat.

58. **Pezizella citrinella** (Schw.) Sacc. *Peziza citrinella* Schw. Syn. 1198 p. 177. — Gregaria ac sparsa minuta, sessilis junior subglobosa, demum applanata, semper margine crassiusculo obtuso distincta varie flexuoso et linea albida notato, ceterum unicolor citrinella.

Hab. in lignis cariosissimis saliciniis, Bethlehem Amer. bor. — Affinis *Call. chrysocomæ*: an ergo *Calloria*?

59. **Pezizella crispa** (C. et Phill.) Sacc. *Peziza crispa* C. et Phill. 1199

Grev. VIII, p. 62. — Aurantiaca, sparsa, sessilis, 1-4 mm. lat.; cupulis concavis, crispulis, subtus pallidioribus, tenuibus; ascis cylindraceis; sporidiis linearibus, 8-3 utrinque rotundatis.

Hab. ad lignum, Maungaroa Novæ Zelandiæ.

60. **Pezizella exidiella** (B. et C.) Sacc. *Peziza exidiella* B. et C. 1200

North Am. Fung. n. 740 et Cooke Disc. U. S. I, p. 297. — *Gregaria regularis* extus intusque flavo-rufa; ascis clavatis; sporidiis oblongis angustis hyalinis.

Hab. ad ramos Corni floridæ in Carol. inf. Amer. bor.

61. **Pezizella Aquifolii** (C. et E.) Sacc. *Peziza Aquifoliae* C. et E. 1201

Grev. VI, pag. 91. — Foliicola, sparsa, subaurantia, ceracea; cupulis sessilibus, concavis, margine incurvo, ad basim fibrillis albis radiantibus affixis; ascis cylindraceis; sporidiis arcte ovalibus, hyalinis 7-10 ≈ 2,5; paraphysibus vix distinctis.

Hab. in foliis emortuis Ilicis opacæ, New Jersey Amer. bor.

62. **Pezizella citrinula** (Karst.) Sacc. *Helot. citrinulum* Karst. M. F. I, 1202

pag. 139. — Sparsa, sessilis plana, subaurantiaco-lutea vel citrina, 0,7 mm. lat.; ascis cylindraceo-clavatis, 45-60 ≈ 6, apice jodo vix vel obsolete cœrulecentibus; sporidiis fusoideo-elongatis, curvulis vel subrectis, vulgo guttulis paucis (plerumque 2 apicalibus) præditis, 6-12 ≈ 1,5-3; paraphysibus parcis, gracilibus sursum leviter incrassatulis.

Hab. in culmis graminum exsiccatis pr. Helsingfors et Mustiala Fenniæ et in vaginis culmisque Phragmitis communis in Gallia. — Pulchra et facile distincta species, cupula carnosoceracea, integerrima. Ad *Ph. alabastrinam* Fr. ut videtur, vergit. Ascomata (in specim. gallicis) sessilia, patellulata, integra v. subsinuosa, glabra, luteola, ceracella; asci teretiusculi 50-55 ≈ 4-5, paraphysati, octospori; sporidia elongata, curvula 8-9 ≈ 2-2,5, hyalina.

63. **Pezizella flaveola** (Cooke) Sacc. *Peziza flaveola* Cooke Grev. I, 1203

p. 131. — Sessilis, minuta, vivide flava, submembranacea tenera, mox discoidea glabra levis; ascis cylindraceis; sporidiis subfuscoides rectis v. curvulis 8 ≈ 2.

Hab. in frondibus Pteridis aquilinæ King's Lynn Britanniæ (PLOWRIGHT).

64. **Pezizella chrysostigma** (Fr.) Sacc. *Peziza chrysostigma* Fr. S. 1204

M. II, p. 128 (*Peziza*), *Helot. chrysostigma* Fr. Summa p. 355, *Calloria chrysostigma* Phill. Disc. p. 328, *Pez. mellina* Pers. M. E. I, p. 290 (f. major), *Pez. versicolor* Desm. — Minima, punctiformis, aurea, disco plano immarginato, stipite brevissimo; ascis

elongato-clavatis, stipitatis, 8-sporis, 25-40 = 6-8; sporidiis oblique monostichis, oblongo-ovatis, inæquilateralibus, 4-guttulatis, hyalinis, continuis, 10-12 = 2-3; paraphysibus filiformibus.

Hab. ad stipites putrescentes *Aspidii Filicis-maris et Filicis-feminæ* locis umbrosis in Suecia, Germania, Fennia, Italia. — Pulcherrima, nudo oculo vix discernibilis. Stipes constanter adest, paullo dilutior, sursum dilatatus in cupulam planam, margine nullo prominente cinctam. Ascomata (in exempl. italicis) per exigua, vix ac ne vix puberula. Ascii clavulati $35 = 6\frac{1}{2}-7$, tetraspori (mihi saltem visi!), paraphysibus bacillaribus, raro furcatis obvallati; sporidia curvula, 9-11 = 2, hyalina. Ab hoc *H. aureolum* meum differt ascomatibus nitidis, sporidiis octonis, multo longioribus etc.

65. **Pezizella? Hypnorum** (Fr.) Sacc. *P. Hypnorum* Fr. Syst. Myc. II, 1205 p. 149, Wallr. Crypt. n. 9489. — Sessilis, minor, cupulata, sicca, lutea, margine subflexuoso.

Hab. in silvis ad saxa majora irrigua supra *Hypnum cressiforme* etc. in Suecia et Germania. — Sparsa, parce obvia, 1 mm. lata, glabra, demum convexa et irregularis. Scutella *Biatoræ* refert nec cum *P. muscorum* commutanda.

66. **Pezizella minima** (Fr.) Sacc. *P. minima* Fr. Elench. II, p. 13. — 1206 Minima, globosa, levissima, glaberrima, firmula, spadicea, ore connivente albido.

Hab. in *Hydnofarinaceo* copiose hieme primoque vere in Suecia. — Species bella, eximia, omnino libera, fere mobilis, intense spadicea, pæne nigrescens, nitida!, urceolata persistens. Nudo oculo inconspicua.

IV. Disco virescente.

67. **Pezizella mycophila** (Fr.) Sacc. *P. mycophila* Fries Obs. I, p. 164 1207 Summa p. 353. — Sessilis, ceraceo-mollis, glabra, pallide viridis, cupulis concavis marginatis.

Hab. in *Russula adusta* putrida, semel visa, in Suecia. — An, ex ipso auctore, eadem ac *Hum. Jungermanniæ?* Conferta, passim confluens, *Hum. Jungerm.* in mentem revocans sed tenuior ceracea. *Pez. diluta* γ. *fungorum* A. S. pag. 338 ad præsentem accedit, sed cupula substipitata roseo-rubella; hinc propria species.

V. Disco fulvo, rufo, brunneo-flavo, cinereo-rufo.

68. **Pezizella versatilis** (Fr.) Sacc. *Pez. versatilis* Fr. Summa p. 353. 1208

— Ascomate sessili, ceraceo-membranaceo, concavo, dilute fulvo, mycelio brevi radiante cincto, hymenio exoleto, reticulato-poroso.

Hab. ad folia betulina putrida coacervata in Suecia. — Ascoma 2-3 mm. lat., margine in siccis involuto et flexuoso, mycelium maculam orbicularem sistit.

69. **Pezizella Spirææ** (R. et Desm.) Sacc. *Pez. Spirææ* R. et Desm. 1209

24 Not. p. 7. — Hypo-rarius epiphylla, sessilis, minutissima, ceraceo-mollis, glabra, sparsa, punctiformis, junior globosa, adulta aperta, concava v. plana, sicca rufo-brunnea, humida disco subcinerescente, margine subdenticulato; ascis clavatis, rectis v. arcuatibus, 60 μ . longis; sporidiis ellipsoideis, 2-guttulatis, 10 \times 5.

Hab. in foliis vetustis Spirææ Ulmariae in Gallia bor. — Ascoma $\frac{1}{5}$ - $\frac{1}{4}$ mm. diam.

70. **Pezizella Leguminum** (Schw.) Sacc. *Pez. Leguminum* Schw. Syn. 1210

Amer. bor. p. 176. — Minuta, sparsa, emergens, globosa, ceracea, sessilis, glabra, rufa, disco excavato concolori, margine albescente.

Hab. in leguminibus Catalpæ et Tecomæ, Bethlehem Amer. bor., parce.

71. **Pezizella Pastinacæ** (Schw.) Sacc. *Peziza Pastinacæ* Schw. Syn. 1211

pag. 176. — Miniata, stipite brevissimo crasso, interdum sessilis, cupulis sparsis apertis, margine tumidulo obtuso, albescenti-luteo, disco subconvexulo rufo-lutescente.

Hab. in caulinis exsiccatis duris Pastinacæ, Bethlehem Americae borealis.

72. **Pezizella Aconiti** (Saut.) Sacc. *Peziza Aconiti* Saut. in Rabenh. 1212

Crypt. Fl. I, p. 344 et Wint. in Hedw. 1881, p. 134. — Sparsa, erumpens, punctiformis, subclausa, initio orbicularis, dein globoso-depressa, brunneo-flava, expallens, corrugata, margine inflexo; ascis anguste clavatis, 70-75 \times 5; sporidiis oblongo-fusoideis, hyalinis, 7-11 \times 1,5.

Hab. in caulinis emortuis Aconiti, Pinzgau.

73. **Pezizella juncina** (Pers.) Fuck. Symb. Myc. p. 299, *Peziza jun-* 1213

cina Per. Myc. Eur. I, p. 314. — Sessilis, sparsa glabra, subceracea, applanata, margine prominulo, obscure rufa; ascis stipitatis, oblongis, 8-sporis, 32 \times 4; sporidiis distichis, cylindraceis, 8 \times 2, quandoque curvatis.

Hab. in calamis putrescentibus Junci effusi et J. conglomerati prope Estrich Rhenogoviae et in Belgio.

74. **Pezizella lutiseda** (Karst.) Sacc. *Helotium lutisedum* Karst. M. 1214

F. I, pag. 126. — Sparsa, sessilis, convexa, testacea vel subflava,

demum obscurior, fulvo-fusca, circiter 1 mm. lat.; ascis cylindraceo-clavatis, 100 \approx 9-10, apice summo jodo haud v. leviter cœrulecentibus; sporidiis oblique mono- v. distichis, fusoideo-oblongatis, plerumque inæquilateralibus, 13-21 \approx 3,5-4,5, guttulis 2-pluribus præditis; paraphysibus gracilibus.

Hab. supra terram locis uliginosis prope Mustiala Fenniæ.

PSEUDOHELOTIUM Fuck, Symb. mycol. p. 298 [1869] (Etym. *pseudes* falsus et *Helotium*), *Mollisia* sect. *Mollisiella* Phill. Disc. p. 193, *Helotii* spec. Karst. — Ascoma ceraceum, tenue, cupulatum v. scutellatum, sessile v. subsessile, extus brevissime puberulum; granulosum vel pruinatum. Ascii tereti-clavati, filiformi-paraphysati, octospori. Sporidia hyalina, oblonga vel teretiuscula vel (in subg. *Mollisiella*) globulosa.

A. Pseudohelotium: sporidiis tereti-oblongis.

I. Disco albo, albido, mox sæpius flavicante v. fuscescente.

1. **Pseudohelotium hyalinum** (Pers.) Fuck. Symb. p. 298, *P. hyalina* 1215 Pers. Syn. p. 655, Myc. Eur. pag. 316 excl., Nees Syst. f. 270, extus granulosa, *Helotium* Karst. M. F. I, pag. 144, *Lachnella* Phill. Disc. p. 267, t. VIII, f. 48. — Sessile, punctiforme, subglobosum, humidum vitreo-pellucidum, extus puberulum; ascis linearioribus, 8-sporis, 24 \approx 4; sporidiis oblongis, hyalinis, 6-9 \approx 2-3.

Hab. ad truncos putridos (ligna et cortices) in Suecia, Germania, Britannia, Gallia, Italia, Belgio, Amer. bor., Cuba, Ceylon, Tasmania. — Juniore ætate subglobosum clausum, demum appannatum marginatum, variat irregulare, aquose albidum vel omnino candidum.

2. **Pseudohelotium microstomum** (Wallr.) Sacc. *Pez. microstoma* Wallr. 1216 Fl. crypt. n. 2589, ? *Peziza hyalina* β. *granulata* Pers. myc. II, p. 316, *P. hyalina* Nees syst. f. 270, nec Pers. — Ascomatibus sessilibus, exilissimis, albidis, extus concoloribus, tenuiter papulosis, dein hemisphæricis in osculum circinatum immarginatum angustum dilutum reclusis, urceolaribus.

Hab. ad Coryli truncos vetustos in Thuringia.

3. **Pseudohelotium vitreolum** (Karst.) Sacc. *H. vitreolum* Karst. M. 1217 F. I, p. 145. — Gregarium, sessile, convexulum, immarginatum, pilosellum vel demum subglabrum, hyalinum vel aquose albidum, siccum hemisphærico- vel disformi-contractum, lutescens vel lutescenti-pallidum, 0,4-1 mm. lat.; ascis cylindraceo-clavatis vel cla-

vatis, 60-65 \approx 8-11, apice jodo vix tinctis; sporidiis plerumque distichis, fusoideo-elongatis, rectis vel leniter curvulis, guttulas 2-4 sat minutas, rudes foventibus, 10-15 \approx 3-4; paraphysibus haud bene discretis, gracilibus.

Hab. ad lignum putrescens Betulæ et Padi in Fennia meridionali. — A *P. hyalino*, cui proximum, ascis sporidiisque majoribus, pilis cupulæ subtilioribus et brevioribus (30-40 μ . long.), dein evanescentibus aliisque notis differt. Ascomata interdum confluunt.

4. **Pseudohelotium Bullii** (Sm.) Sacc. *Peziza* Sm. in Gard. Chron 1873, 1218
Mollisia Phill. Disc. p. 194, tab. 6, fig. 35, Grevil. I, p. 120, t. 8, f. 3. — Subhemisphæricum, demum inæquale, sessile vel brevissime stipitatum, albidum, margine infexo, pulverulentum, disco sæpe proliero; ascis subclavatis; sporidiis subellipticis minutis 5-7 \approx 3.

Hab. in aquæ ductibus ligneis in Britannia. — Ascoma $\frac{1}{3}$ mm. lat.

5. **Pseudohelotium ceracellum** (Fr.) Sacc. *Peziza ceracella* Fr. S. 1219
M. II, p. 114. — Cupulis gregariis, ceraceis, duriusculis, $\frac{1}{2}$ - $1\frac{1}{4}$ mm. diam., subsessilibus, subtus turbinatis v. prorsus sessilibus, patellaribus, ex albo pallidissime (præcipue disco) flavescentibus, extus margineque setulis tenuissimis subvelutinis; disco e concavo plano marginato; ascis cylindraceis, breve attenuato-stipitatis, 80 \approx 5-6, paraphysatis, 8-sporis; sporidiis oblique monostichis v. distichis fusoideis v. subclavulatis, 10-12 \approx $2\frac{1}{2}$, hyalinis.

Hab. ad ligna populea et quercina in Suecia, Germania, Italia. — Sec. Fries l. c.: gregarium, subinde cæspitosum, tenue, margine tenuissime ciliato; discus albidus, 2-4 mm. latus, junior et siccus urceolatus, humidus expansus planiusculus subrepandus.

6. **Pseudohelotium opalinum** (Quél.) Sacc. *Mollisia opalina* Quél. J2 1220
Suppl. p. 13, tab. VII, f. 9 et in Grevillea 1879, p. 177. — Ascomate discoideo, 1 mm. diam., ceraceo, firmo, pruinoso, glabro, dia-phano, hyalino vel opalino; hymenio plano, lilacino vel glauco, in sicco amethysteo; sporidiis bacillaribus, 12 μ . longis, hyalinis.

Hab. in ramis putrescentibus in turfosis, Alsace.

7. **Pseudohelotium firmulum** (Karst.) Sacc. *Helotium firmulum* Karst. 1221
Symb. XXVII, pag. 32. — Gregarium, sessile, subiculo dematiego (*Tapesia fusca*?) insidiens, sphæroideum vel late obovoideum, ore minuto apertum, prosenchymatico-contextum, firmulum, album vel lutescens, albo-puberulum, demum cupulari-apertum et sordidatum, 0,2-0,3 mm. latum; ascis cylindraceo-clavatis, 6-7 μ . crassis; sporidiis distichis, fusoideo-elongatis, rectis vel curvulis, continuis,

eguttulatis, hyalinis, $7-12 \approx 1,5$; paraphysibus filiformibus coalescentibus.

Hab. in caulis aridis Rubi idæi in ditione mustialensi Fenniæ. — *Lachn. inquininæ* proximum.

8. **Pseudohelotium scrupulosum** (Karst.) Sacc. *Helotium scrupulosum* Karst. M. F. I, p. 152. — Gregarium, tenue, sessile, subhemisphæricum v. subsolenioideum, albidum v. lutescenti-albidum v. cinerescens, papillose exasperatum, circiter 0,2 mm. lat.; ascis cylindraceo-clavatis vel subcylindraceis, $20-30 \approx 3,5-4,5$, obturaculo jodo vix cœrulescente; sporidiis elongato- vel aciculari-elongatis plerumque 2-4-guttulatis, rectis, $4-8 \approx 1-2$; paraphysibus gracilimis.

Hab. supra caules Rubi idæi et R. odorati putrescentes in Mustiala in Thornwood Britanniæ (MASSEE). — Var. **Carpini**: ascomate sessili, suburceolato, per exiguo, flavidò, margine pallidiore, pruinoso-puberulo; ascis $35-38 \approx 4$, filiformi-paraphysatis; sporidiis 8, suballantoideis, $4-6 \approx 1$, hyalinis. In ligno decorticato Carpini, Fontainebleau (THERRY). — Var. **caulium**: ascis brevioribus et crassioribus, nempe $24-28 \approx 5$; sporidiis $5-6 \approx 1,5-2$. In caule Spirææ Ulmariae Rouen (LETENDRE).

9. **Pseudohelotium cæspitosum** (Bres.) Sacc. *Helotium cæspitosum* Bres. Fung. Trid. p. 70 et 105, t. 75, f. 2 a. — Ascomatis sessilibus, dense gregariis, ceraceo-carnosis, e concavo explanato-convexis, glabris v. vix in junioribus extus puberulis, albidis, in sicco lutescentibus, $\frac{1}{2}-1$ mm. latis, substantia concolore; ascis clavatis, jodo obturaculo minutissimo obsolete cœrulescente, $60-85 \approx 6-7$; paraphysibus apice clavato, guttulis flavis majusculis repletis, $3-3\frac{1}{2} \mu$. crasso; sporidiis subfusoideis, rectis v. curvulis, hyalinis, $10-12 \approx 3$.

Hab. ad caules sicclos Aconiti Napelli in alpibus Monte Peller in Valle di Sole. — Videtur *Helotio carnosulo* Rehm affine. Subiculum: **Chalara Aconiti** Bres. l. c., cæspitulis atro-olivaceis, congestis, ex hyphis subcylindraceis, basi attenuatis, interdum ramosis, septatis, fuligineo-olivaceis, $100-120 \approx 5-7$ compositis, conidiis, hyalinis, cylindraceis, utrinque truncatis, biguttulatis $8-10 \approx 2$.

10. **Pseudohelotium deparculum** (Karst.) Sacc. *Helotium deparculum* Karst. M. F. I, p. 150. — Gregarium, primitus sphæroideum, dein planiusculum, siccum hemisphæricum v. concavum, sessile, surfurello-puberulum, pallidum v. flavo-pallidum, siccum ochraceo-luteum v. testaceo-flavum; $0,3-0,4$ mm. lat.; ascis cylindraceo-cla-

vatis, 4-sporis, 30-45 \approx 4-5, apice jodo non cœrulecentibus; sporidiis fusoideo-bacillaribus, rectis vel curvulis, simplicibus vel spuriis tenuiter septatis, 12-15 \approx 1,5; paraphysibus parcis gracilibus.

Hab. ad caules Spirææ Ulmariæ exsiccatos in Fennia et Germania.

11. **Pseudohelotium effugiens** (Desm.) Sacc. *Hel. effugiens* (Rob.) Karst. 1225

M. F. I, p. 149, *P. effugiens* Desm. XXII, Not. p. 19. — Sparsum, sessile, o sphæroideo-hemisphæricum, in statu humido et vegeto concaviusculum vel planiusculum, siccum subsphæroideo-contractum, farinoso-subtomentellum, tenue, albidum, siccum lutescens, margine subtilissime fimbriato, 0,3-0,4 mm. lat.; ascis cylindraceo-subclavatis, apice attenuatis, 25-35 \approx 4-5; sporidiis fusoideo vel aciculare-elongatis, simplicibus, incoloribus 6-9 \approx 1-1,5; paraphysibus filiformibus, gracillimis.

Hab. ad caules plantarum, ad ferulas Cerefolii silvestris exsiccatas, in Gallia, Fennia, Britannia. — Pro *Ps. micacei* varietate possit haberi, at cupula tenuiore, colore vix numquam fulvo siccitateque minus regulariter contracta distinctum.

12. **Pseudohelotium tenuiculum** (Karst.) Sacc. *Helotium tenuiculum* 1226

Karst. M. F. I, p. 149. — Sparsum v. subsparsum, tenue, sessile, planiusculum v. concaviusculum albidum v. lutescenti-albidum, siccum difformiter contractum, lutescens furfuraceum vel furfuraceo-tomentellum, margine fimbriatum, 0,5-1 mm. lat.; ascis fusoideo-clavatis, 70-80 \approx 8, obturaculo minutissimo, jodo obsolete cœrulecente; sporidiis distichis, fusoideo-elongatis v. fusoideo-bacillaribus, ut plurimum curvulis, incoloribus, simplicibus v. raro tenuiter 1-3-septatis, 14-20 \approx 3; paraphysibus gracillimis.

Hab. supra ferulas Cerefolii silvestris exsiccatas in Mustiala Fenniæ.

13. **Pseudohelotium millepunctatum** (Lib.) Sacc. *Moll. millepunctata* 1227

(Lib.) Sacc. Fung. Ard. p. 8, nec *Pyrenop. millepunctata* Sacc. *Peziza millepunctata* Lib. Exs. n. 128. — Gregaria, punctiformis, sessilis, globoso-urceolata, tenerrima, albo-hyalina, extus exasperata; ascis cylindraceis; sporidiis ovatis (Libert).

Hab. ad caules putres Seneciois Fuchsii in Arduennis. — Ascis et sporidiæ non vidi.

14. **Pseudohelotium Teucrii** (Fuck.) Sacc. *Niptera Teucrii* Fuck. 1228

Symb. App. I, p. 355. — Cupulis quandoque dense gregariis, $1\frac{1}{2}$ -1 mm. lat., sessilibus, aquoso-diaphanis, pallidissime flavo-albis, subochroleucis, primo subclausis, excavatis; demum planis, orbiculari-

bus, marginatis, extus margineque vix puberulis, disco concolori, siccis flavescentibus; ascis globuloso-stipitatis, cylindraceis, 8-sporis, $50 \approx 8$; sporidiis in asci superiori parte dense congestis, cylindraceis, parum curvatis, continuis, $8 \approx 1-2$; paraphysibus filiformibus, guttulatis.

Hab. in caulis emortuis Teucri Scorodoniiæ in Germania. — Sec. Rehm Ascom. Ill. n. 757, excipulum tenui-parenchymaticum, flavidum; asci 30-36 ≈ 5 ; sporidia 5-6 ≈ 1 .

15. **Pseudohelotium punctiforme** (Grev.) Sacc. *H. punctiforme* (Grev.) 1229
Phill. Disc. p. 169, *Helotium punctatum* Fr. Summa p. 356,
Sacc. F. it. tig. 1351, *Peziza punctiformis* Grev. Scot. t.
63 nec Auctorum. — Cupulis minutis, $1/4-1/5$ mm. diam., sessilibus, centro punctiformi adfixis, denique applanatis, disco albido, subinde lutescente, extus margineque crenulato albido-pilosellis; ascis cylindraceo-clavatis, deorsum longiuscule attenuatis, $40 \approx 4$, octosporis; sporidiis distichis fusoideis, $6 \approx 1\frac{1}{2}$, utrinque minute 1-guttulatis, hyalinis.

Hab. in foliis quercinis dejectis in Scotia et Padova e Conegliano Ital. bor. — *Hel. punctoideum* Karst. Fenn. I, 151, huic peraffine videtur. — Var. **robustellum** Sacc. Mich. I, p. 427, F. it. f. 1352: recedit ascomatibus majoribus, eximie margine sinuosis, ascis sporidiisque duplo majoribus (nempe primis $60 \approx 7-8$, postremis clavulato-bacillaribus $12-15 \approx 1\frac{1}{4}-1\frac{3}{4}$); excipuli contextu anguste parenchymatico, fuscescente ad marginem extusque prosenchymatico-spinuloso, dilutiore. In foliis dejectis et arescendo (ob mycelium?) aurantiaco-coloratis quercinis in silva Montello Ital. bor.

16. **Pseudohelotium punctoideum** (Karst.) Sacc. *Pezizella punctoidea* 1230
(Karst.) Rehm Ascom. n. 162, *Pseudohelotium chloroticum* Kunze:
Cfr. Grevill. V, p. 61, *Hel. punctoideum* Karst. M. F. I, p. 151.
— Gregarium, sessile vel substipitulatum; cupula subimmarginata, subnuda, convexa, sicca plana vel concaviuscula, hyalino-albida, disco siccitate testaceo, 0,4 mm. lat.; ascis e basi tenuata, flexuosa, cylindraceo-clavatis vel clavatis, $38-43 \approx 5-6,5$, apice jodo non tinctis; sporidiis distichis, fusoideo-elongatis, ut plurimum rectis, subeguttulatis, $5-8 \approx 2-3$; paraphysibus gracilescentibus.

Hab. ad folia Epilobii angustifolii putrida locis uidis in Mustiala Fenniæ et Germania. — Ascomata siccitate interdum subhemisphærico- vel subsphæroideo-contracta, leviter furfurella, sessilia vel stipite punctiformi.

17. **Pseudohelotium Pineti** (Batsch) Fuck. Symb. pag. 298, *Peziza* 1231
Batsch Elench. p. 201, f. 140, Fr. S. M. II, p. 101, *Helotium* Karst.
M. F. I, p. 142, Pat. Tab. f. 493, *Pez. abieticola* Nyl. — Sessile,
adnatum, subplanum, firmulum, puberulum, sordide album, disco
albido; ascis stipitatis, subclavatis, 8-sporis, $24 \approx 8$; sporidiis fu-
soideo-cylindraceis, rectis, $8-12 \approx 2-3$, hyalinis; paraphysibus fili-
formi-clavulatis.
- Hab.* ad acus Pinuum putrescentes in Suecia, Germania, Fen-
nia, Britannia, Belgio. — Ascoma $\frac{1}{2}-1$ mm. lat., minute urceola-
tum, mox explanatum, margine integro inflexo obtuso. Sec. Rehm
Ascom. n. 561, sporidia 18-20 \approx 1,5.
18. **Pseudohelotium dispersellum** (Karst). Sacc. *H. dispersellum* Karst. 1232
M. F. I, p. 135. — Sparsum, sessile vel subsessile, cyathoideum
aut subhemisphæricum, albidum, marginem versus furfurellum,
siccum obovoideo-contractum, fuscescens, margine connivente, fir-
mulum, 0,6 mm. lat.; ascis cylindraceo-clavatis, sessilibus, $70 \approx 6-7$;
sporidiis elongatis v. fusoideo-elongatis, vulgo rectis, $14-21 \approx 2,5-$
 $3,5$; paraphysibus filiformibus, sursum incrassatulis.
- Hab.* ad folia Caricis elongatæ emortua prope pagum
Kytö par. Tammela Fenniæ.
19. **Pseudohelotium Jerdoni** Sacc. *Helot. pruinosa* Jerdon in B. et 1233
Br. Ann. Nat. Hirt. n. 1174, t. V, f. 33 nec Wallr. — Sessile,
planum, crassiusculum, firmum, marginatum v. immarginatum, cal-
ceo-album, pruinosa; ascis tereti-clavatis; sporidiis oblongis v.
oblongo-ellipsoideis, triquadriguttulatis, demum (spurie?) bisepta-
tis, $20-22 \approx 4$; paraphysibus filiformibus.
- Hab.* in Hypoxylon fusco et Diatrypè Stigmate ad ligna
putrida in Britannia et Belgio. — Ascoma 1 mm. lat., hymenium
subinde cyanescens.
20. **Pseudohelotium versicolor** (Desm.) Sacc. *Pez. versicolor* Desm. 1234
22 Not. p. 18. — Minutissimum, sessile, albo-hyalinum, vix tactum
flavescens, fragile, glabrum, primo globosum, dein turbinato-con-
cavum, extus leviter furfuraceum, margine subdenticulatum; ascis
clavatis, 30μ . long.
- Hab.* ad folia stipitesque Aspidii Filicis-maris in Gallia.
— Ascoma $\frac{1}{2}$ mm. diam. Ad *Cyathiculam* vergit.
21. **Pseudohelotium miliare** (Wallr.) Sacc. *P. miliaris* Wallr. Fl. Crypt. 1235
n. 2605. — Ascomatibus sessilibus sparsis microscopicis pallenti-
bus, extus tenuiter pruinosis immarginatis, cum disco demum con-
vexusculo concoloribus.

Hab. ad *Peltigeræ caninæ* paginam superiorem in locis uidis Thuringiæ passim.

II. Disco flavo, flavido, stramineo-aurantio.

22. **Pseudohelotium pruinatum** (Wallr.) Sacc. *P. pruinosa* Wallr. 1236

Crypt. n. 2564, nec Jerd. — Ascomatibus sessilibus gregariis stramineis carnoso-membranaceis, extus tenuiter farinosis, primum subglobosis vertice hiantibus deinde discum pallide lutescentem recludentibus scutellaribus, demum dilatatis planiusculis, marginem flexuosum reclinantibus glabrescentibus, facile decoloribus.

Hab. ad *Solani tuberosi* caules putres agri Northusani, vere (WALLR.)

23. **Pseudohelotium eurotioides** (Karst.) Sacc. *Helotium eurotioides* 1237

Karst. M. F. I, pag. 146. — Subsparsum, sessile, hemisphæricum, siccum sphæroideo-contractum, stramineo-aureum vel sulphureum, albo-puberulum, circiter 0,2 mm. lat.; ascis cylindraceo-clavatis v. clavatis, 31-43 \times 5-6; sporidiis ovoideo-oblongatis v. aciculari-elongatis, guttulis majusculis præditis, 5-8 \times 2-2,5; paraphysibus obsoletis.

Hab. ad caules *Artemisiæ vulgaris* et *Cerefolii silvestris* aridos in Mustiala Fenniæ.

24. **Pseudohelotium pulchellum** (Quél.) Sacc. *Mollisia pulchella* Quél. 1238

11, Suppl. p. 21, t. XII, f. 10. — Ascomate globuloso, urceolato, 100-200 μ . diam. tenui, glabro, nitide aureo, aculeis tenuibus pulchre marginato; hymenio concavo, concolori; sporidiis cuneatis, 5-6 μ . longis, hyalinis.

Hab. in caulis exsiccatis *Orobi verni*, *Dentariae pinnatae* in Jura. — *Helotio punctato* et *H. eurotioidi* simile.

25. **Pseudohelotium Absinthii** (Lasch) Sacc. *Peziza Absinthii* Lasch 1239

in Klotzsch H. M. n. 1317 et in Bot. Zeit. 1850, pag. 438. — Ex flavido fuscescens, sessile, subrotundum, ruguloso-fibrillosum, margine infexo albido-simbriato, disco dilutiore flavido.

Hab. in caulis *Artemisiæ Absinthii*, Driesen Germaniæ.

26. **Pseudohelotium separabile** (Karst.) Sacc. *H. separabile* Karst. 1240

M. F. I, p. 118. — Confertum vel gregarium, sessile, primitus leviter puberulum, dein glabrescens, ex subsphæroideo appланatum, album vel albidum, disco flavescente, margine integrum, subinde flexuosum, 0,5-1 mm. lat.; ascis cylindraceo-clavatis, apice jodo non tinctis, 60-80 \times 5-6; sporidiis, fusoideo-elongatis, 2-guttulatis

vel spurie tenuiter uniseptatis, rectis vel curvulis 8-15 × 2-3; paraphysibus gracilibus.

Hab. ad caules Rubi idæi exsiccatos prope Mustiala Fenniæ.

27. **Pseudohelotium gemmiferum** (Wallr.) Sacc. *P. gemmifera* Wallr. 1241
Fl. Crypt. n. 2596. — Ascomatibus gregariis parvis subcarnosis
flavicantibus, extus gemmulis pellucidis plurimis splendentibus ob-
sessis, primum obconicis clausis, dein in discum marginibus inflexis
cinctum saturatiorem reclusis sensimque in basim stipitiformem de-
sinentibus.

Hab. ad caudices quercinos in Hercynia inferiore passim. —
Habitus externus *P. citrinæ*; nec quoque *P. pallida* Schum. Saell.
II, 428, dubia Friesio, dissimilis esse videtur.

28. **Pseudohelotium sacchariferum** (Berk.) Sacc. *Peziza saccharifera* 1242
Berk. North Am. Fungi n. 738, Cooke Disc. U. S. I, p. 196. —
Mollis, gregaria, pallide aurantiaca, irregularis, extus saccharina;
margin'e tumidula; disco concavo.

Hab. in truncis Liquidambaris, Alabama in America boreali
(PETERS).

29. **Pseudohelotium fibrisedum** (B. et C.) Sacc. *Peziza fibriseda* B. 1243
et C. North Am. Fungi n. 737, Cooke Disc. U. S. I, pag. 296. —
Irregularis, aurantiaca, extus saccharino-granulosa; margin'e fracto,
laciniato; disco concavo.

Hab. ad truncos Ulmi americanæ in montanis Virginiae
Amer. bor.

30. **Pseudohelotium minutum** (Pass.) Sacc. *Peziza minuta* Pass. Fung. 1244
Abiss. n. 17, t. IV, f. 7, Fl. Bog. p. 137. — Sparsum v. subgre-
garium, pusillum, ceraceo-carnosum, flavescens vel subcarneum,
subhyalinum, disco concavo, margine furfuraceo, siccitate versi-
formi contracto; ascis cylindricis, basi attenuata incurvis, apice
obtuso subtruncato, 8-sporis; sporidiis minutis obscure distichis
fusiformibus, rectis, hyalinis, apicibus acutis, 8 × 1-1,5 (ex iconæ);
paraphysibus tenuibus parcis.

Hab. ad fragmenta vetusta corticis, Keren Abyssiniae.

31. **Pseudohelotium nipteroides** (Speg.) Sacc. *Helotium? nipteroides* 1245
Speg. Fung. Fueg. n. 335. — Cupulis densiusculis sparsis, udis
applanato-concavis 1-2 mm. diam., siccis subcontractulis angulosis
sessilibus crassiusculis rigidulis, extus pulchre griseo-fuscis minu-
tissime pruinosis, papillis pusillis 1-cellularibus obtusis flexuosis
10-25 × 5 vestitis, contextu subparenchymatico carnoso subindistin-
cto, disco plano ceraceo flavescente, margine acutiusculo integro

v. repandulo; ascis cylindraceis, antice acutato-rotundatis v. subtruncatis, postice longiuscule attenuato-pedicellatis, 80-90 \approx 7-8; paraphysibus paucis filiformibus ovallatis, octosporis; sporidiis recte v. oblique mono- v. distichis ellipticis v. elliptico-ovoideis utrinque obtusis 12-13 \approx 3-3,5, hyalinis, continuis (an per ætatem medio 1-septatis?).

Hab. in truncis cariosis putrescentibus fagineis? in silvis in Staten island Fuegiæ.

32. **Pseudohelotium succineum** (Quél.) Sacc. *Mollisia succinea* Quél. 1246 X Suppl. p. 13, t. IX, f. 11, Grevillea 1879, p. 38. — Ascomatibus turbinatis, obconicis, 1 mm. diam., breviter stipitatis, subtilibus, in sicco subscariosis, diaphanis, flavo-albis, margine pruinoso albescente, hymenio concolori; sporidiis bacillaribus, 8 μ . longis.

Hab. in lignis exsiccatis ad ripas rivorum, Jura.

33. **Pseudohelotium stramineum** (B. et Br.) Sacc. *Peziza* B. et Br. 1247 Ann. N. H. n. 571. — Confertum vel sparsum, sessile hemisphaericum dein expansum, pallide flavum, granulis crystallinis conspersum; disco rubro-flavo v. carneo; margine inflexo; ascis teretico-clavatis; sporidiis fusoides rectis vel curvulis, 3-4-guttatis, 7-10 \approx 1,5; paraphysibus tenuissimis.

Hab. in vaginis foliisque graminum in Britannia. — Ascoma 1 mm. lat.

34. **Pseudohelotium epicalamium** (Fuck) Sacc. *Pez. epicalamia* Symb. 1248 mycol. App. II, p. 61. — Cupulis carnosis, sparsis seu dense gregariis, subsessilibus, primo globosis, clausis, demum explanatis, vix $1\frac{1}{2}$ mm. latis, orbicularibus, integris, extus margineque subtilissime lacerato demum involuto, striato, albicante-flavescente, furfuraceis, disco concavo seu explanato, luteo-ochraceo; ascis elongato-clavatis, sessilibus, 8-sporis, 52 \approx 6; sporidiis distichis, fusiformibus, continuis, rectis, 2-3-guttulatis, 9 \approx $2\frac{1}{2}$, hyalinis.

Hab. in culmis Tritici repentis, Iudensand prope Estrich Rhenogoviæ.

35. **Pseudohelotium arundinellum** (S. et Sp.) Sacc. *Helotium arundinellum* Sacc. et Sp. Mich. I, p. 426, Sacc. F. it. f. 1353. — Ascomatibus gregariis minutissimis, sessilibus, concaviusculis, $1\frac{1}{6}$ mm. diam., pallidis, extus tomentello-furfuraceis, disco sordide luteolo; ascis cylindraceo-clavatis, 40 \approx 5-6, breve crassiuscule stipitatis, apice obtusiusculis; paraphysibus bacillaribus, guttulatis, octosporis; sporidiis distichis v. oblique monostichis, cylindraceo-oblongis utrinque rotundatis, curvulis, 8-10 \approx $1\frac{1}{2}$ -2, 2-guttulatis, hyalinis.

Hab. in vaginis Arundinis Donacis putrescentibus Conegliano Italiae bor. — *Pseud. micaceo* (Pers.), *P. microspi* Karst.

H. parvula (De Not.) etc. affine sed notis nonnullis satis diversum.

36. **Pseudohelotium costatum** (Boud.) Sacc. *Helotium costatum* Boud. 1250

in B. S. B. Fr. 1881, p. 95, t. III, f. 6. — Ascomate perexiguo, griseo-albido, sessili, urceolato, marginato, disco griseo-flavido, vix $\frac{1}{3}$ mm. diam., extus præcipue inferius pubescente, pilis albidis, septatis in costulas conniventibus; ascis perexiguis, 22-26 \approx 5-7, cylindraceis, apice obtusis; sporidiis fusoideo-elongatis, 8-9 \approx 1-1,5 pluriguttulatis, hyalinis; paraphysibus bacillaribus.

Hab. in calamis putridis Junc acutiflori, Montmorency.

37. **Pseudohelotium elaphoides** Sacc. *Mollisia elaphines* Quél. Champ. 1251

Norm. pag. 32 nec Berk. et Br. — Tenue, discoideum, dein expansum, undulatum, fulvum, margine pubescenti pruinoso, hymenio plano, flavido, dein obscuriore; sporidiis ellipsoideo-oblongis, 4 μ . longis.

Hab. in caulis sicca Equiseti, Jura Galliae. — Sec. Quél.
l. c. ascoma 0,1 mm. lat. Videtur diversum ab homonymo Berk.

38. **Pseudohelotium strumosum** (E. et E.) Sacc. *Helotium strumosum* E. et E. Journ. Mycol. 1888, p. 56. — Ascomatibus gregariis, sessilibus, læte citrinis, initio clauso-subsphæricis, dein aperitis, cupulatis, demum expanso-subplanis, $\frac{1}{6}$ - $\frac{1}{3}$ mm. diam., extus tomentoso-putrescentibus basique hyphulis albidis, affixis; ascis clavatis, 70-80 \approx 10-12; sporidiis distichis, ellipsoideis vel ovato-ellipsoideis, granuloso-multinucleatis, dein 2-3-nucleatis simulateque uniseptatis, 10-12 \approx 3-4,5.

Hab. in Dichæna strumosa Newfield N. J. Amer. bor. — Substantia mollis, subcarnosa; margo acutus.

39. **Pseudohelotium puberulum** (Lasch) Fuck. Symb. p. 298, *Peziza puberula* Lasch in Kl. Hb. myc. 1529. — Ceraceum, minutum, sessile v. substipitatum, luteum vel albidum, cupula ex hemisphærico planiuscula, extus puberula, disco demum convexo; ascis oblongis, curvato-stipitatis, 8-sporis, 64 \approx 8; sporidiis in ascii parte superiore inordinatis, oblongis, utrinque attenuatis, subcurvatis, hyalinis, 10 \approx 3-4.

Hab. in foliis putridis arborum frondosarum in Rhenogovia.

III. Disco melleo, virente, olivaceo.

40. **Pseudohelotium pomicolor** (B. et R.) Sacc. *Peziza pomicolor* B. et R. North Am. Fungi n. 735, Grev. t. 51, f. 260. — Sparsum,

subhemisphaericum, sero apertum, extus pomicolor furfuraceum; disco olivaceo; ascis cylindraceis filiformi-paraphysatis; sporidiis

Hab. in cortice *Taxodii* distichi in Carolina inf. (RAVENEL). — Ascoma extus colore viridi pomorum tinctum.

41. **Pseudohelotium melichroum** (Cooke) Sacc. *Pez. melichroa* Cooke 1255
Grev. VII, pag. 47. — Sparsum, sessile; cupulis minimis, vix $\frac{1}{2}$ mm., mellicoloribus, demum applanatis, extus saccharino-granulatis; margine leniter elevato; ascis cylindraceis; sporidiis subellipticis 5 μ . longis; paraphysibus filiformibus.

Hab. ad cortices, Alabama Amer. bor. (PETERS).

42. **Pseudohelotium Mali** (Rehm) Sacc. *Pezizella Mali* Rehm Ascom. 1256
n. 460, *Mollisia* Phill. Disc. p. 194. — Gregarium, minimum, primitus subglobosum, dein urceolatum, denique plus minusve exsum, siccum margine involuto, dilutiore scabro, disco viridulo-flavo, humidum orbiculare, margine crasso, disco plano, c. 0,3-5 mm. diam., contextu prosenchymatico tenerrimo, fere hyalino; ascis clavatis, subcurvatis, 4-8-sporis, 40 \approx 8; sporidiis oblongis, obtusis, hyalinis, plerumque subcurvatis, distichis, 9 \approx 3; paraphysibus filiformibus, superne sensim clavulatis.

Hab. in superficie corticis interioris distantis Piri Mali in Franconia bavarica et in cortice Ilicis, Shrewsbury Britanniæ. — Ascoma sec. Phill. pruinatum, disco melleo.

43. **Pseudohelotium elaphines** (B. et Br.) Sacc. *Mollisia elaphines* 1257
B. et Br. Ann. Nat. Hist. n. 1325, t. 19, f. 18, Phill. Disc. p. 179.
— Ascomate subgloboso, pallide cervino, margine pallidiori, granulis saccharinis consperso; disco aquose albido; ascis 45 μ . longis; sporidiis fusoideis, levibus, hyalinis, monostichis, 8,5-10 μ . longis.

Hab. in ligno putri prope Shrewsbury Britanniæ.

44. **Pseudohelotium alligatum** (B. et Br.) Sacc. *P. alligata* B. et Br. 1258
Fung. of Ceyl. n. 954. — Sparsum v. congestum, subtus albo-pulverulentum, flexuosum, hymenio melleo; paraphysibus apice curvatis; ascis clavatis; sporidiis uniseriatis ellipticis 15 μ . long.

Hab. in foliis emortuis, quibus arcte adpressum, Peradenya Ceylon.

45. **Pseudohelotium incrustatum** (Ellis) Sacc. *Mollisia incrustata* Ellis 1259
Am. Nat. 1883, p. 192. — Gregarium, rarius subconfluens, sessile vel in basim stipitiformem contractum, disco immarginato convexo vel concaviusculo in siccis, melleum, pruinatum, crystallis minutissimis saccharinis quasi conspersum, $\frac{1}{6}$ mm. diam.; ascis clavatis te-

nuiter stipitatis, 30-35 = 4; sporidiis partim distichis, ovato-oblongis, 5-6 = 1, hyalinis.

Hab. in Polyporo quodam resupinato ad truncum Juniperi virginianæ dejectum, New Jersey Amer. bor.

IV. Disco rubente, testaceo, fusco, lurido, livido.

46. **Pseudohelotium micaceum** (Pers.) Sacc. *Hel. micaceum* (Pers.) 1260

Karst. M. F. I, p. 149, *Peziza micacea* Pers. Myc. eur. I, p. 268, Fr. Syst. Myc. II, p. 105, *Pez. melleopallens* Karst. Monogr. Pez. p. 190. — Gregarium vel sparsum, sessile, e sphæroideo-hemisphæricum, marginem versus farinoso-subtomentellum vel furfuraceum, pallidum vel pallide helvolum vel subfulvum, subnitens 0,3-0,4 mm. lat.; ascis cylindraceo-clavatis 30-38 = 4-5; sporidiis fusoideo-vel aciculari-elongatis, 5-11 = 1-2; paraphysibus filiformibus, sat numerosis, 1 μ . cr.

Hab: ad caules putrescentes Cardui nutantis, Ligustici Levistici et Artemisiæ vulgaris in Germania, Gallia, Fennia.

47. **Pseudohelotium incarnatum** (Quél.) Sacc. *Mollisia incarnatina* 1261

Quél. XIII, Suppl. p. 9, t. VIII, f. 22. — Ascomatibus hemisphæricis, 200-300 μ . diam., dein concavis, ceraceis, margine puberulis, pallide roseo-carneis, hymenio roseo; sporidiis navicularibus 10 μ . longis, subfusiformibus, 2-guttulatis, hyalinis.

Hab. in caulis plantarum majorum e. g. Cirsii, prope Paris et Jura Galliæ (BOUDIER). — A *Ps. roseolo* fabrica et minutie differt.

48. **Pseudohelotium Fairmani** (E. et E.) Sacc. *Mollisia Fairmani* E. 1262

et E. Journ. Myc. 88, p. 56. — Sessile, centro affixum, gregarium concavum, margine subincervo, dein expansum, planum vel convexulum, $1\frac{1}{3}$ - $\frac{3}{4}$ mm. diam., disco livido, margine pallidiore extusque pilis glandularibus brevibus, hyalinis, obtusis, basi longioribus, demum obscuris obsessum; ascis tereti-clavatis sessilibus, paraphysatis, 35-40 = 5-6; sporidiis subdistichis, ovato-ellipsoideis, 3,5-5 = 2-2,5.

Hab. in cortice interiore dejecto Lyndonville N. Y. Amer. bor.
— In sicco fere clausum.

49. **Pseudohelotium xylitum** (Karst.) Sacc. *Helotium xylitum* Karst. 1263

M. F. I, p. 144. — Gregarium, sessile, primo sphæroideum, dein planiusculum vel concaviusculum, pallide helvolum vel testaceum,

albido-furfuraceum, siccum hemisphærico- v. difforini-contractum, rosello-helvolum, 0,4-1,2 mm. lat.; ascis fusoideo-clavatis, 37-45 \approx 4,5-5,5, apice jodo non tinctis; sporidiis distichis v. oblique monostichis, aciculari- vel fusoideo-elongatis v. oblongatis, rectis v. subrectis, eguttulatis vel 2-guttatis, 6-10 \approx 2,5-1,5; paraphysibus gracilescentibus sursum leviter incrassatis, 2 μ . cr.

Hab. supra lignum induratum vetustum Alni, Betulæ et Populi in Fennia meridionali.

50. **Pseudohelotium granulosellum** (Karst.) Sacc. *Helotium granulosellum* Karst. M. F. I, p. 142. — Gregarium v. subsparsum, sessile, tenue planiusculum, hyalino-albidum, siccum subsphæroideo-hemisphæricum, albidum v. stramineo-albidum, disco subtestaceum, 0,4-0,6 mm. lat.; ascis cylindraceo-clavatis, 38-54 \approx 5-6, obturaculo minutissimo jodo cœrulecente; sporidiis oblique monostichis, fusoideo- vel aciculari-elongatis aut ovoideo-oblongatis, eguttulatis, rectis, 5-9,5 \approx 2-2,5; paraphysibus sat raris, gracilibus sursum leviter incrassatis.

Hab. in ligno betulino vetusto ad Mustiala Fenniæ.

51. **Pseudohelotium Aspidiorum** (Rehm) Sacc. *Helotium Aspidiorum* 1265 Rehm Ascom. n. 563. — Minutissimum, gregarium, c. 0,1 mm. altum, turbinato-globosum, fusco-atrum, sæpe subpulverulentum, disco plus minusve gracili et dilutiore, prosenchymatico-contextum, fibris c. 5 μ . crassis, obtusis, sæpe distantibus, fuscis; ascis teneris, clavatis, apice subincrassatis, 36-45 \approx 5-6, 8-sporis, apice jodo cœrulecente; sporidiis elongato-ellipticis, obtusis, rectis, binucleatis, hyalinis, distichis, 12-15 \approx 2,5; paraphysibus teneris, filiformibus; jodo cœrulecente apice ascorum.

Hab. ad putrescentem frondem Athyrii Filicis-feminæ, Rindalphorn alpium Allgoviæ c. 1500, mm. alt. (BRITZELMAYR).

52. **Pseudohelotium hymeniophilum** (Karst.) Sacc. *H. hymeniophilum* 1266 Karst. M. F. I, p. 143. — Gregarium, sessile, primo sphæroideum, dein planiusculum, siccum hemisphærico-applanatum, sæpiissime difforme, furfurellum, cinerescenti-albidum vel pallens, disco sublurido, circiter 0,4 mm. lat.; ascis cylindraceo-clavatis, 30-40 \approx 4,5-5,5, apice jodo non tinctis; sporidiis elongatis vel oblongatis 4-6 \approx 1,5.

Hab. supra hymenium Polypororum resupinatorum in Fennia et Lapponia orientali. — Substratum colore roseo-sanguineo late pertingens, qua nota facillime distinctum.

B. Mollisiella Phill. emend.: sporidiis globulosis.

53. **Pseudohelotium ilicinolum** (B. et Br.) Sacc. *Peziza* B. et Br. 1267
Ann. N. H. n. 958, tab. 16, f. 17, *Mollisia ilicincola* Phill. Disc.
p. 193. — Fasciculatum, furfuraceum, hemisphaericum dein expan-
sum, extus sordide album; disco concavo pallide brunneo vel pur-
pureo v. roseo-cinereo; margine erecto v. incurvo; ascis clavatis;
sporidiis globosis, 1-guttatis 5 μ . diam.; paraphysibus filiformi-
clavulatis.

Hab. in ramis Ilicis sæpe in thallo Myriangii, Cheiselhurst
et Forres Britanniæ.

54. **Pseudohelotium apicale** (B. et Br.) Sacc. *Pez. apicalis* B. et Br. 1268
Fung. of Ceyl. n. 951. — Pallide aurantiacum, congestum, subglo-
bosum, margine incurvo, extus pulverulentum; ascis clavatis; spori-
diis, minimis subglobosis, 3-4 μ . d.

Hab. ad caules Macromitrii sulcati in ins. Ceylon. — Per-
exiguum, nonnisi plura specimina congesta oculo inermi visibilia:
excipulum e cellulis longitud. seriatis constans.

55. **Pseudohelotium hydnicolum** (B. et Br.) Sacc. *Pez. hydnica* B. 1269
et Br. Ann. N. H. n. 1327, t. XX, f. 20, *Mollisia hydnica* Phill.
Disc. p. 194. — Orbiculare, dein irregulare, planum, atro-virens;
ascis cylindraceis; sporidiis subglobosis 1-seriatis, 10 \approx 7,5-9; para-
physibus filiformibus ramulosis.

Hab. in Hydno ochraceo in Britannia. — Conidia adsunt
obtusa 2-4 μ . long. Ascoma a cl. B. et Br. furfuraceum non de-
scribitur, sed subg. *Mollisiella*, quo a cl. Phillips includitur et
ascomate furfuraceo-pulvereo dignoscitur.

CYATHICULA De Not. Prof. Disc. in Comm. I, p. 381, ex p. (1864)
Sacc. Consp. Disc. p. 5 ex p. (ut subg.) [Etym. *Cyathus*], *Trichoscy-
pha* et *Peristomialis* Phill. Disc. p. 126 et 201 (ut subg.), *Calycelle*
et *Cistellæ* spec. Quél. — Ascoma ceráceo-membranaceum, tenué,
initio suburceolatum, mox apertum, stipitatum vel sessile, margine
distincte dentatum. Ascí fusoides v. teretiusculi, 8-spori, filiformi-
paraphysati. Sporidia oblonga v. teretiuscula, continua, hyalina.

I. *Eu-Cyathicula*. Stipitatæ.

1. **Cyathicula coronata** (Bull.) De Not. Discom. p. 381, *P. coronata* 1270
Bull. Champ. p. 251, t. 416, f. 4, *P. armata* Roth. cat. I, p. 240,
P. radiata Pers. Myc. Eur. p. 237, Nees Syst. f. 293, *P. subu-*

lata Schum. Saell. 2, p. 418, Fl. Dan. t. 1380, f. 1, Pers. l. c. p. 286, *Helotium coronatum* Karst. Pat. Tab. 593, *P. denticulata* Fl. Dan. t. 1016, fig. 1 (forma lignatilis). — Stipitata, pallens, cupula concava, margine dentibus setaceis coronata; ascis oblongis, stipitatis, 8-sporis, 88 \approx 8; sporidiis distichis, oblongo-fusiformibus, curvatis, hyalinis, 16-18 \approx 3-4.

Hab. ad caules dejectos plantarum Urticæ Sed i, Scrophulariæ etc. in Suecia, Gallia, Britannia, Italia, Belgio, Amer. bor. — Sparsa, subgregaria, levis, glabra, albida, sicca pallescenti-fusca. Stipes 2-5 mm. longus, subincurvus; cupula 2 mm. lata, dentibus primo conniventibus, dein erectis.

2. **Cyathicula inflexa** (Bolt.) Sacc. *Peziza inflexa* Bolt. t. 106, Sow. 1271 t. 306, Fr. Syst. II, 120, *P. Coronilla* Pers. M. E. I, p. 287. — Ascomatibus majusculis stipitatis albidis, cupulæ subhemisphæricæ margine dentibus triangularibus brevibus cincto.

Hab. in caulibus Urticæ etc. in Britannia, Belgio. — A præcedente, cum qua confusa fuit, differre videtur statura majori, colore dilutiori dentiumque forma. Ita Fr. l. c. Sec. Phill. Disc. p. 127 est var. *C. coronatæ*.

3. **Cyathicula alba** (Pat.) Sacc. *Calycella alba* Pat. Tab. anal. Fung. 1272 II, p. 37, f. 594. — Sparsa, plana v. leniter convexa, stipitata, glabra, omnino alba, margine in dentes acutos erectos diviso; stipite basim versus attenuato; ascis clavatis; paraphysibus linearibus, guttulatis; sporidiis fusoideis rectis v. curvulis, guttulatis, 14-16 \approx 3.

Hab. in caulibus herbaceis putridis in Gallia. — Stipes $\frac{3}{4}$ mm. longus.

4. **Cyathicula petiolorum** (Desm.) Sacc. *Helotium petiolorum* (Desm.) 1273 De Not. Comm. Critt. I, p. 378, *Peziza* Desm. IX Not. p. 6. — Majuscula, stipitata, sæpe solitaria, cupula fulva, concava, dein plana, margine dentibus subtriangularibus cincta; stipite plus minusve longo, basi brunneo; ascis clavatis, 100 μ . long.; sporidiis oblongis, curvatis, 2-guttulatis, 10 μ . longis.

Hab. in petiolis foliorum emortuorum Fagi etc. in Gallia. — Ascoma 2 mm. lat.

5. **Cyathicula clavuligera** Speg. Fung. Fueg. n. 329. — Sparsa, amphigena, minutissima, discoideo-applanata, 600-800 μ . diam., rufescens, inferne subturbinata, glabra pulchre rubescenti-ochracea, margine acutiuscula hinc inde denticulato-subfimbriata, stipite breviusculo ac crassiusculo, 700-1000 \approx 130-160, inferne non vel vix incrassato sæpius nigricante rigidulo; ascis cylindraceis v. cylindra-

ceo-subfusoideis, antice truncatis vel truncato-rotundatis, postice attenuato-pedicellatis, 75-80 \approx 7-9, octosporis; paraphysibus filiformibus non vel vix longioribus continua apice clavulatis, ascorum obturaculo minutissimo iodi ope cœrulecente; sporidiis oblique monostichis, ellipticis v. ovoideis, utrinque plus minusve obtusatis 9-10 \approx 4 non v. nubiloso-farctis, hyalinis.

Hab. ad folia dejecta putrescentia Fagi betuloidis in silvis in Staten island Fuegiæ.

6. **Cyathicula incarnata** (Quél.) Sacc. *Phialea incarnata* Quél. V 1275 Suppl. p. 329, t. VI, fig. 9. — Ascomatibus hemisphæricis, 2 mm. diam., coriacellis, albis, margine fibrilloso, denticulato-fimbriato; stipite 1 mm. alto, glabro albo; hymenio roseo-pallido; sporidiis 20-25 \approx 4, oblongo-bacillaribus, (spurie?) 5-septatis.

Hab. ad culmos in silvis humidis Juræ. — Si sporidia vere septata, species ad *Belonidium* accederet, sed habitus omnino *Cyathiculæ*.

7. **Cyathicula Queletii** Sacc. *Phialea ciliata* Quél. Esp. nouv. Champ. 1276 Fr. 1878, p. 291 et X Suppl. p. 13, t. IX, f. 6. — Stipite filiformi, 1-2 cm. alto, glabro, brunneo; ascomatibus 2-4 mm. diam., griseis, ciliatis, h. e. denticulis subulatis, brunneis præditis; sporidiis fusiformibus, 15 μ . longis, sæpe curvis.

Hab. in caulibus herbarum, Jura.

8. **Cyathicula denigrans** (Fuck.) Sacc. *Peziza denigrans* Fuck. Symb. 1277 mycol. p. 309. — Cupulis stipitatis, 4-6 mm. altis, hemisphæricis, in stipitem contractis, fuscis 2 mm. diam., margine fasciculato-ciliatis, fasciculis triangularibus, fuscis, stipite in foliorum petiolo nigrificato insidente, basi nigro, disco pallide fusco; ascis elongatis, stipitatis, 8-sporis, 70 \approx 8 (p. sporif.); sporidiis monostichis, oblongis, utrinque obtusis, continua, rectis, 8 \approx 2-3, hyalinis.

Hab. in nervis foliorum dejectorum Fagi silvaticæ in silva Æstrich Rhenogoviæ. — Status conidicus, sec. Fuckel, est *Sphaeridium flavo-virens* Fuck.

II. *Peristomialis* Phill. (*Cistellæ* spec. Quél.). Sessiles.

9. **Cyathicula ciliata** (Cr.) Sacc. *Helotium ciliatum* Cr. Finist. p. 47, 1278 t. ult. f. 6. — Ascomate flavo-aurantio pallido, sessili, subcylindrico, $1\frac{1}{5}$ mm. lat., margine ciliato-dentato; hymenio aurantio-flavo; ascis subclavatis, crassis octosporis; sporidiis fusoideo-elongatis.

Hab. in caulibus emortuis Urticæ, Finistère Galliæ. — Celulæ excipuli prosenchymatico-anastomosantes.

10. **Cyathicula aquilina** (Rehm) Sacc. *Coronellaria aquilina* Rehm 1279
in Hedwigia 1885, p. 231. — Ascomatibus sparsis in pagina inferiore frondis, sessilibus, siccis fere globosis, extus flavidulis et dilutius marginatis, humidis concavis v. subplanis, extus et margine subfimbriato dilute flavidulis, contextu parenchymatico flavido, tenuerrimo, fimbriis marginalibus hyalinis discoque hyalino, 2-3 mm. diam.; ascis clavatis, apice obtuse acutatis, latis basi sessilibus, 8-sporis, $33 \approx 6-7$; sporidiis oblongis haud acutatis, rectis, 1-cellularibus, hyalinis, distichis, $9 \approx 2$; paraphysibus paucis, teneris, filiformibus, haud clavatis, hyalinis.

Hab. ad frondem siccam Pteridis aquilinæ, Gruneswald prope Berlin, in Germania (P. Sydow).

11. **Cyathicula Marchantiæ** (Somm.) Sacc. *P. Marchantiæ* Sommerf. 1280
Lapp. p. 295, Fr. Elench. II, p. 12. — Sessilis, infundibuliformis, glabra, albida, margine ciliata, ciliis erectis distinctis.

Hab. in Marchantia polymorpha marcescente ad rivulos subalpinos Nordlandiæ. — Magnitudine seminis sinapeos et minor; ob defectum stipitis et minutiem cum *C. coronata* confundi non potest.

12. **Cyathicula pseudo-ciliata** (Phillips) Sacc. *Helotium* Phill. Grev. 1281
XV, p. 17. — Sparsa; ascomatibus 1-2 mm. lat., sessilibus v. substipitatis, concavis carneo-rufis, margine squamulis setiformibus fimbriato; ascis cylindraceo-clavatis, octosporis; sporidiis oblongo-fusoides v. clavatis, subinde curvulis, $20-25 \approx 5-6$; paraphysibus filiformibus, gracilibus.

Hab. in ligno putri New Zealand (COLENZO).

13. **Cyathicula crystallina** (Quél.) Sacc. *Helotium crystallinum* Quél. 1282
V Suppl. pag. 329, t. VI, fig. 12. — Ascomatibus semper apertis, 300 μ . diam., concavis, tenuissimis, eleganter dentato-crenulatis, glabris, diaphanis, nitentibus; sporidiis teretiusculis curvulis 12 μ . longis.

Hab. inter strata librosa Salicis et Quercus prope Paris et Jura Galliæ.

14. **Cyathicula dentata** (Pers.) Sacc. *Pez. dentata* Pers. ic. et descr. 1283
p. 5, t. 1, f. 6, 7, Fries. Syst. Myc. vol. II, p. 147, *Calloria* Phill.
Brit. Disc. p. 332. — Sessilis, minutissima, albido-pallens subfarinacea, margine dentato; ascis clavatis; sporidiis biseriatis, fusoides, $5 \approx 1-2$.

Hab. in ligno carioso, materia virescente saepius obducto, in Suecia, Germania, Britannia. — Ab initio subglobosa, dein hemi-

sphaerica, tenera, 300-400 μ . diam., margine dentato-inciso. — Var. **straminea**, *Nipt. dentata* var. Fuck. Symb. App. I, p. 335: ascis stipitatis, primo oblongis, demum magis elongatis, 8-sporis, 56-80 \approx 9-10; sporidiis primo oblique monostichis, dein in ascis superiori parte stipatis, subclavatis, biguttulatis, continuis, hyalinis, 8 \approx 3; paraphysibus filiformibus. — In ligno putri Juglandis pr. Johannisberg Rhenogoviæ.

15. Cyathicula peristomialis (B. et Br.) Sacc. *Peziza peristomialis* 1284

B. et Br. Ann. Nat. Hist. n. 1169, t. V, f. 32, Phill. Disc. t. VI, fig. 37. — Minuta, sessilis, cylindracea, pallida, margine dentibus longis triangularibus, albis ornato, disco subplano; ascis lanceolatis; sporidiis distichis, fusoideis, multiguttulatis, 20-27 \approx 4-5.

Hab. in cortice putri Ilicis, Penzance Britanniæ. — Ascoma 300-400 μ . alt., 200 μ . lat.

16. Cyathicula lepidula March. Fung. coproph. Belg. p. 38. — Asco- 1285

matibus subgregariis, patellæformibus, demum valde expansis, glabra, albis, 250-450 μ . diam., margine tenui, laciniato, laciniis irregularibus, elongatis, approximatis, 20 μ . et ultra longit, disco concavo ad marginem ascis paucis leviter papillato; paraphysibus tenuibus, hyalinis, cyntinuis, apice non incrassatis, ascos æquantibus v. eos vix superantibus; ascis fusoideo-clavatis, sæpe curvatis, breviter stipitatis, 55-65 \approx 7,5-8,5 μ .; sporidiis subdistichis, oblongis, utrinque rotundatis, primo 2 guttulis præditis, dein spurie vel distincte 1-septatis, 10,5-11 \approx 3,3-3,5 μ .

Hab. in fimo putrido damarum Tervueren Belgii. — An hujus generis?

17. Cyathicula pusilla (Oed.) Sacc. *Helv. pusilla* Oed. Fl. Dan. t. 779, 1286
f. 1 (nec 3), *Pez. Oederi* Pers. Syn. p. 668, *P. pusilla* Fr. Syst. Myc. II, p. 148. — Sessilis, minuta, glabra, subtus albida, disco cœrulecente, margine crenato.

Hab. ad truncos putridos in Dania.

ALLOPHYLARIA Karst. Monogr. Pez. Fenn. pag. 103 [1869] (Etym. *allophylos* heterogeneus), *Helotidium* Sacc. Consp. gen. Disc. p. 4. — Ascoma clavato-turbinatum vel obconice stipitatum, glabrum, subdiaphanum, lutescens vel ochraceo-pallidum, disco concavo-truncato. Excipulum ceraceum. Ascis clavati, paraphysati, octospori. Sporidia oblonga, continua, hyalina. — Ab *Helotio* differt ascomatibus turbinatis, apice subtruncatis, subdiaphanis. *)

*) Post *Allophylariam* cl. Karst. collocat suam *Clibanitem*, quæ vero, teste

1. **Allophylaria nana** Sacc. *Helotium nanum* Sacc. Fung. Ven. Ser. 1287 IV, n. 91, F. ital. f. 115. — Cupulis gregariis cum stipite crassiusculo obconicis, ceraceis, subdiaphanis, flavidis, in sicco extus longitudinaliter striolatis, glaberrimis, $\frac{1}{2}$ mm. alt., disco aperto concaviusculo, margine acutiusculo crenulato, excipulo e fibris longis, fusoideis, parallelis efformato; ascis clavatis longiuscule stipitatis, 100 μ . long., p. s. $50 \div 9$, paraphysibus copiosis filiformibus, apice clavulatis obvallatis, 8-sporis; sporidiis distichis cylindraceo-fusoideis, $12 \div 3-3\frac{1}{2}$, rectis curvulisve, continuis, hyalinis.
- Hab.* in caule Galeobdolonis, Senecionis nemorensis et Dulcamaræ in silva Montello (Treviso) Ital. bor.
2. **Allophylaria byssacea** (Karst.) Rev. mon. p. 131, *Pezicula* Karst. 1288 M. F. I, p. 165. — Sparsa v. subgregaria, primitus obconice elongata, demum deorsum substipitato-constricta, hyalina, sicca subtestaceo-pallida vel concolor, disco concaviusculo, circiter 0,3 mm. alt., 0,1-0,2 mm. lat.; ascis clavatis, $85 \div 9-10$, obturaculo minutissimo ope jodi leviter violaceo-fuscescente; sporidiis distichis, fusoideo-elongatis, guttulis 2-4 majusculis præditis v. spurie tenuiter septatis, rectis v. leniter curvulis, $15-20 \div 3-4,5$; paraphysibus sat numerosis, gracilibus, $0,5-1 \mu$. cr.
- Hab.* supra folia Caricis putrescentia in Tammela Fenniæ.
3. **Allophylaria phyllophila** (Karst.) Rev. Monogr. p. 131, *Pezicula* 1289 *phyllophila* Karst. M. F. I, p. 167, vix *Peziza phyllophila* Desm. (Cfr. *Phialea phyllophila*). — Gregaria v. subsparsa, breviter stipitulata vel subsessilis, albido-hyalina, sicca lutescenti-albida vel albida, disco plano-concavo, demum convexiusculo, 0,3 mm. lat.; ascis clavatis, $105 \div 8-10$, obturaculo minuto, ope jodi leviter violaceo-fuscescente; sporidiis distichis, oblongatis, rectis v. lenissime curvulis, guttulas 2 sat minutis foventibus, $10-16 \div 3,5-5$ plerumque 4 μ . cr., chlorino-hyalinis; paraphysibus numerosis, 1 μ . cr.

Cooke Grev. XIII, pag. 8, inter Pyrenomycetes juxta *Hypocream* aptius inserenda sit. Donec fungi indoles melius innotescat, ejusdem diagnosim refiero. — **Clibanites** Karst. M. F. I, p. 14. Ascoma sessile, glabrum, irregulare et difforme, ceraceum. Ascii fusoideo-cylindracei, membrana tenuissima instructi. Sporidia cylindracea, hyalina, guttulata vel spurie tenuiter septata. Paraphyses parum notabiles. — **Clibanites paradoxa** Karst. l. c. Gregaria v. confluens, sessilis, tuberculosa, fulvo- vel rufo-testacea, 0,2-0,6 mm. lat.; ascis teretifusoideis, $30-36 \div 4$; sporidiis subdistichis, bacillaribus, 4-guttulatis v. spurie septulatis, $6-10 \div 1-1,5$; paraphysibus obsoletis.

Hab. in ligno quercino vetusto in insula Runschala pr. Abo Fenniæ.

Hab. ad folia et petiolos Aceris platanoidis in Fennia australi. — Ascomata primitus subobconica, dein deorsum stipitato-constricta.

4. **Allophylaria clavuliformis** Karst. Rev. mon. p. 131, *Pezicula clavuliformis* Karst. M. F. I, p. 164. — Gregaria, stipitata, firmula, hyalino- v. lutescente-pallida, subinde sublutea v. rufescens, sicca lutescens, circiter 0,5-0,6 mm. alt., cupula convexa, subimmarginata, sicca concaviuscula, 0,3-0,5 mm. lat., stipite breviusculo, sensim in cupulam dilatato; ascis clavatis, 135-140 \approx 10-11, obturaculo jodo leviter violaceo-fusco; sporidiis plerumque distichis, oblongatis, spurie uniseptatis vel guttulis 2 majusculis praeditis, curvulis v. rectis, 11-20 \approx 4-4,5; paraphysibus gracilibus, 1,5 μ . cr. sursum leniter incrassatis.

Hab. ad caules Rubi idæi putrescentes, Mustiala Fenniæ.

5. **Allophylaria subliciformis** Karst. Rev. mon. p. 131, *Pezicula subliciformis* Karst. M. F. I, p. 164. — Gregaria, stipitata, firmula, 0,3 mm. alt., cupula lutescente, cum stipite brevissimo, crasso, pallido v. lutescente-pallido, obconica, disco planiusculo, 0,1-0,2 mm. lat.; ascis clavatis, 160-170 \approx 11-15; sporidiis subdistichis v. oblique monostichis, fusoideo-ellipsoideis v. fusoideo-oblongatis, guttulas 2 majusculas distincte foventibus, inæquilateralibus vel leniter curvulis, 15-26 \approx 6-7; paraphysibus numerosis, gracilibus, 1,5 μ . cr., apicem versus leniter incrassatis.

Hab. ad caules Artemisiæ vulgaris et Cerefolii dejectos aridos in Mustiala, ad caules Epilobii angustifolii in par. Messuby Fenniæ.

PEZICULA Tul. S. F. C. III, p. 182 [1865] (Etym. demin. a *Peziza*), *Dermatea*, *Cenangium*, *Peziza* Auct. — Ascoma mediocre, ceraceo-tenacellum, sessile, subinde erumpens v. furfurellum, patellatum v. cupulato-patellare, haud nigrum. Ascii paraphysati, typice majusculi, clavati, 4-8-spori. Sporidia ellipsoidea v. oblonga, hyalina, nucleata vel guttulata, continua, subhyalina. — A *Dermatella*, cui analoga, differt sporidiis continuis nec septatis, sed adsunt species sporidiis pseudoseptatis ambiguæ, hinc sub utroque genere inquirendæ.

1. **Pezicula carpinea** (Pers.) Tul. Carp. III, p. 183, Sacc. F. I, 1419, 1292 Pat. Tab. 221, *Peziza* Pers. Syn. p. 673, *Dermatea* Fr. Summa p. 362, *Cenangium Carpini* Wallr. — Ascomatibus subcæspitosis exiguis, primum clausis, mox in discum planum carneo-flaventem, margine dilutiore tenuiter farinoso cinctum apertis, sessi-

ibus, patellaribus, strato carnoso primum subcutaneo; ascis clavatis apice incrassatis obtusis, 200 ± 18 , jodoque viola scentedibus; sporidiis octonis ovato-oblongis, subinæquilateralibus, subdistichis, continuis nucleatis, hyalinis, $18-22 \pm 9-10$.

Hab. in ramis corticatis *Carpini Betuli* et *americanæ* in Germania, Gallia, Belgio, Italia, Amer. bor. — Ob ascomata initio erumpentia epidermideque fissâ cincta ad *Ocellariam* vergit. Quoad status secundarios cfr. Tul. loc. cit. — ***tetraspora** Sacc. Mich. II, pag. 613: asci cylindracei 120 ± 20 , brevissime stipitati, paraphysati; sporidia quaterna, oblique monosticha, oblongo-ellipsoidea, $24 \pm 12-14$, granulosa, hyalina. In cortice *Carpini* in Gallia (GILLET).

2. **Pezicula acericola** Peck in litt. et Sacc. Misc. Myc. Ser. II, n. 66. 1293

Nodularia acericola Peck 25 Rep. 98. — Cæspitosa, tenuis, subcarnosa, irregularis, pallide flava, primitus aperta, disco plano vel convexo, subpruinosa, margine obsoleto; ascis clavulatis, 4-8-sporis, $80-90 \pm 10-18$; sporidiis oblongo-clavulatis, sæpe curvulis, $24-26 \pm 8$, hyalinis; paraphysibus filiformi-clavulatis, subflexuosis.

Hab. in cortice *Acerum* North Elba Amer. bor. — Cæspituli e 3-7 cupulis formati, 2 mm. lat.

3. **Pezicula cinnamomea** (DC.) Sacc. Mich. II, p. 332, *Pez. cinnamomea* DC. fr. 6, p. 23, Pers. Myc. Eur. p. 268, *Dermatea* Phill. Disc. p. 342. 1294

— Erumpens, sessilis, carnoso-coriacea, tenuis, rubiginoso-lateritia, subtus pruinosa, disco subimmarginato; ascis clavatis, octosporis; sporidiis oblongis, utrinque rotundatis inæquilateralibus, 2-3-guttatis simulateque septatis, $20-30 \pm 8-10$.

Hab. ad *Quercus caudices* inter Lichenes ad Neufchatel et ad cort. *acerinum* pr. Bristol Britanniæ (BUCKNALL). — Gregaria v. fasciculata, parum carnosa, plana aut margine undulato-sinuata, 1-2 mm. lata, subtus glabra, sed pulverulenta, ut vix cum Persoonio ad *Lachneas* referri possit. — Forma **Castaneæ** Sacc. l. c, et Fung. ital. f. 1420: ascomatibus erumpenti-superficialibus, flavocinnamomeis, subfragilibus, sessilibus, ccomplanatis; ascis crasse clavatis 80 ± 18 , subsessilibus, apice rotundatis, octosporis; sporidiis distichis ellipsoideo-oblongis inæquilateralibus $20-25 \pm 8-9$, crasse 3-4-guttatis, hyalinis. In cortice *Castaneæ* vescæ, sociâ *Endothium gyroso* in Amer. bor. (RAVENEL).

4. **Pezicula rhabarbarina** (Berk.) Tul. S. F. C. III, p. 183. *Patellaria Rubi* Lib. Exs. 231, *Peziza rhabarbarina* Berk. Engl. T. V. 2, p. 197, *Lachnella rhabarbarina* Fr. S. v. Sc. p. 365, *Helotium*

Rubi Sprée in Rbh. F. eur. 717, *P. Rubi* Lasch. Bot. Zeit. 1845, p. 65, *Pez. arduennensis* Mont. Ann. Sc. nat. V, p. 287, t. XIII, f. 9. — Sessilis, patellata, flavo-fulva, sparsa, pilis clavulatis flavis minutis adspersa, carne rhabarbarina; ascis cylindraceo-clavatis, breve crasseque stipitatis, 96-110 \approx 13-18, paraphysibus filiformibus apice subincrassatis obvallatis, octosporis, apice jodo violascens-tibus; sporidiis oblongo-cylindraceis, utrinque rotundatis, 18-20 \approx 6-7, curvulis, granulosis, hyalinis.

Hab. ad ramos putrescentes Rubi fruticosi, cæsii et fastigiati in Gallia, Germania, Britannia, Italia, Belgio, in Rubo nec non in Fago, Corno in Amer. bor. — Quoad status secundar. cfr. Tul. loc. cit. An huc pertineat *P. rhab.* Hazsl. Magg. Kögg. t. XII, f. 20 sporidiis cuboideo-4-nucleatis, dubium. — ***P. Rosæ** Sacc. Mich. II, p. 323, Fung. ital. f. 1421: cupulis sparsis erumpentibus minutis, $1\frac{1}{6}$ - $1\frac{1}{8}$ mm. diam., brevissime stipitatis, crasse lenticularibus, disco subplanis, lutescenti-rubiginosis; ascis crasse clavatis substipitatis, 70 \approx 20, 8-sporis; sporidiis subdistichis oblongo-ovatis v. subcylindraceis, 28 \approx 10, continuis, hyalinis, tandem fuscescentibus. In ramis emortuis Rosæ caninæ in silva Montello Ital. bor. et Rouen Galliæ.

5. **Pezicula laricicola** Fuck. Symb. myc. p. 279. — Cupulis sparsis 1296 coacervatisve, subsessilibus, lentiformibus convexis, 1 mm. diam., emarginatis, carneo-ochraceis, pulverulentis; ascis elongatis, subclavatis, 8-sporis, 104 \approx 18; sporidiis oblongis, inæquilateralibus, 1-nucleatis, hyalinis, 24 \approx 8; paraphysibus linearibus.

Hab. in ramis siccis corticatis Laricis prope Oestrich Rheno-goviæ.

6. **Pezicula minuta** Peck Bull. N. Y. St. Mus. n. 2, p. 21 (1887). — 1297 Minuta, sparsa v. 2-3-aggregata, puncto exiguo matrici affixa, grisea, pulverulenta, margine obtuso v. obsoleto, disco plano v. convexo, subochraceo; ascis oblongo-clavatis; sporidiis stipatis oblongo-ellipsoideis, hyalinis, 20-25 μ . long.; paraphysibus filiformi-clavulatis.

Hab. in ramis emortuis Viburni lantanoidis in monte Catskill Amer. bor. — Ascoma $1\frac{1}{4}$ - $1\frac{1}{2}$ mm. diam.

7. **Pezicula myrtillina** Karst. M. F. I, p. 165. — Sparsa v. subgredaria, sessilis, convexa, immarginata, testaceo-pallida v. lutescens vel testacea, 0,4 mm. lat.; ascis clavatis, 90-100 \approx 14-18, obturaculo jodo leviter violaceo-fusco; sporidiis di- vel subdistichis, fusideo-oblongatis v. elongatis, rectis v. curvulis, guttulatis 1-2 ma-

jusculis præditis, 17-30 \approx 6-7; paraphysibus 1,5 μ . cr., apice sphæ-roideo v. obovoideo-incrassato, 8 mm. cr.

Hab. ad caules Myrtilli nigræ aridos in Mustiala Fenniæ.

8. **Pezicula saccharina** (Bresad.) Sacc. Mich. II, p. 536, *Peziza saccharina* Bresad. F. Trident. II, p. 24, t. XXIX, f. 1. — Convexa dein convexo-marginata vel explanato-marginata, subtus planiuscula, ligno adhærens, ceraceo-subgelatinosa, sessilis, 4-8 mm. lata, extus pallida, centro colore sacchari crystallizati, in siccis nigra, sub lente granulosa, carne hyalino-fuscidula, subpellucida, inodora; ascis cylindraceis, basi constrictis, 8-sporis, 360 \approx 18; paraphysibus apice clavatis, intus luteolo-granulosis; sporidiis oblongo-ellipticis, hyalinis, biguttulatis, oblique monostichis, 21 \approx 12; excipuli cellulis hexagonis.

Hab. in raminis corticatis Tiliæ parvifoliæ in agro Tridentino. — A *Pezicula carpinea*, cui valde accedit, distinguitur ascis cylindricis, sporidiis monostichis etc.

9. **Pezicula lilacina** Bresad. Trid. p. 68, t. 74, f. 1. — Sessilis, 1300 rassisime substipitata, ceraceo-tenacella, extus puberula, dein glabra, plano-concava, marginibus elevatis tumidiusculis, tota dilute lila-dea, 2-4 mm. lata, gregaria v. subcæspitosa, substantia ceraceo-carnosa subconcolore, inodora; ascis cylindraceo-stipitatis, apice operculo minuto jodo cœrulescente, 8-sporis, 180-210 \approx 8-12; paraphysibus filiformibus, ramosis, guttulatis, æqualibus, 1-2 μ . cr.; sporidiis ellipticis v. subcylindraceis, interdum uno latere subcurvatis, hyalinis, 14-16 \approx 7-8, intus minutissime granulosis.

Hab. ad ramulos Alni gelatinosæ locis humidis v. in rivulis sub aqua, Valle di Sole, Trento. — Ab *Ombrophila violacea* et *lilacina* certe distincta.

10. **Pezicula nectrioides** (Phill.) Sacc. *Dermatea nectrioides* Phill. 1301 Disc. p. 340. — Gregaria v. cæspitosa, erumpens, sessilis, subglobosa dein planiuscula, subcircularis, rubro-brunnea, disco aurantio-rubro, demum pallide brunneo; ascis tereti-clavatis; sporidiis oblongo-ellipsoideis vel oblongis utrinque rotundatis, 1-2-guttatis rectis curvulisve, 20-25 \approx 7; paraphysibus filiformibus ramulosis.

Hab. in conis Pini silvestris pr. Birmingham Britanniæ (W. B. GROVE). — Ascoma $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam. initio nectriæforme.

11. **Pezicula dryina** (Cooke) Sacc. *Dermatea dryina* Cooke Grev. 1302 VII, p. 62, Phill. Disc. p. 340. — Sparsa vel subinde 3-4-aggregata, convexo plana, carnea, 1 mm. diam., erumpens, margine vix vel non elevato, substantia carnosulo-lenta; ascis clavatis; spori-

diis ellipsoideis, utrinque subattenuatis, rectis curvulisve, 30-35
 \approx 10-12.

Hab. in cortice quercino pr. Hereford Britanniæ.

12. **Pezicula populea** Hazsl. Einig. Disc. p. 11, Magyar. szab. Disc. 1303
 tab. I, f. 18. — Stromati tubiculariaceo conidiophoro pulvinato
 inserta, scutellata, margine exiguo, obtuso, luteola; ascis tereti-
 clavatis, octosporis; sporidiis ovoideis, hyalinis, continuis; paraphy-
 sibus bacillaribus, apice conidium («?»), subreniforme exserentibus.

Hab. in radicibus defossis Populi nigræ pr. Eperies Hun-
 gariæ. — Ascoma habitu *Peziculæ carpineæ*.

13. **Pezicula rivularis** (Cr.) Sacc. *Patellaria rivularis* Crouan Fl. 1304
 Finist. p. 45. — Ascomate extus brunneo, 2-4 mm. diam., cras-
 siusculo, disciformi, margine sublobato, disco ochraceo; ascis cy-
 lindraceis, octosporis; sporidiis ovoideis, granulosis, monostichis,
 limbatis; paraphysibus ramulosis, sursum crassiusculis, granulosis.

Hab. in ramis emortuis Quercus rivulo immersis, Finistère
 Galliæ. — Contextus prosenchymatico-granulosus, densus.

14. **Pezicula Herminiæ** Rabenh. in Hedwigia 1878, p. 146. — Gre- 1305
 garia, plus minus approximata vel sparsa, sessilis, nuda, glabra,
 ceraceo-carnosa, vix 1 cm. lata, sicca plana v. concaviuscula, hu-
 mectata concaviuscula; disco aurantio-luteo, non nunquam expal-
 lescente, margine tumidulo, integerrimo dilutiore; ascis breve cla-
 vatis, jodo non tinctis; paraphysibus filiformibus intertextis, octo-
 sporis; sporidiis oblongo-fusiformibus, recte v. oblique monostichis
 (passim subdistichis), 6 \approx 3, hyalinis, medio spurie uniseptatis.

Hab. ad folia emorta, Muscos etc. in Guadalupa (L' HERMINIER).

15. **Pezicula Kalmiæ** (Peck) Sacc. *Peziza Kalmiæ* Peck 25 Rep. St. 1306
 Mus. p. 99, *Dermatea* Cooke Disc. U. S. II, p. 23. — Minuta, ses-
 silis, subplana, marginata, extus furfuracea, sordide grisea, margine
 initio incurvato, disco rufo-brunneo; sporidiis ellipsoideis, plerum-
 que nucleatis, 10 \approx 5.

Hab. in ramis emortuis Kalmiæ angustifoliæ subinde su-
 pra Dothidellam Kalmiæ, Sandlake Amer. bor.

16. **Pezicula Cenangium** (De Not.) Sacc. *Peziza Cenangium* De Not. 1307
 Dec. IV, n. I, E. C. It. n. 637, Rab. et Gonn. M. E. p. 10, t. VI.
 f. 1. — Ascomatibus gregariis, flavo-virentibus erumpenti-superfi-
 cialibus, 1 mm. lat., rarius cæspitulosis, subcoriaceis, cupulatis, dein
 expansis subsessilibus subcontortis, disco albido, nitidulis; ascis te-
 reti-clavatis, filiformi-paraphysatis; sporidiis oblongo-ellipsoideis,
 ex hyalino citrinis, 12-14 \approx 4-5 (De Not.), 9-12 \approx 3-4 (Rab.).

Hab. ia ramulis dejectis Rhododendri et forte Laricu m
in monte Cenisio et Valsesia. — «Quoad genus dubia stirps».

17. **Pezicula viridi-atra** (B. et C.) Sacc. *Pez. viridi-atra* B. et C. 1308
Cub. Fungi n. 692, nec Fr. — Sessilis, atro-viridis, subtus granu-
lata, undulata, irregularis; sporidiis ellipticis demum subfuscis, 5
 μ . longis.

Hab. in ligno putrescente, cum Polypori specie in insula
Cuba. — Ascoma 8 mm. lat., lignum haud fœdans. Verisimiliter novi
generis typus.

18. **Pezicula? Philadelphi** (Schw.) Sacc. *Peziza Philadelphi* Schw. 1309
Syn. 177. — Cupulis sessilibus aut stipite papillæformi præditis,
interdum aggregatis, siccis subrotundis, sparsis superficialiter epi-
dermidi adnatis, extus ferrugineo-lutescentibus, subpulverulentis,
marginè subprominulo, disco demum convexo, punctato, umbrino
2 mm. lat.

Hab. in ramulis Philadelphi inodori culti in hortis Beth-
lehem Amer. bor.

Species indescripta.

19. **Pezicula Cratægi** (Auersw. in Sched.) Fuck. Symb. App. II, p. 56 1310
Hazsl. Mag. szab. Disc. p. 220.

Hab. in ramis duris corticatis Cratægi Oxyacanthæ, rarius,
pr. Neuchâtel Helvetiæ et in Hungaria. — Sporidia sec. Hazsl. l. c.
22-24 = 10.

CHLOROSPLENIUM Fr. Summa p. 356 [1849] (Etym. *chloros* viridis
et *splen* lien, ob colorem). Ascoma breve stipitatum, ceraceum,
cupulatum vel subplanum, saepè flexuosum, æruginosum v. oliva-
ceum, matricem saepius colorans. Ascii tereti-cavati, paraphysati,
octospori. Sporidia ovoidea v. fusideo-oblonga, continua, guttulata,
subhyalina.

1. **Chlorosplenium æruginosum** (Eder) De N. Disc. pag. 22, nec Tul. 1311
Helvela æruginosa Ed. Fl. Dan. t. 534, f. 2, *Peziza æruginosa*
Vahl Flor. Dan. t. 1260, f. 1, Pers. Syn. Fung. p. 617, *Hel. æru-
ginosum* Fr. Summa p. 355, Sacc. F. it. f. 1348. — Breviter sti-
pitatum, æruginosum; cupula turbinata, dein expansa et subflexuosa,
disco albicante, 1-4 mm. lat.; ascis tereti-clavatis, 65-75 = 6, obtu-
raculo jodo leviter cœrulecente; sporidiis plerumque distichis, fu-
soideo-elongatis, rectis v. curvulis, hyalino-chlorinis, plerumque
2-guttatis, 10-14 = 3,5-4; paraphysibus sat numerosis.

Hab. ad lignum mucidum *Quercus*, *Betulae*, *Alni*, *Fagi*, *Populi* in *Suecia*, *Fennia*, *Germania*, *Britannia*, *Sibiria*, *Italia*, *Gallia*, *Belgio*, *Amer. bor.*, *Africa austr.*, *Australia*, *Fuegia*. — Quoad spermogonia cfr. *Tul.* l. c. — Var. **subgriseum** Holmsk. *Ot.* pag. 28, t. 12. In *Dania*. — Admodum versiforme, lignum colore æruginoso late profundeque tinctum passim adhibetur in usum fabrile.

2. ***Chlorosplenium æruginascens*** (Nyl.) Karst. M. F. I, p. 103, *Pez.* 1312
æruginascens Nyl. Obs. *Pez.* p. 42, *Ch. æruginosum* Multorum, Tulasne Carp. III, p. 187, t. XX, f. 15-19. — Stipitatum, æruginosum, cupula concava, dein expansa et flexuosa, stipite plus minus elongato aut abbreviato, subinde fere nullo; ascis cylindraceo-clavatis v. cylindraceis, tenellis, 50-60 \approx 4, obturaculo jodo levisime cœrulecente; sporidiis plerumque distichis, fusoideo-elongatis, rectis v. curviusculis, ut plurimum guttulatis, 6-8 \approx 1.5-2.5; paraphibus gracilibus.

Hab. ad lignum mucidum arborum frondosarum et *Pini* per totam *Fenniam*, *Laponiam*, *Galliam*, *Germaniam*. — Spermogonia discreta v. cæspitosa, interdum ascomatibus recentibus sociata, sessilia sphæroidea, glaberrima, nonnihil rugosa, atro-virentia multilocularia, loculis angustis periphericis, poris numerosis minutissimis instructa sterigmata filiformia, cæspitosa, ramosa, circiter 16 μ . long., sporulæ elongatæ, rectæ, simplices, circiter 3.5-5 μ . long.

3. ***Chlorosplenium omnivirens*** (Berk.) Cooke Austr. Fung. p. 51. — 1313
Peziza omnivirens Berk. Ph. Tasn. p. 275. — Æruginosum, cupula breviter stipitata subturbanata, hymenio plano; sporidiis amplis 18-20 μ . longis.

Hab. in ligno putrescente, Leith Creek Tasmaniae (ARCHER). — Præcipue forma magis regulari et sporidiis magnis dignoscitur a *Ch. versiformi* et *æruginoso*.

4. ***Chlorosplenium æruginellum*** (Nyl.) Karst. M. F. I, p. 104, *Peziza* 1314
æruginella Nyl. Obs. *Pez.* p. 40. — Sparsum, stipitatum, æruginosum, cupula planiuscula, tenui, subinde subfle Xuosa, 0.4-0.9 mm. lat., stipite æquali, 0.3-0.5 mm. long.; ascis cylindraceo-clavatis, 46-56 \approx 5-7, apice jodo vix vel non tinctis; sporidiis distichis vel oblique monostichis, fusoideo-elongatis, guttulatis, 7-12 \approx 2-3.

Hab. ad caules *Spireæ* *Ulmariæ* putrescentes prope Mustiala Fenniae.

5. ***Chlorosplenium versiforme*** (Pers.) Karst. M. F. I, pag. 102, *P.* 1315
versif. Pers. ic. et deser. p. 25, t. 7, f. 7, *Syn.* p. 647, *Fr. Syst.*

Myc. II, p. 130. — Gregarium, substipitatum, cupula difformi sub-contorta, flavo-olivacea, externe subpurpurascens; ascis tereti-clavatis, 90-100 \approx 7-8; sporidiis submonostichis, ovoideo-oblongatis, saepe curvulis, 9-14 \approx 3-4, guttulatis spurieve septatis; paraphysibus gracilibus.

Hab. in truncis abiegnis rarius betulinis, quercinis et fraxineis in Britannia, Germania, Belgio, Gallia, Helvetia, Amer. bor. — Primo subvillosum, mox nudum, colore olivaceo-flavescente, forma nunc regulari, nunc compressa subundulata, disco convexo v. depresso.

6. **Chlorosplenium brasiliense** B. et Cooke Fung. Bres. n. 434. — 1316
Æruginosum; cupula turbinata demum expansa, stipite brevi æquali; ascis cylindraceis, 45 \approx 2; sporidiis linearibus 3-4 μ . longis; paraphysibus filiformibus.

Hab. in lignis Rio Tapajos Brasiliæ (TRAILL). — Sporidia non-dum matura, tamen tenuiora videntur quam *Ch. aeruginosi*. Matrix non decolorata.

7. **Chlorosplenium fuegianum** Speg. F. Fueg. n. 338. — Cupulis hinc 1317 inde gregariis, udis infundibuliformi-explanatis 1-2,5 mm. diam., siccis contractulis involutis, extus intusque sordide atro-olivaceis, extus in juventute rufescenti- v. fulvescenti-pruinulosis dein glabratris, margine integerrimis, tenacellis, rigidulis, postice in stipitem brevem et crassiusculum, 1-2 mm. long., 0,5 mm. cr., saepe basi connatum concolorem glabrum vel olivaceo-pruinulosum cuneato-attenuatis; ascis clavulatis, antice rotundatis crassiuscule tunicatis non v. minute bifoveolatis, postice longe stipitato-attenuatis, p. sp. 40 \approx 8, ped. 70-90 \approx 3-5, octosporis; paraphysibus filiformibus densiusculis; sporidiis apice ascorum subconglobatis, recte distichis, elliptico-fusoideis, utrinque acutato-obtusatis, 10 \approx 4-5, hyalinis, continuis (an per ætatem 1-3-septatis?)

Hab. ad ramos decorticatos putrescentes Fagi betuloidis prope Port Vancouver in Staten island Fuegiæ. — Jodi vis nulla; lignum maculis latiusculis fusco-olivaceis, saepe furfure rufescenti conspersis donatum.

8. **Chlorosplenium Puiggarii** Speg. Fung. Arg. Pug. III, n. 226. — 1318
Stipitatum, pulchre viridi-æragineum, glaberrimum; cupulis primo regularibus, concavis, dein lateraliter expansis, 5-15 mm. diam., irregularibus, undulatis, crassiuscule membranaceis, coriaciusculis; stipite brevi, 1-3 \approx 0,5 mm. quandoque fere nullo, excentrico, sursum æragineo, inferne sordide fuscescente; ascis cylindraco-clava-

tis, sursum rotundatis crasseque tunicatis, deorsum brevissime stipitatis, 40-45 \approx 3-3,5; paraphysibus simplicibus, filiformibus, hyalinis obvallatis, octosporis; sporidiis elliptico-fusoideis, utrinque attenuato-rotundatis, sœpe inæquilateralibus, 5-7 \approx 1-1,5, 2-guttulatis, hyalinis; obturaculo ascorum jodi ope cœrulescit; substratum profunde æruginascens.

Hab. in truncis ramisque decorticatis, dejectis putrescentibus, in nemorosis prope Apiahy, Brasilia meridionali (Dr. J. PUIGGARI). — An hoc *Ch. brasiliense* Berk. et Br. ducendum? An unum et alterum *Ch. æruginascentis* (Nyl.) Karst. tautum varietates?

9. **Chlorosplenium æragineum** (Berk.) Sacc. *Peziza æraginea* Berk. 1319 Dec. of Fungi n. 474. — Cupulis cyathiformibus, subregularibus, extus æragineo-obscuris, disco pallidioribus; ascis linearibus; sporidiis subfusiformibus, 7-8 μ . longis, mycelio lignum superficialiter viridi-tingente.

Hab. in lignis putridis Khasia Indiae (HOOKER et THOMPSON). — Ascomata 2 mm. lata, sparsa, extus metallico-viridia; stipes cupulam vix æquans, atro-virens. Affine *Ch. æruginoso* sed magis regulare, tenuius, obscurius sporidiisque minoribus. Sec. Cooke spor. 10-12 μ . long.

10. **Chlorosplenium Kriegerianum** (Rab.) Sacc. *Ciboria Kriegeriana* 1320 (Rab.) Rehm Asc. n. 660, *Ombrophila?* *Kriegeriana* Rab. Hedw. 1878, p. 31. — Turbinatum demum profunde cyathiforme, tremulum, distincte stipitatum, olivaceum, nudum subnitens, margine initio inflexo dein erecto, nigro-denticulato, stipite crassiusculo, solido, concolore; ascis elongatis clavatis, substipitatis, 96-100 \approx 6; sporidiis ellipsoideis, rectis vel subcurvulis, biguttatis, hyalinis, 12-18 \approx 6; paraphysibus filiformibus, clavulatis, subolvaceis.

Hab. ad ramos in Germania et Helvetia. — Sec. Rhem l. c., ascomate distincte prosenchymatico ad *Ciboriæ* spectat, teste Winter (Hedw. 1881, p. 70) a *Chl. elatino* (A. et S.) vix differt.

11. **Chlorosplenium elatinum** (A. et S.) Sacc. *P. elatina* A. S. p. 330, 1321 t. 2, f. 3, Pers. Myc. Eur. p. 291, Fr. Syst. Myc. II, p. 112, *Helotium* Quél. Ench. pag. 309. — Obconicum, viride, extus nigro-striatum, margine lato infexo, stipite firmo sursum incrassato.

Hab. ad ramos emortuos humi jacentes Abietum, frequens in Lusatia, Helvetia etc. — Ceraceum, firmum, crassiusculum, 4-7 mm. altum, 6-10 mm. latum, extus fibrillis nigris striatum vel venoso-pilosum, margine subserrato. Initio, præsertim intus, fusco-virens; dein totum viride, disco demum explanato sordide lutescente.

12. ***Chlorosplenium lividum*** (A. et S.) Karst. Rev. mon. pag. 124, *P.* 1322
versiformis *β. livida* A. et S. Consp. p. 314, *Pez. versiformis* *β.*
nigrescenti-olivacea Weinm. Hymen. Gast. pag. 467, *Chlor. nigre-*
scenti-olivac. Karst. Symb. M. F. IX, pag. 56, *Rutströmia bulga-*
rioides (Rab.) Karst. M. F. I, pag. 105, *Peziza bulgarioides* Rab.
Fung. eur. II, 1008, *Peziza ciborioides* Fr, *P. strobilaria* Nyl. Obs.
 p. 36. — Gregaria, stipitata subsessilis, primum cyathoidea et regula-
 ris dein expansa, difformis et undulata, initio livida, dein fuliginea v.
 subolivacea, tandem nigrescens v. cinerescens, disco concolori, saepe
 ruguloso, margine integro, 2-8 mm. lat.; ascis subcylindraceis, 90-
 125 \approx 6-8, obturaculo jodo dilute cœrulecente; sporidiis monostichis,
 ellipsoideis vel oblongatis, rectis, eguttulatis, 6-11 \approx 4-4,5;
 paraphysibus gracilibus.

Hab. ad strobilos Abietis e Fennia meridionali ad prov. Öster-
 botten (Wasa, Jakobstad) usque, in Germania, Hungaria. — Cum
Chl. versiformi, cuius varietatem *b. lividam* Fr. Syst. Myc. II,
 p. 131, forsan sistit, multum commune habet.

13. ***Chlorosplenium repandum*** Fr. Summa pag. 356 in nota, *Peziza* 1323
chlorascens Schw. Syn. 175, Berk. Cub. Fung. n. 690. — Maju-
 sculum, 7-10 mm. diam., gregarium, siccum valde contortum, cra-
 teriforme, sessile aut brevius stipitatum, extus floccoso-fibrosum,
 nigrum aut nigro-fuscum (statu sicciori), madefactum autem ærugi-
 nosum, disco lato chlorascenti aut luteo-viridi, demum subpulve-
 rulento; ascis, teste Berk., clavatis; sporidiis fusoideis 12-15 μ . long.

Hab. ad cortices Taxodii, Quercus etc. in Carolina, Pen-
 sylvania et ins. Cuba.

14. ***Chlorosplenium amenticolum*** Karst. Fung. nov. Fenn. p. 1 (Rev. 1324
 myc. 1887). — Sparsum, ceraceum, stipitatum, pruinatum v. fur-
 furaceum, subsulphureo-virens, cupula e concavo explanata, saepe
 repanda, 1-2 mm. lata, stipite cupulæ latitudine vel infra; ascis
 cylindraceo-subclavatis, 55-60 \approx 6; sporidiis oblique monostichis,
 ellipsoideis v. suboblongatis, simplicibus, 6-7 \approx 3; paraphysibus pa-
 rum notabilibus.

Hab. in amentis Alni glutinosæ emortuis prope Rouen in
 Gallia (LETENDRE).

15. ***Chlorosplenium chloromelum*** (Phill. et Hark.) Sacc. *Peziza chloro-* 1325
mela Phill. et Hark. Grevill. vol. XIII, pag. 22. — Sparsum v.
 congregatum, stipitatum, 1 mm. long., glabrum, atro-viride, sti-
 pite pallidiori, disco plano, immarginato, luteo-viridi 0,6 mm.; ascis
 cylindraceo-clavatis, octosporis; sporidiis fusiformibus v. clavatis;

plerumque curvulis, hyalinis, viridi-tinctis, 20-25 \times 4-5; paraphysibus filiformibus, indistinctis cohærentibusque, hymenio granuloso, subgelatinoso.

Hab. in foliis *Sequojae semperfirantis* in California (HARKNESS).

16. ***Chlorosplenium epimyces*** Cooke Disc. U. S. I, pag. 299. — Ses- 1326
sile, viridi-flavum dein olivaceum, pulverulentum, 2 mm. latum,
planum, disco pallidiore; ascis clavatis; sporidiis distichis, late fu-
soideis, 20 \times 10.

Hab. ad Corticium vetustum New Jersey Amer. bor. (ELLIS).

— Pulvis e granulis globosis brunneis 7 μ . diam. constans.

17. ***Chlorosplenium luteo-virens*** (Fr.) Sacc. *P. luteo-virens* Fr. Syst. 1327
Myc. II, p. 133, *Helotium* Fr. Summa p. 356. — Turbinatum, ses-
sile, carnosо-ceraceum, glabrum, viridi-luteum, disco plano immar-
ginato.

Hab. ad ramos putridos dejectos Fagi, Feinsio in Suecia. —
Distinctissimum, 1 mm. tantum latum, substantia carnosо-ceracea,
pallescente. Cupula late turbinata, absque stipite distincto, junior
quidem concava, sed vetusta etiam convexa.

18. ***Chlorosplenium Schweinitzii*** Fr. Summa Veg. p. 356, nota, *Pez.* 1328
Chlora Schw. Carol. n. 1235. — Sessile, carnosо-ceraceum, luteo-
virens, margine plicato-crenato, extus squamuloso-pulveraceum.

Hab. frequens ad latera interiora truncorum Carolinæ. — Cfr.
Ellis N. A. F. n. 664.

19. ***Chlorosplenium tortum*** (Schw.) Fr. Summa p. 356, in nota, *Pez.* 1329
ziza torta Schw. Syb. p. 175. — Subfasciculata aut sparsa; sti-
pite crasso sulcato, torto ferrugineo-fusco æque ac cupula extus,
disco explanato undulato, immarginato, æruginoso-viridi, pulveru-
lento, substantia demum subcoriacea.

Hab. raro in lignis vetustis, Salem Amer. bor. — Totus fungus
8 mm. latus et altus. Affinis forte *Cib. bolari*.

CORONELLARIA Karst. M. F. I. pag. 14 [1871] (Etym. *corona*). —
Ascoma obconicum, sessile, glabrum, margine pallide ciliato v. fimbriato,
excipuli contextu fuscescente. Asci oblongo-clavati, octo-
spori, paraphysati. Sporidia fusoideo-oblonga v. acicularia, conti-
nua guttulata, hyalina. — A gen. *Cyathicula* (Sect. II) differt exci-
pulo fuscescente, sporidiis majoribus.

1. ***Coronellaria Delitschiana*** (Auersw.) Karst. M. F. I, p. 183, *Pez.* 1330
Delitschiana Auersw. in Rab. F. E. n. 912 Hedw. V, p. 189. —

Gregaria aut sparsa, sessilis, subobconica, margine fimbriato, testaceo-pallida v. pallida, dein fusco-pallida v. interdum (sicca) nigrescens, disco planiusculo, griseo-pallido v. lutescente, subinde (sicco) albicante, 0,2-0,5 mm. lat.; ascis ellipsoideo-oblongatis aut oblongato-clavatis, sessilibus, 75-110 = 16-20; sporidiis di-tristichis, oblongatis aut fusoideo-oblongatis v. fusoideo-elongatis, rectis v. inaequilateralibus, guttulas 1-4 majusculas foventibus v. demum spurie tenuiter septatis, 25-40 = 6-7; paraphysibus crassiusculis.

Hab. in calamis *Scirpi lacustris* vetustis in Germania et Fennia.

2. **Coronellaria pulicaris** Karst. M. F. I, p. 184. — Sparsa vel sub- 1331 gregaria, obconica aut hemisphærico-cupulata, albida v. lutescens v. pallide testacea v. viridi-lutea dein fusco-pallida v. lutescens vel (sicca) nigrescens, margine fimbriato, sicco albicante, 0,2-0,3 mm. lat.; ascis oblongato-clavatis, sessilibus, apice saepe attenuatis, 55-85 = 12-16, obturaculo violaceo-fuscescente; sporidiis di-tristichis, interdum oblique monostichis, oblongatis aut fusoideo-oblongatis v. aciculari-oblongatis, rectis v. inaequilateralibus, vulgo 2-guttulatis, 14-24 = 3,5-4,5; paraphysibus mollibus, flexuosis, 1,5 p. cr. apicem versus incrassatis.

Hab. in calamis et foliis *Scirpi silvatici* et *Sc. lacustris* vetustis in Fennia australi. — *C. Delitschianâ* omnibus partibus minor, ceterum peraffinis.

3. **Coronellaria caricinella** Karst. Symb. p. 248, M. F. I, p. 185. — 1332 Gregaria v. sparsa, obconica, pallida v. fuscescente-pallida, margine albo-ciliato, disco pallido, 0,1-0,2 mm. lat.; ascis ellipsoideo-oblongatis aut oblongato-clavatis, 42-50 = 10-13; sporidiis di-vel tristichis, oblongatis, spurie uni-triseptatis aut guttulis 2 majusculis præditis, rectis, 13-16 = 3,5-4; paraphysibus crassiusculis, sursum incrassatis.

Hab. in foliis *Caricum majorum* putrescentibus in Fennia meridionali. — Ascomata interdum lutea v. aurea conspiciuntur.

MOLLISIA Fr. Syst. Myc. II, p. 137 [1822], ut subg., emend. (Etym. *mollis*), *Niptera* et *Micropeziza* Fuck. ex p., *Mollisia* Karst. et Sacc. ex p., non *Niptera* Fr. — Ascoma superficiale v. suberumpenti-superficiale, sessile, minutum, ceraceum scutellato-patellare, excipulo fusco, duriusculo, disco initio pallente. Asci filiformi-paraphysati, 4-8-spori. Sporidia ellipsoidea oblonga, fusoidea, teretiuscula, continua, hyalina.

A. Eu-Mollisia. Ascoma subcirculare, subsuperficiale. Sporidia breviuscula.

I. Caulicolæ.

1. **Mollisia Polygoni** (Lasch) Gill. Disc. p. 120, *Pez. Polygoni* Lasch 1333
in Rab. H. M. n. 1127, Desm. 24 Not. pag. 7, *Niptera Polygoni*
Rehm Ascom. n. 70 (1871), Wint. Flora 1872, pag. 542, *Peziza*
luctuosa Cooke in Hedwigia 1875, pag. 83. — Gregaria, sessilis,
primo hemisphærica, margine incurvo, crenulato, demum aperta,
diformis, extus fuligineo-badia, hyalino-papillata, basi pilis ara-
chnoideis repentibus hyalinis cincta, disco e concavo piano conco-
lore, dein pallente, 0,5-1 mm. lato; ascis tereti-oblongis, sessili-
bus, 30-40 \approx 5-6; sporidiis elongato-clavulatis rectis curvulisve,
utrinque uniguttulatis, 7-9 \approx 2; paraphysibus filiformibus.

Hab. in caulis Polygonorum putridis in Hungaria, Transylania, Germania, Britannia, Gallia, America boreali. — Var.
Rumicis Sacc. F. Gall. n. 2231: ascis tereti-clavatis, paraphysatis,
35-38 \approx 7; sporidiis distichis, oblongo-fusoideis, 6-8 \approx 1 $\frac{1}{2}$ -2, 2-gut-
tulatis, hyalinis, rectis. In caule Acetosæ emortuo Rouen Gal-
liæ (LETENDRE).

2. **Mollisia atro-cinerea** Cooke in Phill. Disc. p. 176. — Gregaria v. 1334
sparsa, sessilis, leniter concava v. plana, extus nigra, glabra, disco
cinereo; ascis clavatis; sporidiis fusoideis vel cylindraceis, rectis
curvulisve, 5-7 \approx 1; paraphysibus non visis.

Hab. in caulis herbaceis e. g. Dipsaci in Britannia et Hun-
garia et in caulis Solidaginis, Hyperici, Desmodii in
Amer. bor. — Ascoma $\frac{1}{3}$ - $\frac{1}{2}$ mm. lat. Sec. Rehm Asc. Lojk. n. 38:
asci 24-30 \approx 5; sporidia 6 \approx 1,5.

3. **Mollisia Solidaginis** Karst. Rev. mon. atque Syn. p. 173. — In- 1335
nato-erumpens, concaviuscula, sicca contracta, margine lacero-fi-
broso, fuscescens, disco albido; ascis fusoideo-clavatis, obturaculo
jodo cœrulecente, 55-60 \approx 7-8; sporidiis distichis, elongatis, cur-
vulis, 12-15 \approx 2,5-3, 2-guttulatis.

Hab. in caulis aridis Solidaginis Virgaureæ, Kola Lap-
poniæ rossicæ.

4. **Mollisia fusco-rubra** (Rehm) Sacc. *Niptera* Rehm Ascom. Ill. n. 69, 1336
Winter Flora 1872, p. 526. — Cupulis gregariis vel basi congre-
gatis, hemisphaericis, ovatis v. diformibus, glabris, rugulosis, sessi-
libus extus fuscellis v. fusco-rubris, margine involuto v. incurvo,
fisso et hyalino-fimbriato (fimbriis basi pallide luteolis, antice hya-
linis, marginem dense fasciculatimque vestientibus), disco concavo-

luteo-rubro, 0,8-1,0 mm. latis; ascis linear-i-cylindraceis, sessilibus 8-sporis, 50-64 \approx 5-6; paraphysibus crassis, filiformibus, guttulatis, ascos æquantibus; sporidiis breviter cylindraceis, rectis subcurvatisve, hyalinis, utrinque uni-guttulatis et paullulum attenuatis, 10 \approx 2.

Hab. in caulis putridis *Artemisiæ campestris* prope Sugenheim Germaniæ.

5. **Mollisia erigeronata** Cooke Hedw. 1875, pag. 83, Disc. U. S. I, 1337 p. 290. — Gregaria, ceraceo-mollis, sessilis, cupula hemisphærica, demum applanata, atro-umbrina, nigrescens, $\frac{1}{2}$ mm. lata, disco livido cinereo, margine leniter elevato; ascis brevibus clavatis; sporidiis biseriatis v. stipatis, elliptico-linearibus, 10 \approx 2.

Hab. ad caules *Erigeronis*, New Jersey Amer. bor. (ELLIS).

6. **Mollisia cœrulans** Quél. X Suppl. p. 13, t. IX, f. 10, Grev. 1879, 1338 p. 37. — Ascomatibus gregariis sæpe confluentibus, discoideis, 500-800 μ . diam., crassis, ceraceis, teneris, griseis, margine pruinoso, albescente; hymenio convexo-plano, obscure cœruleo-nigrescente; sporidiis acicularibus, 10 μ . longis, curvatis.

Hab. gregarie in caulis plantarum majorum e. g. *Eupatoriæ*, Jura Galliæ.

7. **Mollisia astericola** C. et E. Grev. VI, p. 90. — Sparsa, mollis; cupulis demum applanatis, extus atro-brunneis; disco aquoso; ascis cylindraceo-clavatis; sporidiis linearibus, leniter curvulis, 10 μ . longis; paraphysibus filiformibus.

Hab. in caulis *Asteris*, New Jersey Amer. bor. — Ascoma $\frac{1}{2}$ -1 mm. lat. Forte huc spectat *P. Artemisiæ* Lasch.

8. **Mollisia sarmentorum** Sacc. F. Ven. Ser. IV, n. 72 (*Niptera*), F. 1340 ital. fig. 1383. — Cupulis minutissimis, gregariis, plano-scutellariis, sessilibus, $\frac{1}{6}$ mm. diam., griseis margine pallidiore, excipulo fuscescente; ascis clavatis, 32 \approx 8, deorsum attenuatis, subsessilibus; paraphysibus cylindraceis obvallatis, 4-sporis; sporidiis fusoideis, 11-12 \approx 3, curvulis initio 2-3-guttulatis, hyalinis.

Hab. in sarmentis *Humuli Lupuli*, Selva (Trev.) Ital. bor. — Affinis *M. Mercurialis* et *M. Polygoni* sed sporidiis quaternis amplioribusque præ ceteris differt.

9. **Mollisia urticicola** Phill. Disc. p. 177. — Gregaria, sessilis, sub-hemisphærica, demum subplana, extus glabra, pallide rufo-brunnea, margine fimbriato, disco e pallido-cinereo brunneolo; ascis cylindraceo-clavatis; sporidiis fusoideis vel cylindraceis, 7 \approx 1; paraphysibus filiformibus.

Hab. in caulis Urticæ locis inundatis, Shelton Rough Britanniae. — Ascoma 0,5 mm. lat.

10. **Mollisia lycopincola** Rehm Asc. n. 910. — Ascomatibus poculi-¹³⁴² formibus, dein discoideis, circularibus, rarius angulosis, margine tenui albido, mox inflexo, disco appланato extus nigricante, $\frac{1}{2}$ -1 mm. lat.; ascis clavatis, 30-40 \approx 5-6; sporidiis oblongis, rarius clavatis, rectis curvulatis, utrinque uniguttulatis, 7-10 \approx 2-2,5; paraphysibus filiformibus.

Hab. in caule putri Lycopi europæi prope Berlin Germaniæ (SYDOW). — Excipulum sphærico-parenchymaticum, brunneum; ascomata macula nigra incidentia.

11. **Mollisia brachyspora** Sacc. Fung. Ven. Ser. IV, n. 74 (*Niptera*) ¹³⁴³ F. ital. f. 111. — Cupulis sparsis gregariisque, minutis, $\frac{1}{6}$ - $\frac{1}{8}$ mm. diam., totis ochraceo-pallidis, margine pallidore, ex urceolato patellaribus, sessilibus, levibus, glabris, disco concolori concaviusculo, siccis varie tortis v. conchiformibus, excipuli contextu anguste prosenchymatico lutescente; ascis cylindraceo-clavatis vel subfusoideis 50 \approx 6, p. s. 35 long.; paraphysibus parcis, filiformibus, obvallatis, 8-sporis; sporidiis inordinatis, ovoideis, 6-7 \approx 3 $\frac{1}{2}$ -4, plerumque 2-guttulatis, hyalinis.

Hab. in caule putrescente Galeopsidis grandifloræ, Selva (Treviso) Ital. bor. — Sporidiis subovoideis, brevibus, a ceteris speciebus mihi notis diversa. Ad *Pezizellam* accedit.

12. **Mollisia Grappensis** Sacc. Mich. I, pag. 442 (*Niptera*), F. ital. ¹³⁴⁴ f. 1385. — Ascomatibus ceraceo-tenacellis, erumpenti-superficialibus, gregariis, sessilibus, scutellaribus, brunneis, disco margineque breve fibrilloso pallidioribus, $\frac{3}{4}$ - $1\frac{1}{4}$ mm. diam., contextu excipuli minute celluloso fuligineo, marginis subradiante dilutiore; ascis cylindraceis, breve stipitatis, apice, tunica integra, rotundatis, 70-80 \approx 12-13; paraphysibus bacillaribus obvallatis, octosporis; sporidiis distichis, oblongo-cylindraceis, inaequilateralibus, 15-17 \approx 5, utrinque rotundatis, 3-nucleatis simulateque 1-septatis, hyalinis.

Hab. in caulis emortuis Myrrhidis odoratæ in Monte Grappa, Bassano, Ital. bor. (BIZZOZERO). — Ascomatibus majusculis tenacellis, sporidiis ascisque crassis ab affinibus speciebus dignoscitur.

13. **Mollisia fuscidula** Sacc. F. Ven. Ser. IV, n. 73, F. ital. f. 1381 ¹³⁴⁵ (ut var. *M. Dehnii*). — Cupulis sparsis scutellaribus, minutis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., sessilibus, siccis margine contortis, ceraceo-duriusculis, disco griseo-lutescente, concavo, marginato, excipulo obscure fuli-

gineo, villo nullo insidentibus; ascis cylindraceo-clavatis, 50-60 \approx 6-7, brevissime stipitatis; paraphysibus filiformibus obvallatis, 8-sporis; sporidiis inordinate distichis, fusoideis, 10-12 \approx 2½-3 rectiusculis hyalinis.

Hab. in caule emortuo *Vincetoxicum* a Vittorio (Treviso) Ital. bor. — Affinis quoque *M. Polygoni*, a qua cupula pallidiore, villi defectu et sporidiis longioribus satis differt.

14. **Mollisia Euphrasiæ** (Fuck.) Sacc. *Nipt. Euphrasiae* Fuck. Symb. 1346 mycol. p. 293. — Cupulis gregariis, sessilibus, 2 mm. latis, irregularibus, extus rugulosis, fusco-nigris, margine distincto, denticulato, demum complicato, disco griseo; ascis anguste elongatis, 8-sporis, 64 μ . long.; paraphysibus asci longitudine, simplicibus; sporidiis farctis, anguste clavatis, saepe curvatis, continuis, multiguttulatis, hyalinis, 18 \approx 2-3.

Hab. in caulis siccis adhuc stantibus *Euphrasiæ* luteæ pr. Budenheim Rhenogoviæ.

15. **Mollisia Dehnii** (Rabenh.) Karst. M. F. I, p. 20, *Peziza Rabenh.* 1347 bot. Zeit. 1842, p. 12. — Gregaria, innato-sessilis, demum planiuscula, fusca, sicca nigricans, disco livido-cinereo, sicco saepe subgilvo, 0,5-1 mm. lat.; ascis cylindraceo-clavatis, 50-60 \approx 6-7,5; sporidiis distichis, fusoideo-elongatis, simplicibus, saepe guttulatis, leniter curvulis v. subrectis, 10-16 \approx 3.

Hab. in foliis caulisque *Potentillæ argenteæ*, *pulchellæ*, *norvegicæ*, *emarginatæ* in Germania, Fennia, Spetsbergia et Amer. bor. — Sporidia in exempl. amer. 20 μ . long. Sec. Rehm Ascom. n. 759, paraphyses clavulatae ascos superantes; sporidia 12-15 \approx 3; asci 45-50 \approx 8-9.

16. **Mollisia montiæcola** (Berk.) Sacc. *Peziza montiæcola* Berk. Fl. 1348 New Zeal. p. 201. — Atra, expansa, margine angusto; ascis clavatis amplis; sporidiis subcymbæformibus endochromate partito.

Hab. in *Montia fontana* putri, Ahuriri, Eastern Coast Novæ Zelandiæ (COLENZO). — *Pyrenopezizæ atratæ* affinis; an ergo hujus generis?

17. **Mollisia microstigmæ** Passer. Diagn. Fungh. nuov. pag. 258. — 1349 Sparsa vel gregaria, punctiformis, mollis, fusca, cupula madore aperta, disco concavo pallidiore, margine tumido integro, vix extus rugulosa; ascis clavato-fusiformibus, 8-sporis; sporidiis fusiformi-lanceolatis integris; paraphysibus ascorum longitudine.

Hab. in caulis siccis *Trifolii pratensis*, Vigheffio pr. Parma Ital. bor.

18. **Mollisia Mercurialis** (Fuck.) Sacc. *Nip. Mercurialis* Fuck. Symb. 1350
mycol. pag. 293. — Cupulis sessilibus, 1 mm. latis, primo clausis
globosisque, dein apertis, subplanis, extus atris, squamulosis, mar-
gine connivente, albido, fimbriato, disco griseo; ascis oblongo-cla-
vatis, 8-sporis, $44 \approx 4$; sporidiis oblique monostichis, subcylindra-
ceis, rectis, hyalinis, $10 \approx 2$.

Hab. in caulis putridis *Mercurialis perennis* in Rhenogovia et Britannia.

19. **Mollisia melanogramma** (Boud.) Sacc. *Spilopodia melanogramma* 1351
Boud. Disc. charn. p. 32. — Ascomatibus minutis 0,25-0,50 mm.
diam., nigris, margine irregulari, fimbriato macula nigra elongata,
lineari insidentibus, extus rugosis, intus cinereis v. olivaceis; ascis
magnis, cylindraceis, basi vix attenuatis, octosporis, $80-82 \approx 9-10$,
paraphysatis; paraphysibus indistincte septatis, fasciculatis; spori-
diis oblongis, granulosis, $10-12 \approx 3-4$, hyalinis.

Hab. in caulis senescentibus emortuis *Mercurialis perennis* in Gallia. — An *Pyrenopeziza*?

20. **Mollisia rubidula** Rehm Ascom. Alp. n. 23. — Minuta, sessilis, 1352
plerumque oblonga, rarius rotundata, fusca v. fusco-nigricans, hu-
mida explanata, dilacerato-marginata, margine fusca, disco rubi-
dulo, c. 0,2 mm. diam., gregaria contextu parenchymatico, fusi-
dulo; ascis clavatis, subcurvatis, 8-sporis, $70 \approx 10$; sporidiis obtuse-
ellipticis, 1-cellularibus, hyalinis, distichis, $12-14 \approx 6$; paraphysibus
filiformibus, subcurvatis, apice haud incrassatis.

Hab. ad caulem siccum Campanulæ?, Mittagskogel pr. Mit-
telberg in valle Piz alpium tyrolens. alt. c. 2000 m. (REHM).

21. **Mollisia plicata** (Rehm) Sacc. *Niptera plicata* Rehm Ascom. n. 1353
570. — Solitaria vel gregaria, sessilis, primitus sphæroidea, sub-
clausa, dein urceolata, sicca plicata, fusco-cervina, margine albido
connivente, humectata concaviuscula, margine tenui albido subdi-
lacerato cincta, orbicularis; disco livido-albo, contextu parenchyma-
tico fuscidulo, 1 mm. diam.; ascis clavatis, apice incrassatis, 8-
sporis, $45-60 \approx 6-9$, jodo cœrulecentibus; sporidiis ellipticis, plus
minusve obtusis, rectis vel subincurvatis, distichis, 1-cellularibus,
hyalinis, $9-12 \approx 2-3$; paraphysibus obtuse lanceolatis, plus minusve
ascos superantibus, 4-5 μ . crass., hyalinis, in fungo evoluto vix
rite cognoscendis.

Hab. ad caules putrescentes Aconiti Napelli et Angelicae in M. Rigii Helvetiae, in alpinis Tyroliae, Allgoviae et in al-
pibus transylvanicis. — Forma **albo-marginata** Rehm in Hedwigia

1885, p. 231: ascomatibus plerumque gregariis, saepe confluentibus, distincte albo-marginatis, majoribus quam in typo saepe elongatis, parenchymatico-fusco-contextis, fibris marginalibus distantibus septatis, c. 30-40 = 3, subhyalinis, creberrimis. In caulis Aconiti in Hungaria.

22. **Mollisia ? Alismatis** (Pers.) Sacc. *Peziza Alismatis* Pers. M. E. 1354 I, p. 301, *Patellaria* Fr. Syst. Myc. II, pag. 161. — Carnoso-tremellosa, sessilis, minuta, glabra, ex cinereo purpurascens, cupulis depressis, margine subcomplanatis.

Hab. in caulis siccis Alismatis Plantaginis emortuis in aqua adhuc persistentibus, vere in Gallia. — Cupulæ discus passim pulvere rubello tectus.

23. **Mollisia ? Browniana** (Blox.) Sacc. *Peziza Browniana* Blox. in B. 1355 et Br. Ann. N. H. 1865, n. 1072. — Cupulis hemisphæricis, sessilibus, colore corneo; margine pallidiori, ciliato; disco pallido; sporidiis breviter fusiformibus, hyalinis, 11 μ . long.

Hab. in caulis emortuis Epilobii hirsuti Twycross. Britanniæ. — *M. lacustri* proxima dicitur sed margo pilis flexuosis delicatis ciliatus. «An *Trichopeziza*?»

II. Foliicolæ (non graminicolæ).

24. **Mollisia Pumilionis** Rehm Ascom. Alp. n. 18. — Gregaria, sessilis, 1356 substipitata, sphæroidea, demum concavula, sicca fusco-cervina, humida disco fuscidulo, contextu parenchymatico, fusco, c. 0,8 mm. diam.; ascis clavatis, apice incrassatis, 8-sporis, 75 = 9, apice jodo violascente; sporidiis obtuse ellipticis, rectis v. subcurvatis, 1-cellularibus, hyalinis, 12-14 = 3; paraphysibus filiformibus, laxis apice haud incrassatis, hyalinis.

Hab. ad acus Pini pumilionis putridos, Haspelmoos prope Augustam Vindelicorum seu Augsburg (BRITZELMAYR). — «An *Pyrenopeziza*?»

25. **Mollisia Pinastri** (C. et P.) Sacc. *Peziza Pinastri* C. et P. Disc. U. S. I, p. 297. — Sessilis, molliuscula, sparsa, initio pruinâ albida evanidâ conspersa, mox nuda, griseo-brunnea, hemisphærica dein plana, margine saepe lacerato; disco pallide griseo; ascis tereti-clavatis; sporidiis anguste fusoideis, obtusis, 12-15 μ . longis.

Hab. in foliis Pini rigidæ New York Amer. bor. (PECK).

26. **Mollisia minutissima** Karst. Symb. Fenn. XXII, p. 15. — Sparsa 1358 v. subgregaria, sessilis, cupula plana v. concaviuscula, nuda, pale-

scens, sicca fusco-pallida, disco dilutiore, circiter 0,1 mm. lata; ascis clavatis, 36-40 \approx 6-7; sporidiis 7-8 \approx 1-2.

Hab. in foliis putrescentibus *Comari palustris* in regione mustialensi juxta lacum Salois (K. STARBAECK).

27. **Mollisia Earliana** (Ell. et Ever.) Sacc. *Peziza Earliana* Ell. et 1359 Ever. Torr. Club. 1884, p. 74. — Erumpens, sparsa, hepatica, sicca obscurior, epidermide arcta cincta, dein expansa, disco pallido, margine vix eroso; ascis sessilibus oblongis, 50-55 \approx 12; sporidiis tereti-oblongis, obtusiusculis, 3-4-nucleatis, hyalinis, 15-18 \approx 2-3; paraphysibus clavulatis, obtusis.

Hab. in pag. inf. foliorum emortuorum *Fragariæ*, Anna Illinois Amer. bor. (EARLE). — Ascoma 1,2-1,7 mm. lat.

28. **Mollisia Cotoneasteris** Starbäck in Bot. Centralblatt 1888, I, p. 1360 351. — Superficialis v. basi innata, sparsa vel subgregaria, sphæroidea et clausa, demum ore orbiculari v. perlato aperta, cinereo-fusca, sicca nigricans, disco pallidore, 0,3-0,5 mm. lat.; ascis cylindraceo-clavatis, 55-64 \approx 8-10; sporidiis octonis, ellipsoideis vel elongato-ellipsoideis, monostichis vel superne distichis, eguttulatis, 10-15 \approx 5-6; paraphysibus numerosis, 2-2,5 μ . cr. apice subclavato.

Hab. ad folia arida *Cotoneasteris vulgaris* in regione upsaliensi Sueciæ. — «An Pyrenopeziza?»

29. **Mollisia Trollii** (Wettst.) Sacc. *Micropeziza Trollii* Wettst. Fung. 1361 nov. Austr. I, p. 14, t. II, f. 16-19. — Gregaria v. solitaria, carnosula, orbicularis vel elliptica, 0,5-1 mm. diam., initio velata, mox erumpens, sessilis, plana v. subconcava, luteola, demum fuscescens, margine obscuriori, crenulato; ascis oblongo-clavatis, 60-70 \approx 10-14; sporidiis cylindraceis, curvulis, obtusis, 14-20 \approx 5-6, sœpe 2-guttulatis.

Hab. in foliis petiolisque putridis *Trollii europæi* ad pedem montis Hochswab Austriæ. — Sec. Rehm non differt a *Nævia rubella*.

30. **Mollisia singularis** Peck 25 Rep. St. Mus. pag. 142. — Minuta, 1362 ceracea, sessilis, applanata v. convexa, vix marginata, macula indefinita pallida insidens, sordide grisea v. pallida, succineo-brunnea; ascis tereti-clavatis, 50-70 \approx 7-10; sporidiis confertis vel distichis, oblongis, subinde uno apice acutioribus, 10-15 \approx 4-5; paraphysibus filiformibus.

Hab. in pagina inf. foliorum viventium *Ranunculi hispidi* East Berne Amer. bor. — «An forma *Pseudopezizæ Ranunculi*, monente ipso auctore?»

31. **Mollisia islebiensis** (J. Kunze) Sacc. *Pyrenopeziza islebiensis* J. 1363
 Kunze Fung. sel. exs. n. 64 (absque diagnosi). — Ascomatibus superficialibus, patellatis, sessilibus, vix $\frac{1}{2}$ mm. lat., plumbeis, dein nigricantibus; ascis....; sporidiis ovoides, inæquilateralibus, hyalinis, 9-10 \times 4.
Hab. in foliis aridis *Helianthemum Chamæcisti* in Saxonia.
32. **Mollisia Rehmii** Sacc. *Mollisia fuscidula* Rehm Ascom. Alp. n. 1364
 17, nec Sacc. — Gregaria, minuta, sessilis, c. 0,2 diam., orbicularis, humida marginata, patellaris, sicca corrugata, fusca, disco vix dilutiore, contextu parenchymatico; ascis clavatis, apice incrassatis, 8-sporis, 45 \times 8, apice jodo cœrulecente; sporidiis fusoideo-ellipticis, 1-cellularibus, rectis, hyalinis, distichis, 8 \times 2; paraphysibus apice subincrassatis et fuscidulis.
Hab. ad paginam infer. foliorum putridorum *Vaccinii uliginosi* prope glaciem æternam, Taschach-Gletscher in valle Piz alpium tyrolens., c. 2000 m. (DR. REHM).
33. **Mollisia Uredo** Rehm Ascom. Alp. n. 22. — Gregaria in acervulis 1365
Uredinis instar, patellæformis, submarginata vel concava in statu sicco, humida convexula, immarginata, sicca rufo-fusca v. fuscidula disco parum dilutiore, humida flavo-fuscidula, sessilis, c. 0,3 mm. diam., contextu parenchymatico fuscidulo; ascis clavatis, apice incrassatis et obtuse acuminatis, 8-sporis, 50-60 \times 8; sporidiis haud plane evolutis visis, fusoideo-elongatis, 1-cellularibus (?), nucleis 3 præditis, parallelis, hyalinis, 16-18 \times 1-5; paraphysibus ascos superantibus, subramosis, subincurvatis, apice non incrassatis.
Hab. ad paginam inf. foliorum putridorum *Vaccinii uliginosi*, Taschach-Gletscher in valle Piz alpium tyrolensium, alt. c. 2000 m. (DR. REHM).
34. **Mollisia nemophila** Sacc. Fung. Ven. Ser. IV, n. 75, Fung. ital. f. 1366
 1382 (*Niptera*). — Cupulis amphigenis, præcipue nervicolis, sparsis, scutellaribus, minutis $\frac{1}{5}$ - $\frac{1}{7}$ mm. diaen., totis fulvo-ferrugineis, sessilibus, minute rugulosis, basi hyphis brevibus parcis articulatis fuligineis cinctis; excipuli contextu laxe parenchymatico, fuligineo; ascis oblongo-clavatis, subsessilibus, 55 \times 8, 8-sporis pseudopara-physibus crasse filiformibus, obvallatis; sporidiis distichis, cylindraceis v. subfusoideis, 14 \times 3- $\frac{1}{2}$, utrinque rotundatis, 4-guttulatis, rarius 2-nucleatis simulateque 1-septatis, hyalinis.
Hab. in foliis putrescentibus *Carlinæ vulgaris* in silva Montello (Treviso) Ital. bor.
35. **Mollisia parasitica** (Wint.) Sacc. *Niptera parasitica* Wint. Hedw. 1367

1884, pag. 173. — Cupulis in macula irregulari, plerumque angulo-rotundata, 2-6 mm. lata, in foliorum pagina inferiori pallide lutea, in pagina superiori arescendo albida, fusco-cincta et zona luteola indeterminata circumdata, hypophyllis, gregariis, sessilibus, disciformibus, primo submarginatis, demum immarginatis, convexiusculis, subrepandis, ceraceis, luteo-fuscis, disco pallidiore siccis, irregularibus, saepe umbilicatis undulatisque, obscurioribus, fusco-atris, usque $\frac{1}{2}$ mm. latis; ascis clavatis, in pedicellum sublongum attenuatis, 8-sporis, 75-88 \approx 16; sporidiis inordinatis, ovato-ellipticis, parum inaequilateralibus, hyalinis, 14 \approx 7; paraphysibus filiformibus, ascos æquantibus.

Hab. in foliis vivis Melastomaceæ cujusdam prope S. Francisco prov. S. Catharina Brasiliæ.

36. **Mollisia nervicola** (Desm.) Gill. Disc. p. 128, *Peziza nervicola* 1368 Desm. 8 Not. p. 5. — Amphigena, subnervisequa, sessilis, ceraceo-mollis, minuta, glabra, griseo-brunnea, primo subglobosa, ore albicante, adulta plana, margine sub prominente denticulato, disco albidio pallescente; ascis clavatis, minutis, 50 μ . longis; sporidiis oblongis, linearibus, 10 μ . longis.

Hab. in foliis exsiccatis Quercus et Fagi in Gallia bor.

37. **Mollisia prinicola** E. et E. Journ. Myc. 1888, pag. 99. — Hypophylla, subgregaria, carnosso-membranacea, hepatico-brunnea, 0,5 mm. diam., fibroso-contexta, puncto centrali affixa; ascis teretio-oblängis, sessilibus, 35 \approx 6; sporidiis distichis, clavato-fusoideis, continuis, hyalinis, 8-11 \approx 1,5; paraphysibus filiformi-clavulatis.

Hab. in foliis emortuis Quercus Prini, Louisiana Amer. bor. (LANGLOIS).

38. **Mollisia olivacea** (Speg.) Sacc. *Pyrenopeziza olivacea* Speg. F. 1370 Argent. III, n. 80. — Applanato-patellaris, minuta $\frac{1}{4}$ - $\frac{1}{5}$ mm. diam. sicca non vel vix rugulosa, inferne fuligineo-atra, superne (disco) olivascens, tenacella, matrici arcte adnata, sessilis; ascis cylindraceo-clavatis, sursum attenuato-rotundatis, deorsum attenuato-pedicellatis, 50 \approx 5-6, apice jodo cœrulecente; paraphysibus filiformibus apice coalescentibus; sporidiis cylindraceo-ellipticis, inaequilateralibus, utrinque plus minusve acuminato-rotundatis ac 1-guttulatis, hyalinis, 12 \approx 3.

Hab. in foliis caulinibusque dejectis Eryngii agavifolii in paludosis à la Boca del Riachuelo Argentinæ.

39. **Mollisia? Saniculæ** (Wallr.) Sacc. *Phlyctidium Saniculae* Wallr. 1371 Fl. crypt. n. 2328. — Ascomatibus hypophyllis, primum insculptis

fuscescentibus, dein adnatis orbiculatis convexiusculis, nigris opacis, confertissimis, minutis, maculam nigro-fuscum mentientibus, demum deplanatis, verticem irregulariter reclinantibus discumque planum flaventem, margine scisso nigro cinctum recludentibus.

Hab. ad Saniculæ europææ foliorum vegetorum faciem inferior. in Thuringia.

40. **Mollisia Iridis** (Rehm) Sacc. *Micropeziza Iridis* Rehm Ascom. 1372
Lojk. n. 161. — Sicca oblonga v. rotundata, plus minusve hemisphærice contracta, humida explanata v. concava, sessilis, gregaria, subfusca, margine viridi-flavo cincta, disco fuscidulo, contextu parenchymatico fusco, c. 0,2 mm. diam.; ascis clavatis, apice incrassatis, 4-8-sporis, 50 \approx 9, apice jodo cœrulecente; sporidiis elliptico-elongatis, haud acuminatis, rectis, 1-cellularibus, nucleis magnis 1-4 præditis, hyalinis, distichis, 18-21 \approx 4; paraphysibus filiformibus, hyalinis, superne sensim 3,5 μ . incrassatis.

Hab. ad folia et caules putridos Iridis spuriæ prope Tököl Hungariæ (LOJKA).

41. **Mollisia olivaceo-lutea** Berk. North Am. Fungi n. 746 et Cooke 1373
Disc. U. S. I, pag. 297. — Minuta, erumpens, cito decidua, extus olivacea, margine albido; disco concavo, pallide luteo.

Hab. in foliis emortuis in Carolina inf. Amer. bor.

III. Strobilicolæ.

42. **Mollisia fallax** (Desm.) Gill. Disc. p. 119, Phill. Disc. p. 174, *Peziza fallax* Desm. II Not. p. 367. — Suberumpens, minuta, gregaria v. sparsa, sessilis, glabra, extus brunneo-fusca, junior plano-scutellata, dein convexa, flexuoso-sublobata, sicca concava, disco humido griseo, sicco ochroleuco vel brunneo-lutescente; ascis clavatis, 50 μ . long.; sporidiis oblongis, rectis v. subcurvatis, 10 \approx 2.

Hab. in ramis exsiccatis Pin i silvestris, sæpius vero in ejusdem strobilis, in Gallia, Britannia. — Ascoma $\frac{1}{2}$ -1 mm. lat.

43. **Mollisia suecica** Starbäck in Bot. Centr. I, 1888, p. 350. — Superficialis, gregaria, primo sphæroidea et clausa deinde concava, sicca depressa, margine connivente, atra, disco pallidiore, 0,1-0,4 mm. lat.; ascis cylindraceis vel cylindraceo-clavatis, 44-51 \approx 5-9; sporidiis octonis, elongatis vel fusideo-elongatis, biguttulatis vel multiguttulatis v. interdum eguttulatis, distichis-subdistichis, interdum in parte inferiore monostichis, 6-9 \approx 2,5-3,5; paraphysibus numerosis multiguttulatis, apice incrassatis, 2-2,5 μ .

Hab. ad squamas conorum Pin i silvestris in insula Sandon Sueciæ, sat frequens. — «An *Pyrenopeziza*?»

44. **Mollisia rubicunda** Rehm Ascom. Alp. n. 19. — Sparsa, sessilis, 1376 orbicularis, patellaris plana, margine spurio, sicco tumidulo cineta, rubicunda, circ. 1 mm. diam., contextu parenchymatico fuscidulo; ascis clavatis, 8-sporis, $60 \approx 7$; sporidiis obtuse ellipticis, 1-cellularibus, hyalinis, mono- v. subdistichis, $8 \approx 4,5$; paraphysibus filiformibus, ascis superantibus, clavula elliptica, fuscidula.

Hab. ad conos putridos Abietis excelsæ in silva prope Birburg pr. Augustam Vindelicorum (BRITZELMAYR).

IV. Ramicolæ.

45. **Mollisia myocoron** Speg. Fung. Fueg. n. 341. — Cupulis dense 1377 sparsis, applanato-expansis, discoideis, 9,5-1 mm. diam. tenui-membranaceis, sessilibus late arcteque matrici adnatis, glabris, levibus, contextu majuscule parenchymatico olivaceo-fusco; ascis cylindraceo-clavulatis, antice subtruncato-rotundatis crassiuscule tunicatis, postice breviter crasseque attenuato-pedicellatis, $40 \approx 3-3,5$, octosporis; paraphysibus filiformibus obvallatis; jodi ope nulla; sporidiis recte distichis, obovatis vel elongato-ellipticis superne obtuse rotundatis, inferne acutiusculis, $5-7 \approx 1-1,5$, rectis v. inaequilateribus, hyalinis, continuis.

Hab. in ramis fagineis decorticatis ad tuguria incolarum in Gertrude Cove Fuegiae.

46. **Mollisia fallens** Karst. M. F. I, p. 191, *Trochila fallens* Karst. 1378 Symb. p. 217 et 244. — Gregaria, subinde confluens, subimmarginata, sessilis, glabra, plana v. convexiuscula, sicca varie contracta et concava v. concaviuscula, grisea, vulgo basi subfuscescens, sicca fuscescens, margine saepè albidiori, disco griseo, sicco cinerescente, 0,4-0,6 mm. lat.; ascis cylindraceo-clavatis, $50-60 \approx 7-9$, apice jodo vix tinctis; sporidiis distichis, oblongatis, rectis vulgo guttulatis, $9-14 \approx 3,5-4,5$.

Hab. supra corticem ramorum dejectorum Salicis in Mustiala Fenniae. — Ab affini *Mollisia cinerea* cupulis minoribus tenuioribusque, sporidiis majoribus aliisque notis differt.

47. **Mollisia ramealis** Karst. M. F. I, p. 187. — Gregaria, sessilis, 1379 convexa aut plana, subtus fusca v. fuscescens, disco in statu vegeto albido vel albo, siccitate ochraceo, orbicularis aut interdum subirregularis, circiter 1 mm. lat.; ascis anguste clavatis, apice attenuatis, $75-90 \approx 7-8$, obturaculo minutissimo, jodo dilute cœru-

lescente; sporidiis distichis, fusoideo-bacillaribus, rectis v. leniter curvulis, guttulatis vel spurie tenuiter uniseptatis aut eguttulatis, chlorino-hyalinis, 14-30 \approx 2-3; paraphysibus parcis, sursum levissime incrassatulis.

Hab. supra cortices ramulorum Betulæ exsiccatorum in Fen-nia meridionali.

48. **Mollisia stictella** S. et S. Mich. I, p. 425, F. ital. f. 1387 (Ni-
ptera). — Ascomatibus gregariis, initio peridermio velatis dein erumpentibus, denique prorsus superficialibus, scutellatis, undulatis, sessilibus, ceraceis, siccis contortis, disco concaviusculo, sordide lu-tescente; excipuli contextu parenchymatico ad marginem prosen-chymatico, fuligineo; ascis cylindraceo-clavatis, brevissime nodu-loso-stipitatis, 55-65 \approx 6-7; paraphysibus bacillaribus obvallatis. octosporis; sporidiis oblique monostichis vel distichis, oblongo-cy-lindraceis clavulatisve, 8-14 \approx 2-3, granulosis hyalinis.

Hab. in ramulis Alni incanæ emortuis, Pieve di Soligo, Ital. bor. — Affinis *Mollisiae cinereæ*, a qua imprimis ascomatibus ini-tio tectis statim dignoscitur, et Stictæ accedit.

49. **Mollisia benesuada** (Tul.) Phill. Disc. p. 174, *Peziza benesuada* 1381
Tul. Am. Sc. Nat. 1853, t. 15, f. 8-9, *Niptera benesuada* (Tul.) Rehm Ascom. n. 756. — Erumpens, sessilis, gregaria v. subfasci-
culata, subhemisphærica, dein expansa, demum convexa, difformis,
ceracea, extus glabra, aquose cinerea, disco pallido vel brunneo;
ascis tereti-clavatis, 45 \approx 6; sporidiis fusoideis v. oblongo-fusoideis,
hyalinis, 9-10 \approx 2-2,5, distichis; paraphysibus filiformibus, 2 μ . cr.

Hab. in ramis emortuis Alni in Gallia, Britannia et Germa-nia. — Sec. Phillips l. c. sporulæ seu spermatia raro occurrunt.
Species *M. cinereæ* et *fallaci* peraffinis.

50. **Mollisia aquosa** (B. et Br.) Phill. Disc. p. 172, *Peziza aquosa* 1382
B. et Br. Ann. Nat. Hist. n. 1326, t. 20, fig. 19. — Initio clausa,
dein expansa v. concaviuscula, viridi-brunnea, levis; disco aquose-
griseo; ascis elongatis, 50 μ . longis; sporidiis distichis, ovoideo-api-
culatis, 5-6 \approx 2,5-3,5, jodi ope aurantiis.

Hab. ad ramos salicinos ad Lasiosphæriam hirsutam
in Britannia. — Cupulæ $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. diam.

51. **Mollisia tyrolensis** Sacc. *Micropeziza fuscidula* Rehm Ascom. 1383
Alp. n. 26. — Gregaria, sessilis, sicca plerumque longitudinaliter
complicata, fusco-nigra, humida explanata, orbicularis, disco rubro-
fusco, margine sublacerato, obscuriore, tenui cincta, contextu pros-
enchymatico (?), c. 0,4 mm. diam.; ascis clavatis, 8-sporis, 45-

50 = 9; sporidiis oblongis, obtusis, subcurvatis, hyalinis interdum muco involutis, 1-cellularibus, distichis, 9 = 3; paraphysibus filiformibus, clava piriformi, 4-5 μ . crassa, plerumque fuscidula instructis.

Hab. ad ramulos sicclos, decorticatos Rhododendri ferruginei prope Kühtai (Oetz) alpium tyrolensis alt. c. 2000 m. (DR. REHM).

52. **Mollisia rhododendricola** Rehm Ascom. Alp. n. 24. — Gregaria, 1384 erumpens, demum sessilis, plana, ceracea, sicca subcomplicata vel plana, humida explanata, fusco-rubra, margine spurio vix dilutiore, extus fuscidula, contextu parenchymatico, fusco, c. 1 mm. diam.; ascis clavatis, subcurvatis, 8-sporis, 90 = 12, apice jodo cœrulecente; sporidiis ellipticis, nucleis magnis 2 præditis (an demum 2 cellularibus?), hyalinis, distichis, 16-18 = 6; paraphysibus filiformibus, superne sensim 3 μ . crassis et flavidulis.

Hab. ad ramum siccum Rhododendri ferruginei prope Kühtai (Oetz) alpium tyrolensis, alt. c. 2000 μ . (DR. REHM).

53. **Mollisia miltophthalma** B. et C. North Am. Fung. n. 741 et Cooke 1385 Disc. U. S. I, p. 296 (ex errore *multophthalma*). — Minuta hemisphærica extus nigra intus cinnabarina; ascis linearibus paraphysibusque flexuosis; sporidiis oblongis minutis hyalinis.

Hab. ad ramos Corni floridæ in Carolina inf. Amer. bor.

54. **Mollisia discolor** (Mont.) Phill. Disc. p. 175, *Patellaria discolor* 1386 M. et Fries, Notice etc., in Ann., 2, V, pag. 290 et Syll. Crypt. n. 644. — Ceracea, suberumpens, gregaria, sessilis, plano-patellata, extus umbrino-fusca; disco cerino-lutescente; ascis clavatis; sporidiis fusiformibus, hyalinis, 4-guttulatis, 8 = 2 (Phill.).

Hab. ad ramos Corni sanguineæ ad ligna Sedan Galliæ et in Britannia, nec non ad caules Polygoni pensylvanicæ in Carolina sup. — Sec. Karst., sporidia 10-15 = 2-3, vulgo (spurie?) 1-septata. — ***M. chilensis** Nyl. Pez. pag. 79 nota, *Pat. discolor* Mont. l. c. (quoad stirpem chilensem): sporidiis multo majoribus, 3-septatis (spurie?). Ad ramos, Valdivia in Chile.

55. **Mollisia subcornea** Phill. et Harkn. Calif. pag. 2. — Sparsa vel 1387 gregaria, sessilis, patellata, rufo-cornea, margine crasso levi; ascis clavatis; sporidiis distichis oblongo-fusoideis, simulate uniseptatis, 10-12 = 3; paraphysibus filiformibus.

Hab. in ramulis emortuis Eucalypti Globuli in California.

— Ascomate $1\frac{1}{2}$ -1 mm. lat. senectute sæpius coalescentia.

56. **Mollisia viticola** (Pers.) Sacc. *P. viticola* Pers. Myc. Eur. p. 309, 1388

Fr. Syst. Myc. II, pag. 151. — Sessilis, subcoriacea, crassiuscula; cupulis planiusculis sordide nigris, margine tenui cinctis, disco superficiali rufo-carneo.

Hab. in rimis ramorum *Vitis viniferae* in Germania et Gallia. — Subiculum tenue, nigrescens, subpulverulentum. Cupulae subseriatæ, irregulares, *P. atro-virentis*, præsertim colore, haud assimiles.

57. **Mollisia sensitiva** (Hazsl.) Sacc. *Niptera sensitiva* Hazsl. Einig. 1389

Disc. p. 12. — Uda disciformis, adpressa, sicca sphæroideo-contracta, disco flavido, margine nigrescente; ascis cylindraceo-elongatis, sessilibus; sporidiis cylindraceis, $10 \times 2-3$.

Hab. in ramis putridis *Rubi Idæei* pr. Eperies Hungariæ.

58. **Mollisia ribesia** Cooke et Ph. Grevill. IX, pag. 105. — Sparsa, 1390

atra, ceraceo-mollis; cupulis globosis, arcte apertis, dein concavis, hemisphæricis, glaberrimis, 1-4 mm. diam.; ascis clavatis; sporidiis exiguis, hyalinis, 5×1 .

Hab. ad ramos *Ribis rubri* in Arduennis (LIBERT). — «An *Pyrenopeziza?*»

59. **Mollisia umbonata** (Pers.) Sacc. *P. umbonata* Pers. ic. et descr. 1391

fung. p. 35, t. 9, f. 5, Syn. p. 669, Myc. Eur. p. 315, A. S. p. 539, *P. nigro-olivac.* Schwein. l. c. n. 1220. — Sessilis, glabra, albido-cæsia, disco subgelatinoso, ut plurimum umbonato.

Hab. ad ramos putridos et ligna in Germania et Carolina Amer. bor. — Maxime affinis priori, sicca vix distinguenda. In lignis fila observantur nigra, radiatim diffusa, centro cupularum affixa. Cupula 6-10 mm. lata, orbicularis v. lobato-rotundata, margine involuta elevata subcrenulato, extus ceterum glabra. Intus cinereo-nigra, demum tota nigrescit. — Var. *amenticola* cfr. Fuck. Symb. p. 292: ascis elongato-clavatis, stipitatis, 56×4 8-sporis; sporidiis inordinatis, cylindraceis, subcurvatis, 4×2 , hyalinis. In amentis femineis putridis *Alni glutinosæ* in Rhenogovia.

60. **Mollisia cæspiticia** Karst. M. F. I, p. 188. — Plerumque cæspitose aggregata, planiuscula, flexuoso-sublobata, cinereo-pallida, basin versus fuscescens, sicca concava, disco (sicco) centro fuliginoso v. subnigrescente v. ochroleuco, 0,5-1,5 mm. lat.; ascis cylindraceo-clavatis, $30-40 \times 3-4,5$, apice jodo non cœrulecente; sporidiis fusoideo-elongatis, rectis vel leviter curvulis, guttulatis, $4-6 \times 1$; paraphysibus parum notabilibus.

Hab. ad ramos *Salicis*, *Ribis rubri*, *R. nigri* et *Samosciæ*, *Humuli* in Fennia et Gallia. — Pro *Mollisia*

cinerea primo obtutu facile sumitur, sed notis micrologicis et vegetatione admodum differt. Variat sporidiis paullo majoribus, 8 \approx 2. Ad *Pezizellam* vergit.

V. Ligni-corticicolæ.

61. **Mollisia cinerea** (Batsch) Karst. M. F. I, pag. 189, *P. cinerea* 1393

Batsch Cont. I, p. 196, f. 137, Sowerb. t. 64, Nees Syst. f. 269, *P. callosa* Bull. Champ. pag. 252, t. 416, f. 1, Batsch. Cont. II, p. 95, f. 219 (icon non obstat.) Flor. Dan. t. 1490, f. 2, *P. plana*, *P. alnea* et *obconica* Schum. l. c. pag. 417, 423, *P. subcarneopallens* A. S. p. 307, *P. salicaria* Pers. *Nipt. cin.* Fuck. Sacc. F. it. f. 1376. — Sessilis, mollis, minuta, scutellata, cinerea, margine integerrimo albidiore; ascis tereti-clavatis, 40-50 \approx 4-6; sporidiis oblongo-teretiusculis, rectis curvulisve, hyalinis, 6-12 \approx 1-2,5.

Hab. ad ligna, ramos, caules et quascumque quisquiliis putrescentes in Suecia, Germania, Belgio, Gallia, Italia, Britannia, Brasilia, Nova Zelandia, Tasmania, Ceylon, Africa austr., Kamtschatka, America boreali. — Sparsa vel gregaria, carnosomollis, sed non vere tremellosa; junior urceolata, dein dilatata omnino plana, regularis, margine integerrimo. Color variat in humidis fuscus et saepius cinereus, in siccis expallens, albicans. *P. nivea* et *grisea* Batsch El. f. 55 et 56, etiam ad hanc prope accedunt. Plures formas, forte specifice distinguendas notavi in Mich. pl. l. — a. *luteola*: ascomatibus griseis $\frac{3}{4}$ mm. diam., margine undulato, disco amœne cerino; ascis clavulatis, 25-35 \approx 4-5; sporidiis oblongo-cylindraceis, 2-guttulatis, 5-6 \approx 1- $1\frac{1}{4}$. In ramis decorticatis Castaneæ in Italia et Germania. — b. *grisella*: ascomatibus minutis, depresso $\frac{1}{3}$ mm. diam., griseis, disco pallidiore; ascis clavulatis, 40-45 \approx 4; sporidiis cylindraceis, 9-10 \approx 1 $\frac{1}{4}$ -2, 2-guttulatis, rectis. In caulis Eupatoriæ cannabini, Conegliano, Ital. bor. — c. *culmina*: ascomatibus depresso $\frac{1}{2}$ mm. diam., aquose albidis extus fuscidulis; ascis cylindraceo-clavulatis, 35-45 \approx 3 $\frac{1}{2}$ -4; sporidiis cylindraceis, curvulis, 5-6 \approx 1 $\frac{1}{2}$. In culmis putrescentibus Triticorum Selva, Ital. bor. — d. *caricina*: ascomatibus depresso $\frac{1}{3}$ mm. diam. cinereo-albis, extus fuscescentibus; ascis clavato-fusoideis, 55-60 \approx 5; sporidiis fusoideo-cylindraceis, 10-12 \approx 1 $\frac{1}{2}$ -2, 2-nucleatis, curvulis. In foliis Caricis glaucæ Susegana, Ital. bor. — e. *olivascens* Sacc. F. ital. 1377: ascomatibus depresso, undulatis, fuscescentibus, disco atro-olivaceo; ascis cylindraceo-clavatis, 60 \approx 5; sporidiis cylindraceo-fusoideis, 8-10 \approx 2-2 $\frac{1}{2}$, subinæquilateralibus.

In ligno putri quercino in Silva Montello Ital. bor. — f. **macrospora** Sacc. F. it. t. 1378: ascomatibus griseo-fuscis, majusculis; ascis clavatis, $60 \approx 10-12$; paraphysibus quandoque furcatis obvallatis; sporidiis cylindraceis curvatis, $12-14 \approx 2-3$, 2-nucleatis, hyalinis. In ligno putri, Padova. — g. **albo-nigella**: ascomatibus suburceolatis, atris, albo-marginatis; ascis tereti-clavulatis, paraphysatis, $40-50 \approx 7$; sporidiis distichis cylindraceis, curvulis vel rectis, $9-11 \approx 1,7-2$, hyalinis. In caule Atriplicis in Gallia. — h. **minutella**: ascomatibus dense gregariis, suberumpentibus, minutis, applanatis, griseis; ascis $36-40 \approx 5-6$; sporidiis teretiusculis $7-8 \approx 1,5-2$, hyalinis, rectiusculis. In caule Spirææ Ulmariæ in Gallia. — i. **leptospora**: ascis tereti-clavatis, $40-50 \approx 5-6$; sporidiis distichis, bacillari-fusoideis, utrinque acutiusculis, $13-14 \approx 2$, leviter curvis, hyalinis. In caulibus Epilobii in Gallia. — j. **Epilobii** Kunze F. sel. n. 179: ascis $40-45 \approx 5$; sporidiis clavulatis, rectis, $6-7 \approx 2,5$. In caule Epilobii in Germania. — l. **canella** Karst. M. F. I, p. 190: alba, albida vel cinerescenti-albida, $\frac{1}{2}-1$ mm. lat.; ascis $50-67 \approx 6-7$; sporidiis $8-14 \approx 2-3,5$. Ad lignum et corticem Alni et Quercus in Fennia. — m. **revincta** Karst. l. c.: ascomatibus basi ciliis albicantibus adfixis, totis cinereis, $0,5-0,6$ mm. lat.; ascis $40-45 \approx 4$; sporidiis $6-8 \approx 1-1,5$. In caulibus Spirææ in Fennia et Germania. — Buillard notat varietates sequentes: 1) *ardosiaca*, primo subcœrulea, dein nigra; 2) *alba* primo subalbido-cinerea, vetustate plumbeo-cinerea; 3) *viridis* primo viridiuscula, dein dilute cœrulea aut subnigricans.

62. **Mollisia melaleuca** (Fr.) Sacc. *Pez. melaleuca* Fr. Syst. Myc. II, 1394 p. 150, *Niptera melaleuca* (Fr.) Fuck. Symb. p. 292, F. it. f. 1375, *P. leucomela* A. S. p. 342, nec Pers. — Sessilis, ceraceo-sicca, plano-patellata, extus rugulosa nigrescens, disco albicante; ascis oblique stipitatis, linearibus, acuminatis, 8-sporis; sporidiis distichis, cylindraceis, parum curvatis, hyalinis, $8 \approx 2$ (Fuckel).

Hab. ad ligna vetusta decorticata duriora et in ramis putridis Corni in Suecia, Lusatia, Rhenogovia, Gallia, Italia. — Superficialis, nec erumpens, concava, mox explanata, 2 mm. lata, distincte marginata, subtus applanata, rugulosa, atra, vix pulverulenta, disco primo nigricante pallido, mox præcipue centro exalbicante. Margo variat leviter inflexus. Adsunt formæ (distinguendæ?), alia ascis $75-80 \approx 5$, sporidiis $8-10 \approx 1,5-2$ (in ligno quercino et castaneo in Italia bor.), alia ascomate toto nigricante, disco dilutiore, margine pallescente; cellulis marginalibus breviusculis; ascis cylind-

draceis, 80-90 \approx 8; sporidiis fusoideis, 10-12 \approx 1 $\frac{1}{2}$ -2. In ligno putri Galliae. — * **M. echinocola** Sacc. Mich. I, p. 424: ascomatibus adnato-sessilibus, gregariis, $\frac{1}{2}$ - $\frac{1}{3}$ mm. diam., concavis, fuscis, disco margineque pallidioribus; excipuli contextu parenchymatico ad marginem prosenchymatico, fuligineo; ascis cylindraceo-clavatis, 50-60 \approx 7-8, breviter stipitatis, apice rotundatis, paraphysatis, octosporis; sporidiis distichis v. oblique monostichis, 8-10 \approx 2-3, clavulatis v. subfusoideis, brevissime curvis, hyalinis. In spinis capsularum putrescentium Castaneæ vescæ, Susegana Ital. bor. (SPEGAZZINI).

63. **Mollisia Riccia** Sacc. Myc. Ven. Spec. p. 162, t. XVI, f. 3-6. — 1395

Cupulis gregariis sparsisve, sessilibus, applanatis, concaviusculis, ceraceis, 2-3 mm. diam., margine profunde lobato-crenato, disco subsulcato, cinerescenti-luteolo, extus limboque nigricantibus, sublevigatis; ascis dense fasciculatis, tereti-clavulatis, basi attenuatis, 50-55 \approx 4,5-5,5, octosporis; sporidiis inordinate distichis, tereti-oblongis, 8-10 \approx 2,5-3, subcurvatis, hyalinis.

Hab. in ligno putri Robiniae in Ital. bor. et Britannia. — Ob marginem lobatum *Ricciam* in mentem revocat. Sec. Phill. est var. *M. discoloris* sed sat diversa videtur; ceterum *M. discolor* Mont. ut species collecticia censenda est. — Status spermogonius: **Hymenula Riccia** Sacc. cfr. F. Ven. ser. IV, n. 76: pulvillis in ligni superficie et in fungi ascophori consortio, valde extensis, ex albo luteolis, subceraceis; basidiis densissime fasciculatis cylindricis, 40 \approx 1, apice in sterigmata 2 furcatis ibique conidia cylindracea, 3 $\frac{1}{2}$ -4 $\frac{1}{2}$ \approx $\frac{3}{4}$, utrinque obtusa, recta v. curvula hyalina gerentibus. In trunco denudato Robiniae Patavii.

64. **Mollisia cinerella** Sacc. F. ital. f. 1379 et in Bizz. Fl. critt. Ven. 1396

I, p. 334. — Ascomatibus sparsis extus fuscis, disco cinerescente-olivaceo v. cyaneo, cellulis marginalibus excipuli valde elongatis, parenchymaticis, fuligineis; ascis tereti-clavatis, 50-52 \approx 6-6,5 paraphysibusque filiformibus, basidiis brevibus clavulatis suffultis; sporidiis fusoideis, 10-11 \approx 2,5.

Hab. in ligno decorticato Fagi in silva Cansilio Ital. bor. — Ascoma $\frac{3}{4}$ -1 mm. lat.

65. **Mollisia subgilva** Kalch. et Cooke Grev. IX, p. 25. — Sparsa, 1397
sessilis, ceraceo-mollis; cupulis totis subgilvis, $\frac{3}{4}$ -1 $\frac{1}{2}$ mm. diam. concavis; margine rotundato, subtumido; ascis cylindraceis; sporidiis ellipticis, hyalinis, 8 \approx 4; paraphysibus filiformibus.

Hab. ad lignum putre in Africa australi.

66. **Mollisia tasmanica** (Berk) Sacc. *Patellaria tasmanica* Berk Fl. 1398
Tasm. f. 276. — Sessilis; cupulis e concavo planis, hymenio e rufo nigro; ascis clavatis; sporidiis oblongis curvulis, 10-13 μ . longis.
Hab. in ligno emortuo in Tasmania (ARCHER). — Minuta initio subglobosa, concava dein applanata.
67. **Mollisia introviridis** C. et E. Grev. VII, p. 7. — Gregaria; cu- 1399
pulis atro-brunneis, punctiformibus, $\frac{1}{2}$ mm.; disco fuligineo vel subolivaceo; ascis cylindrico-clavatis; sporidiis elongato-ovatis, 7 \approx 3; paraphysibus filiformibus, supra et gelatinæ, viridibus.
Hab. in lignis denudatis, New Jersey Amer. bor.
68. **Mollisia aureo-fulva** Cooke Hedw. 1875, p. 83. — Subgregaria, 1400
ceraceo-mollis, minuta; cupula hemisphærica, dein expansa $\frac{1}{2}$ mm. lata, sicca contracta; disco convexo, aureo-fulvescente, margine extusque brunneo, vix elevato, in statu sicco obscuriore; ascis clavato-cylindraceis; sporidiis ellipticis variabilibus, binucleatis, obtusis, hyalinis, 8-15 \approx 5; paraphysibus superne incrassatis.
Hab. ad ligna vetusta New Jersey Amer. bor. (ELLIS). — «An Pezizella?»
69. **Mollisia? cruenta** (Bong.) Sacc. *P. cruenta* Bong. in Weinm. 1401
Hym. Gaster p. 476. — Gregaria, sessilis, glabra, atra, planiuscula, læsa succum sanguineum fundens.
Hab. ad ligna dejecta humida nuda in Rossia minore (DR. BON-GARD). Adest in herbario Acad. Imperial. scient. Petrop. — Asco-ma applanato-sessile, orbiculare, margine tumidulo, subelevato, vix 1 mm. latum.
70. **Mollisia sanguinolenta** (A. et S.) Sacc. *P. sanguinolenta* A. S. 1402
p. 336, t. VIII, fig. 7, Pers. Myc. Eur. p. 300, Fr. Syst. Myc. II.
p. 145. — Sessilis, glabella, depresso-hemisphærica ochroleuca v. dilute ochracea, ad tactum cruentata.
Hab. ad ligna varia humida in Lusatia. — Sparsa gregariaque, intacta tota unicolor straminea aut pallide ochracea; at læsa sanguine illico emisso maculis persistentibus rubellis pingitur. Latitudo vix 2 mm.
71. **Mollisia complicata** Karst. M. F. I, p. 192. — Gregaria, sessilis 1403
v. substipitata, concaviuscula, cinereo-livida v. pallida, sicca compressa vel marginibus utrinque conniventibus, umbrino-pallens vel nigricanti-fusca, 0,4-1 mm. lat.; ascis cylindraceo-clavatis, 50-60 \approx 5-7, apice jodo non cœruleacente; sporidiis oblongatis vel elongatis, rectis vel leviter curvulis, simplicibus v. spurie tenuiter 1-septatis, 6-12 \approx 2-3; paraphysibus gracilibus, apice incrassatis.

Hab. supra lignum vetustum in Fennia australi.

72. **Mollisia uda** (Pers.) Gill. Disc. p. 127, *P. uda* Pers. syn. p. 634, 140

Fr. Syst. Myc. II, p. 142, *Niptera uda* (Pers.) Fuck. Symb. p. 293.

— Sessilis, ceraceo-aquosa, applanata, immarginata, levis, livido-fuscescens; ascis anguste linearibus, longe stipitatis, 8-sporis, 120 \approx 4; sporidiis elongato-distichis, cylindraceo-fusiformibus, curvatis, continuis, hyalinis, 11 \approx 3 (Fuck.).

Hab. ad ligna prostrata humida in Germania, Gallia, Suecia et Italia. — A *M. cinerea* habitu parum distincta, ad *P. conigenam* etiam accedens. Sparsa, orbicularis, crassiuscula, 2-6 mm. lata, sublivida in olivaceum colorem disco præsertim vergens, subtus demumque fuscescens.

73. **Mollisia decipiens** (De Not.) Sacc. *Trochila decipiens* De Not. 140

Crouam. I, p. 369. — Ascomate parvo patellulato, atro, in secco collabascente subchartaceo; excipulo cellulis parenchymaticis, fuscis, periphericis radiantibus contexto; sporidiis exiguis, ellipsoideis, hyalinis, continuis.

Hab. in palis siccis, Cambiasca Valle Intrasca Ital. bor. — Au Pyrenopeziza?

74. **Mollisia cæsia** (Fuck.) Sacc. *N. cæsia* Fuck. Symb. App. I, p. 335. 140

— Cupulis ascigeris gregariis, 2 mm. lat., sessilibus, planis, demum convexis, extus glabris, griseo-cæsis, margine repando, pallidore, disco concolori; ascis elongatis, 8-sporis, 112 \approx 8; sporidiis in ascii superiore parte stipatis, fusiformibus, biguttulatis, continuis, 16 \approx 3.

Hab. ad lignum putre Fagi prope Estrich Rhenogoviæ. — Fungus conidiophorus in ligni superficie, lineari-effusus, gelatinosus, cæsius; conidiis oblongo-ovatis, continuis, biguttulatis, hyalinis, 4-5 \approx 2.

75. **Mollisia microcarpa** (Fuck.) Sacc. *Nip. (?) microcarpa* Fuck. Symb. 140

mycol. App. I, pag. 334. — Cupulis ascigeris gregariis sparsisve, minutis, 1 mm. diam., sessilibus, diaphano-aquosis, pallide fuscis, extus margineque parum strigosis, primo clausis demum apertis, pallide fuscis, extus margineque parum strigosis, primo clausis demum apertis, concavis, flexuosis, disco concolori; ascis oblongis, sessilibus, 8-sporis, 30 \approx 8; sporidiis distichis, fusiformibus sed utrinque obtusis, continuis, hyalinis, 8 \approx 2 $\frac{1}{2}$; paraphysibus linearibus, simplicibus, multiguttulatis.

Hab. in truncis putridis, madidis, cavis Fagi ad Mittelheimer Äpfelbach Rhenogoviæ. — Stat. conid. sec. Fuck. est *Cladobotryum gelatinosum* Fuck.

3. **Mollisia sublividula** (Nyl.) Karst. M. F. I, p. 193, *Peziza sublividula* Nyl. Obs. p. 61 in not. — Sparsa v. gregaria, explanata, pallido-cinerea v. pallido-albida v. grisea, sicca concava sæpeque anguloso-contracta, obscurior, interdum fere nigricans, 0,5 mm. lat.; ascis fusoideo-clavatis, 32-42 \approx 4,5-5,5; sporidiis fusoideo-oblongatis v. fusoideo-elongatis vulgo guttulatis, 4-7 \approx 1,5; paraphysibus parum notabilibus.

Hab. ad lignum vetustum Pruni et Alni in Mustiala Fenniæ. — *Mollisiæ complicatæ* proxima, sed differt cupulis minoribus, numquam siccitate subplicato-compressis, ascis sporidiisque minoribus etc. Quamquam asci speciminum nostrorum apice jodo non tinguntur, differentia nulla alia inventa, vix dubitare licet, ea ad *Trochilam sublividulam* (Nyl.) pertinere. Ita Karst. l. c.

7. **Mollisia arctata** Karst. Discom. nov. p. 447. — Gregaria, ligno 1409 adnata, obconoidea, ut plurimum distinctor stipitata, disco concavusculo (in statu humido), sicca varie valde contracta et atrata, glabra, circiter 0,3 mm. lat.; ascis clavatis, deorsum in pedicellum pedetentim attenuatis, apice obtusis, 100 \approx 14; sporidiis distichis, elongatis, curvatis, simplicibus, hyalinis, 14-20 \approx 3-4; paraphysibus filiformibus, gracilibus.

Hab. in ligno ramorum Alni incanæ dejectorum ad Mustiala Fenniæ. — Proxime accedit *Pyrenopez. ligni* (Desm.), sed differt cupulis minoribus, ascis sporidiisque majoribus etc. Lignum circa basim ascomatum ut plurimum colore sordide fulvescente vel rufescente tinctum conspicitur.

8. **Mollisia personata** (Karst.) Sacc. *Aleuriella personata* Karst. 1410 M. F. I, p. 186. — Sparsa, sessilis v. (præsertim siccitate) breviter stipitata, fuscescenti-pallida, disco planiusculo, pallido, 0,5-1 mm. lat., circiter 0,5 mm. alt.; ascis cylindraceis, 100-140 \approx 8-10, obturaculo jodo dilute cœrulecente; sporidiis monostichis, ellipsoideis v. oblongato-ellipsoideis, hyalinis, rectis, uniguttulatis, 10-15 \approx 5-7; paraphysibus filiformibus, æqualibus, discretis, 1,5-2 μ . cr.

Hab. supra corticem Betulæ interiorem in Fennia meridionali.

9. **Mollisia perparvula** Karst. M. F. I, p. 191. — Gregaria v. sparsa, 1411 substipitata v. subobconica, albida, deorsum fusca, cupula sicca concavuscula margineque crasso, obtuso, sordide albida, 0,2-0,4 mm. lat.; ascis cylindraceo-clavatis, apice attenuatis, 60-70 \approx 6-8, apice jodo non cœrulecente; sporidiis distichis, oblongatis v. fusoideo-oblongatis, rectis v. curvulis vulgo guttulatis, 7-11 \approx 2-2,5; paraphysibus parum notabilibus.

Hab. supra corticem Tiliæ parvifoliæ et Betulæ in Fen-nia australi.

80. **Mollisia subcorticalis** (Fuck.) Sacc. *Nip. subcorticalis* Fuck. Symb. 1412 mycol. p. 293. — Cupulis gregariis sparsisve, sessilibus, usque ad 2 mm. latis, extus furfuraceis, fuscis, explanatis, margine albido, distincto, disco subconcavo, fusco-rubello; ascis cylindraceis, stipite crasso, brevissimo, subcurvato, 8-sporis, $30 \approx 5$; sporidiis distichis, oblongo-ovatis, subinæquilateralibus, utrinque uniguttulatis, hyalinis, $6 \approx 2-3$.

Hab. in pariete interiore corticis Platani, prope Hattenheim Rhenogoviæ.

81. **Mollisia livido-fusca** (Fr.) Gill. Disc. p. 127, *Niptera livido-fu-sca* (Fr.) Fuck. Symb. App. II, p. 58, *P. livido-fusca* Fr. Syst. Myc. II, p. 147. — Sessilis, membranaceo-mollis, planiuscula, glabra, subtus fusca, supra livido-pallens; ascis subcylindraceis, 8-sporis, $60 \approx 6$; sporidiis elongato-distichis, oblique fusiformibus, continuis, hyalinis, $10-12 \approx 4$; paraphysibus filiformibus, simplicibus, multiguttulatis.

Hab. ad cortices et radices arborum e. g. Carpinii, Betulæ in Suecia, Germania. — Gregaria, majuscula, absque subiculo; ceterum *Tap. fuscæ* proxima. Junior concava, dein plana, constanter integra, sed margo demum non prominens repandus et quasi lobatus; etiam discus albidus subrimosus, ut in *P. rimosa* Sow. Fung. t. 369, f. 12.

82. **Mollisia udula** Mout. et Sacc. (nov. sp.). — Ascomatibus sparsis, 1414 disciformibus, 1 mm. diam., carnosulis, vix marginatis, obscure purpurascenscentibus siccis nigricantibus; ascis clavatis, octosporis, $95 \approx 8$; sporidiis mono-distichis, ellipsoideis, continuis, 2-guttatis, hyalinis, $10-12 \approx 4-5$.

Hab. in ligno abietino mucido pr. Liege Belgii.

83. **Mollisia? viridi-atra** (Fr.) Sacc. *P. viridi-atra* Fr. Obs. I, p. 163, 1415 Syst. Myc. II, p. 142 nec B. et C. — Sessilis, ceraceo-mollis, glabra, globoso-hemisphærica, disco urceolato concolore.

Hab. rarissime ad ligna putrida pinea locis paludosis dejecta in Suecia. — Affinitas dubia ex ipso auct. (Summa Veg. p. 353). Subcarnosa, majuscula, sed mollis, regularis, integerrima, opaca, siccata nigrescens.

84. **Mollisia paullo puncta** C. et E. Grev. VII, p. 7. — Gregaria; eu-pulis atro-brunneis, minimis, $\frac{1}{4}$ mm., demum applanatis; disco ci-nereo; ascis cylindraceis; sporidiis linearibus, $5 \approx 1$.

Hab. in cortice Aceris, New Jersey Amer. bor.

85. **Mollisia glenospora** Ell. et Ever. Torr. Club. 1884, pag. 38. — 1417
 Sessilis orbicularis, $\frac{1}{4}$ - $\frac{1}{2}$ mm. diam., sordide alba dein obscurior; ascis clavato-cylindraceis, 75-80 \approx 10; sporidiis ellipsoideis uno apice acutiusculis uninucleatis, 12-15 \approx 7-8; paraphysibus curvulis clavulatis.
Hab. in trunco putri Magnoliæ, New Field N. J. Amer. bor. — Excipulum g rosse cellulose, apice cellulis erectis, teretiusculis, continuis, 8-10 \approx 2, coronatum.
86. **Mollisia fumigata** Ell. et Everh. Torr. Club. 1884, pag. 41. — 1418
 Dense gregaria, minuta, 1,3-1,6 mm. lat., nigricans, disco subplano, vix marginato, minute papilloso; ascis tereti-oblongis, 50 \approx 7; sporidiis obliquis vel distichis, oblongis, bi- trinucleatis, curvulis, 9-12 \approx 3, hyalinis; paraphysibus clavulatis.
Hab. in ligno putri Magnoliæ, New Field Amer. bor. — Excipulum prosenchymaticum, cellulis apice exerto conidia subglobosa gerentibus.
- VI. Graminicolaæ.
87. **Mollisia Poæ** (Fuck.) Sacc. *Micropeziza Poæ* Fuck. Symb. mycol. 1419 p. 291. — Cupulis subgregariis, subplanis, sessilibus, 1 mm. usque latis, extus margineque crispulo, atro-fusco-rugulosis; disco sordido, demum dilute rubro-fusco; ascis subclavatis, stipitatis, 8-sporis, 54 \approx 8 sporidiis farctis, cylindraceis subcurvatis, multiguttulatis, hyalinis, 12 \approx 2.
Hab. in foliis vaginisque putridis dealbatis Poæ sudeticæ pr. Cestrich.
88. **Mollisia rufula** Sacc. Mich. I, p. 64 (*Micropez.*). — Cupulis superficialibus sparsis, patellari-scutellatis, sessilibus, minutissimis, $\frac{1}{4}$ - $\frac{1}{5}$ mm. diam., margine crassiusculo, levibus, glabris, tenacellis, rufis, extus obscurioribus; ascis cylindraceis deorsum parum attenuatis, 45 \approx 4; paraphysibus filiformibus obvallatis octosporis; sporidiis oblique monostichis v. distichis, cylindraceis, curvulis, 7-9 \approx 1 $\frac{1}{2}$, hyalinis.
Hab. in culmis putrescentibus graminum in silva Montello Ital. bor. — Similis *M. Poæ*, a quibus exiguitate et colore cupulæ non collabentis nec margine crispæ dignoscitur.
89. **Mollisia maculans** Rehm Ascom. n. 155 (sub *Riptera*), cfr. Cooke 1421 in Grevillea IV, p. 41, t. 51, f. 247. — Sessilis, sparsa, primitus urceolata, margine vix albidulo, dein humida explanata, subplano-patellaris, disco albido-cinereo, extus fuscidula, parenchymatico-

contexta, sicca irregulariter rugulosa, plicata, fusco-nigra, c. 0,3 mm. diam.; ascis clavatis, apice subincrassatis, 8-sporis, 30-42 \approx 6-7, jodo apice cœrulescentibus; sporidiis obtuse ellipticis rectis v. subcurvatis, 1-cellularibus, hyalinis, distichis, 9-11 \approx 2-3; paraphysibus filiformibus, apice haud clavatis nec tinctis.

Hab. ad culmos sicclos Nardi strictæ in alpibus editioribus tyrolensibus juxta glaciem æternam. — Culmi, quos incolit, reliquiis v. initiis ascomatum nigro-maculati. An *Pyrenopeziza*?

90. **Mollisia atro-rufa** Sacc. Fung. Alg. Tahit et Gall. n. 17, t. LV, 1422 f. 4. — Ascomatibus sparsis, erumpenti-superficialibus, scutellatis, levibus, $\frac{1}{2}$ mm. diam., atro-rufis, siccis fere nigris, sed contra lucem rubidis; ascis clavatis, longe sensim stipitatis, 50-70 \approx 7, paraphysatis, obturaculo jodii ope cyaneo, octosporis; sporidiis distichis fusoideis, rectis, 12-14 \approx 2-3, plasmate bipartito et tunc spuriæ 1-septatis hyalinis.

Hab. in foliis Coix lacrymæ putridis in Arduennis (LIBERT).

— Affinis *M. rufulæ* Sacc. sed asci et sporidia multo majora.

91. **Mollisia arenivaga** (Desm.) Phill. Disc. p. 186, *Pez. arenivaga* 1423 Desm. 20 Not. pag. 19, Crypt. II, n. 517. — Foliicola erumpens, minuta, sparsa, ceraceo-mollis, glabra, sessilis, junior globosa, adulta plana, extus fulva, margine clavato tumidiusculo, fulvo, sicco nigro, disco aquoso, subalbido, sicco-brunneo; ascis clavatis subcylindricis 75 μ . long. inter paraphyses simplices erectis; sporidiis octonis, oblongo-ovoideis, hyalinis, utrinque obtusis, 15 \approx 5-6.

Hab. in foliis Psammæ arenariæ in Gallia et Britannia.

— Ascoma $\frac{1}{4}$ - $\frac{1}{2}$ mm. lat.

92. **Mollisia cymbispora** Rostr. Fung. Groenl. pag. 538. — Gregaria, 1424 sessilis, concava, rufo-fusca, sicca nigrescens, 0,3-1 mm. lat.; ascis cylindraceo-clavatis, 90-115 \approx 12-13; sporidiis distichis, fusoideo-cymbiformibus spurie 1-septatis, hyalinis, guttulatis, 16-20 \approx 5-8; paraphysibus filiformibus.

Hab. in foliis emortuis Eriophori Scheuchzeri, E. angustifolii, Caricis pullæ in Groenlandia.

93. **Mollisia arundinacea** (DC.) Phill. Disc. p. 177, *Xyloma* DC. Fl. 1425 Fr. VI, pag. 162, *Eustegia* Fr. Elench. II, p. 112, *Stegia* Fuck. Symb. App. I, pag. 328, *Sphaeria Calami* Nees in Myc. Heft II, pag. 63, *Peziza Kneiffii* Wallr. Fl. Crypt. n. 2552. — Gregaria, erumpenti-sessilis, plana, orbicularis, tenuis, ceracea; disco pallido demum fuligineo, margine angusto, atro; ascis tereti-clavatis; sporidiis anguste fusoideis, 10-15 \approx 2; paraphysibus filiformibus.

Hab. in culmis putrescentibus *Phragmitis* in Scania, Gallia, Britannia, Belgio, in culmis *Phalaridis* in Arduennis, ad *Accorum Calatum* in Batavia. — *Ascomia* vix $\frac{1}{2}$ mm. lat., saepius macula atrata insidens. Sec. Phill. l. c., hymenium proiectum ab excipulo sponte separatur, operculi instar. — Forma *Typhæ* Fuck. l. c.: ascis lineare-clavatis, stipitatis, 8-sporis, 78 ± 6 ; sporidiis in ascis superiori parte distichis, fusiformibus, rectis, 1-2-guttulatis, simplicibus, hyalinis, $8 \pm 2\frac{1}{2}$; paraphysibus copiosis, linearibus, utrinque attenuatis, ascis longitudine, 4 μ . cr. 3-4-septatis, indivisis. In calamis *Typhæ angustifoliæ* putridis pr. Budenheim.

94. **Mollisia riparia** Sacc. Mich. II, p. 81, F. ital. f. 1380. — Asco- 1426 matibus gregariis, erumpenti-superficialibus, concavo-scutellatis, sessilibus, siccis compressis $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam. ubique glabris, rufo-brunneis, excepto margine pilosulo albicante; excipuli contextu prosenchymatico, rufo-melleo, cellulis marginalibus liberatis, piliformibus, asperulis; ascis cylindraceo-fusoideis, subsessilibus, $40 \pm 6-6\frac{1}{2}$; paraphysibus fusoideo-bacillaribus, longioribus, melleis ob-vallatis, octosporis; sporidiis distichis, fusoideis v. allantoideis, $8-9 \pm 1\frac{1}{4}-1\frac{1}{2}$, hyalinis.

Hab. in culmis *Phragmitis communis*, Sainter Galliae (BRUNAUD). — *Pyrenop. Karstenii* affinis videtur, differt vero paraphysibus fusoideis melleis, ascomatibus paullo majoribus etc.

95. **Mollisia lacustris** Fuck. Symb. Myc. p. 292 (*Niptera*) nec *P. lacustris* Fries. — Sparsa adnata, glabra, orbicularis, plano-convexa extus nigra, disco nigro; ascis linearibus, 8-sporis, 82 ± 4 ; sporidiis distichis, cylindraceis, rectis, 10 ± 2 .

Hab. in foliis aqua immersis putrescentibus *Typhæ latifoliæ* prope Budenheim Rhenogoviæ.

96. **Mollisia hydrophila** Karst. M. F. I, p. 196, *Nipt. cinerea* v. *donacina* Sacc. Mich. I, p. 425. — Gregaria, tota albida v. albido-pallens, interdum sordide albida v. cinerescens, sicca (extus) colore immutata aut fusca v. fuscescens, convexa, sicca concava, circiter 0,7-1 mm. lat.; ascis fusoideo-clavatis, $50-70 \pm 5-7$, obturaculo jodo obsolete coerulecente; sporidiis aciculati-elongatis, guttulatis, rectis, $9-12 \pm 1,5-2,5$; paraphysibus gracilescentibus, apicem versus incrassatis.

Hab. ad culmos *Phragmitis* emortuos in Fennia, Italia, Gallia, Amer. bor.

97. **Mollisia Typhæ** (Cooke) Sacc. *Peziza Typhæ* Cooke in Grev. I, 1429 p. 131, *Mollisia* Phill. Disc. 188. — Gregaria v. sparsa, minuta,

erumpens, sessilis, atra, molliuscula ceracea, mox plana, sicca lateraliter compressa, disco fuligineo, sicco nigricante, minute papillato, piano; ascis cylindraceis; sporidiis fusoideis, hyalinis, continuis, 12-14 \approx 2,5-3.

Hab. in foliis *Typhæ* King's Lynn Britanniæ (PLOWRIGHT). — *M. epitypha* Karst. vix est diversa, ex ipso Karsten.

98. **Mollisia epitypha** Karst. M. F. I, p. 196. — Gregaria, sessilis, 1430 primitus sphæroidea, mox hemisphærica, demum convexiuscula, primo tota fuliginoso-cinerea, dein cinereo-livida, disco griseo-pallente v. fuliginoso-pallido, sicca concolor v. extus nigricans, 0,6-1 mm. lat.; ascis fusoideo-clavatis, 50-60 \approx 5-6, obturaculo minutissimo jodo obsolete cœrulecente; sporidiis distichis, rectis, aciculari-elongatis, 10-14 \approx 2; paraphysibus gracilescensibus, apice in-crassato.

Hab. ad calamos *Typhæ latifoliæ* putrescentes in Mustiala Fenniæ.

99. **Mollisia palustris** (Rob. et Desm.) Karst. M. F. I, p. 196, *Peziza palustris* Rob. et Desm. XIV Not. p. 187. — Sparsa v. gregaria, erumpens, primitus sphæroidea, tandem aperta, plana vel convexiuscula, applanata, albida vel dein griseo-brunnea, disco griseo-pallente, margine sinuoso, integerrimo, 0,4-0,8 mm. lat.; ascis cylindraceo-clavatis, apice attenuatis, 40-52 \approx 4-5, jodo non cœrulecente; sporidiis fusoideo- v. aciculari-elongatis, guttulatis, rectis, 6-10 \approx 1-2; paraphysibus gracilibus.

Hab. ad calamos et folia *Junci articulati* etc. putrescentia in Fennia, Gallia, Britannia.

100. **Mollisia melatephra** (Lasch) Karst. M. F. II, p. 197, *Pez. me- latephra* Lasch in Rab. Herb. myc. n. 825, non *M. melatephra* Sacc. et Phill. (sporidiis multo brevioribus diversa. Cfr. *M. juncina*), *Pez. conformis* Nyl. Obs. p. 63 (ut videtur). — Gregaria, erumpens aut sessilis, sicca concava nigricans, disco pallido, 0,2-0,4 mm. lat.; ascis fusoideo-clavatis, 50-60 \approx 7-9, obturaculo minutissimo jodo obsolete cœrulecente; sporidiis fusoideo-elongatis v. fusoido-filiformibus s. bacillaribus, guttulis refertis, 12-20 \approx 2; paraphysibus gracilescensibus.

- Hab.* ad *Juncum conglomeratum* in Germania et Belgio. 1433
 101. **Mollisia juncina** Rehm Asc. n. 856, an Pers. Myc. Eur. p. 314? (sub *Peziza*), *M. melatephra* Sacc. Mich. II, pag. 81 et F. ital. f. 1384, Phill. Disc. p. 189, non auct. nec *Pezizella juncina* Fuck. Sh. I, p. 296. — Gregaria, sessilis, subceracea, primitus globosa,

dein urceolata, demum patellaris, sicca irregulariter semiglobosa. flavidula, demum fusco-rubidula v. nigro-fuscidula, contextu parenchymatico, fusco, margine subhyalino-pseudoprosenchymerico, disco pallide flavidulo, 0,3-0,5 mm. diam.; ascis clavatis, apice obtuse subacutatis, 8-sporis, 45-60 \approx 6-8; sporidiis cylindraceis, obtusis, rectis, 1-cellularibus, utroque apice nucleolo parvulo præditis, hyalinis, 9-10 \approx 1,5, distichis; paraphysibus filiformibus teneris apice subramosis et discum dilute flavidulum formantibus.

Hab. ad calamos sicclos Junci effusi, Gfällalpe prope Lofer Salisburgiæ, 1300 m. (NIESSL.), Saintes Galliæ (BRUNAUD), in culmis Poæ et Caricum in Gallia et Britannia. — A *M. melatephra* sporidiis multo brevioribus differt.

102. **Mollisia emergens** Karst. M. F. I, pag. 194. — Gregaria, per 1434 peridermum erumpens, demum sublibera, concava, testaceo-pallida, demum umbrino-pallida, margine lacero v. crenulato-inæquali, subalbicante, disco obscuriore, planiusculo, 0,2 mm. lat.; ascis fusideo-clavatis, 35-45 \approx 6 apice jodo non cœrulecente; sporidiis aciculari-elongatis v. aciculari-filiformibus, sæpe obsolete septatis, 10-20 \approx 1,5-2; paraphysibus cohærentibus, (saltem in statu sicco).

Hab. in calamis Junci conglomerati putrescentibus in Mustiala Fenniæ.

103. **Mollisia simillima** Karst. M. F. I, p. 196. — Gregaria, primitus 1435 sphæroidea, dein appanata, plana v. convexiuscula, sicca concava, lvescente-albida, sicca fuscescente-livida, disco albido, circiter 0,8 mm. lat.; ascis fusoideo-clavatis, 50-60 \approx 5, obturaculo jodo leviter cœrulecente; sporidiis aciculari-elongatis, guttulatis vulgo rectis, 9-14 \approx 1-2; paraphysibus 2-3 μ . cr., apice 4 μ . crass.

Hab. ad calamos Junci conglomerati putrescentes in Fennia australi.

104. **Mollisia junciseda** Karst. M. F. I, p. 198. — Gregaria v. sparsa 1436 plana v. convexiuscula, fuscescens v. pallida, disco pallido-albido, sicca tota nigricans, at tota pallida aut nigricans, disco sordide pallido vel subrufo, circiter 0,4 mm. lat.; ascis fusoideo-clavatis (v. uno apice obtusatis), 60-70 \approx 8-10; sporidiis filiformibus seu bacillaribus, utroque apice attenuatis, guttulis majusculis refertis, rectis v. rectiusculis, 20-28 \approx 2-3; paraphysibus apicem versus incrassatis.

Hab. in culmis Phragmitis in Fennia et calamis Junci in Germania. — ***Mollisia juncinella** Karst. M. F. I, p. 198: typo subsimilis, at sporidiis 23-33 \approx 3-3,5. Ad Juncum articulatum prope Mustiala Fenniæ.

105. **Mollisia paulula** (Rob.) Phill. Disc. p. 186, *Peziza paulula* Rob. 1437 et Desm. 19 Not. p. 29. — Erumpens, sparsa aut gregaria sessilis, ceraceo-mollis, glabra, subglobosa, minutissima, extus umbriuno-olivascens, intus albido-pallens, ore connivente albo integerrimo; ascis brevibus, crassis, subcylindraceis, 40-50 \times 12-15; sporidiis oblongis, obtusis, 12 \times 4, hyalinis 2-guttulatis.

Hab. ad calamos sicclos Junci maritimi in Gallia bor. et Britannia. — Ascoma 150 μ . lat.

106. **Mollisia stictoides** C. et E. Grev. VI, p. 8, t. 96, f. 37. — Ses. 1438 silis, erumpens, sparsa; cupulis minutis, tectis, demum emergentibus, atro-brunneis; disco pallido; ascis clavatis; sporidiis arcte fuscoideis, nucleatis, 16 \times 3.

Hab. in calamis Junci, New Jersey Amer. bor. — Ascoma $\frac{1}{10}$ - $\frac{1}{8}$ mm. diam., marginibus demum longitud. conniventibus.

107. **Mollisia Curreyana** Phill. Disc. pag. 406, *Patellaria palustris* 1439 Curr. Linn. Trans. XXIV, p. 155, t. 25. — Subplana, marginata; disco cyaneo-griseo, subnigricante, circulari vel irregulari, vix 2 mm. lat.; ascis tereti-clavatis, 55 \times 5; sporidiis clavulato-ellipsoideis, curvulis, 8-10 \times 2.

Hab. in calamis Juncorum aqua immersis, Paul's Cray Britanniæ (CURREY).

108. **Mollisia scirpicola** (Fuck.) Sacc. *Micropez. scirpicola* Fuck. 1440 Symb. myc. p. 292. — Gregaria, numerosissima; cupulis sessilibus, minutissimis, punctiformibus, planis excavatisve, extus margineque elevato atro-fuscis, rugulosis, disco fuligineo; ascis sessilibus, oblongis, 8-sporis, 54 \times 8; sporidiis inordinatis, cylindraceis, curvatis, 12-14 \times 3.

Hab. in foliis putrescentibus Scirpi silvatici pr. Mappen Rhenogoviæ.

109. **Mollisia scirpina** (Peck) Sacc. *Peziza scirpina* Peck 28 Rep. St. 1441 Mus. p. 67. — Minuta, sparsa, erumpens, glabra, extus nigra, intus pallidior v. griseola; ascis oblongis; sporidiis confertis, fusoideis, rectis curvulisve, binucleatis, hyalinis 20-25 μ . longis.

Hab. in caulis emortuis Scirpi cæspitosi, M. Adirondack Amer. bor.

110. **Mollisia silvatica** Karst. Symb. Fenn. XXIII, p. 4. — Sparsa v. 1442 subgregaria, superficialis, mollis, patellata v. subconoidea, glaberri- ma, fuscescens, sicca nigrescens margineque elevato, disco planiusculo pallido, parenchymatico-contexta, 0,1-0,2 mm. lata; ascis cla-

vatis sessilibus, 45-50 \approx 10; sporidiis distichis, elongatis, 2-guttulatis, 15 \approx 3; paraphysibus filiformibus.

Hab. in foliis putridis *Scirpi silvatici* pr. Mustiala Fenniæ.

111. **Mollisia puccinioidea** (De Not.) Sacc. *Trochila puccinioidea* De Not. Comm. Critt. I, p. 370. — Hypophylla, discoideo-patellulata, disco fusco-rubiginosa, vix marginato, excipulo tenui, parenchymatico; sporidiis exiguis, oblongatis, hyalinis, continuis.

Hab. in foliis emortuis *Caricis* in Monte Baldo Italiæ borealis.

112. **Mollisia luzulina** Karst. M. F. I, p. 194. — Gregaria, per epidermidem stellatim fissam erumpens, concava, margine lacero, umbrino-pallida, 0,3 mm. lat.; ascis cylindraceo-clavatis, 40-48 \approx 4-5; sporidiis distichis, aciculari-elongatis, simplicibus, 8-15 \approx 1,5-2; paraphysibus haud bene discretis (in statu sicco), gracilibus.

Hab. in foliis *Luzulæ pilosæ* emortuis prope oppidum Wasa Fenniæ.

VII. In Cryptogamis.

113. **Mollisia epibrya** Karst. M. F. I, p. 190. — Demum planiuscula, 1445 subfuliginosa, sicca concava, subhemisphærica, obscurior, circiter 0,7 mm. lat.; ascis cylindraceo-clavatis, 56-65 \approx 7-8, obturaculo jodo cœrulecente; sporidiis ellipsoideo-oblongatis vel oblongatis guttulas 2 apicales foventibus, 7-10 \approx 3-4; paraphysibus filiformibus 1,5 μ . cr.

Hab. supra muscos prope Mustiala Fenniæ (H. A. KULLHEM). — A *Mollisia cinerea* colore, sporidiis et ascis crassioribus recedit. Plantula non satis cognita, nam specimina perpaucia et sicca tantum visa.

114. **Mollisia pteridina** Karst. M. F. I, p. 194. — Subgregaria, sesilis, concava v. concaviuscula, pallida v. pallido-umbrina, margine subcrenulato inæquali, 0,2-0,4 mm. lat.; ascis cylindraceo-clavatis, 35-45 \approx 4-5, obturaculo minuto jodo obsolete cœrulecente; sporidiis elongatis v. aciculari-elongatis, rectis, vulgo guttulatis, 5-10 \approx 1-2.

Hab. ad stipites *Pteridis aquilinæ* vetustos in regione Aboensi Fenniæ.

115. **Mollisia tenella** C. et E. Grev. VII, pag. 40. — Hypophylla, 1447 sparsa; cupulis tenuibus, hemisphæricis, demum applanatis, 0,15-0,2 mm., pallide corneis, margine albidis; ascis clavatis; sporidiis linearibus, rectis 5 μ . longis.

Hab. in frondibus Osmundæ, New Jersey Amer. bor.

116. **Mollisia Lycopodii** Le Bret. et Malbr. Mélang. mycol. p. 5. — 1448

Cupulis sparsis, minutissimis, sessilibus, primo urceolatis, demum undulato-applanatis, ceraceis, micantibus, furfuraceis, fusco-cineris, basi obscurioribus, margine pallidiore v. omnino albo, leviter fimbriato, disco cinereo-fuscescente, siccis nigricantibus; ascis claviformibus, ad basim attenuatis, apice obtusis etiamque truncatis, subsessilibus, obturaculo minuto jodo dilute cœruleo-argentatus, 49-58 \times 4-6; sporidiis distichis interdum parte superiore submonostichis conglobatis, cylindraceis, vulgo unico apice, acicularibus, rectis curvulive, hyalinis, 8-9 \times 1,5-2,5; paraphysibus filiformibus, hyalinis, haud facile conspicuis.

Hab. ad spicas emortuas putrescentesque Lycopodii clavati in ericetis nemorosis, Saint-Jacques prope Rouen, Galliæ. — Punctiformis, *M. cinereæ* affinis.

117. **Mollisia Collematis** Boud. in Bull. Soc. B. Fr. 1881, p. 96, t. III, 1449

f. 7. — Perexigua sessilis, nigricans, margine albo, 0,3-0,5 mm. diam., expansa marginata, extus fibrillis adpressis, parallelis, nigricantibus obsessa, disco plano v. concaviusculo griseo; ascis tereti-clavatis, 70 \times 10; sporidiis ovato-oblongis, subpiriformibus vel medio subcoarctatis, 7-8 \times 3-4; paraphysibus filiformibus, subinde ramulosis, septulatis.

Hab. in Collematis specie ad terram nudam Montmorency Galliæ.

VIII. In corpore humano et in fimo.

118. **Mollisia auriculæ** (Garov.) Sacc. *Peziza auricula* Garov. Arch. 2450

trienn. Lab. critt. I, p. 218, t. XI (nomen). — Ascomatibus planoscutellatis, sessilibus, crassiuscule marginatis, castaneo-rufis, disco concolori, 0,5-0,7 mm. latis; ascis cylindraceo-clavatis, stipitatis, 90-100 \times 8-10; sporidiis oblique monostichis ellipsoideo-oblongis, 8-12 \times 5,5-6; paraphysibus filiformi-clavulatis.

Hab. in cerumine auriculæ hominis in Italia boreali. — Diagnosin ex icone erui.

119. **Mollisia? coprophila** Speg. Fung. Fueg. n. 340. — Cupulis saepius hinc inde densiuscule aggregatis, sessilibus, udis cupulato-explanatis, 0,5-3 mm. diam., siccis contractulis, crassiuscule membranaceis tenacellis rigidulis intus extusque fusco-olivaceis, glaberrimis, margine acutis integris non v. vix repandulis, contextu parenchymatico atro, intus pallidiore; ascis cylindraceis, antice ob-

tuse rotundatis, postice breviter crasseque attenuato-pedicellatis 140-150 \times 8-10; paraphysibus densis, filiformibus hyalinis, apice coalescentibus olivascentibusque obvallatis, octosporis; sporidiis recte v. oblique monostichis, ellipticis, utrinque obtuse rotundatis, 16 \times 8, hyalinis, continuis, levibus non v. 1-guttulatis.

Hab. in fimo equino et vaccino putrescente vulgata pr. Ushuvia Fuegiæ. — Jodi ope tunica ascorum apice dilute cœrulescit.

B. *Belonopsis* Sacc. Ascoma subcirculare, subsuperficiale. Sporidia bacillari-fusoidea, 16-30 μ . long., sæpius simulate septata. — Cfr. genera *Belonium* et *Belonidium*.

120. **Mollisia cyanites** Phill. Disc. p. 176. — Gregaria, minuta, sessilis, concava, dein plana, atra, margine fimbriato, disco cyaneo-griseo; ascis clavatis; sporidiis tereti-fusoideis, elongatis, 5-7-guttulatis, 25 \times 3; paraphysibus filiformibus.

Hab. in caulis herbaceis in Scotia (?) — Ascoma $1\frac{1}{4}$ - $1\frac{1}{2}$ mm. lat.

121. **Mollisia calamicola** Karst. M. F. I. p. 198. — Gregaria, fuscescens, sicca vulgo nigricans, disco pallido vel albido, planiuscula, margine distincto, circiter 0,3 mm. lat.; ascis fusoideo-clavatis, 65-75 \times 5,5-6,5, obturaculo jodo leviter cœrulecente; sporidiis fusoideo-elongatis vel fusoideo-bacillaribus, guttulis fœtis vel spurie tenuiter septatis, 16-24 \times 2-3; paraphysibus gracilescitibus.

Hab. ad calamos Acori Calami semiputres, Runschala Fenniæ.

122. **Mollisia ventosa** Karst. M. F. I. p. 188. — Gregaria, e sub-sphæroideo-applanata, demum convexiuscula, primo tota cinerea, dein cinereo-atra, disco ochraceo-pallido v. tota cinereo-nigricans, margine elevato, sæpissime flexuoso, obtuso, 1,5-2 mm. lat.; ascis cylindraceo-clavatis aut subcylindraceis, 90-120 \times 6, obturaculo jodo leviter cœrulecente; sporidiis fusoideo-elongatis, plerumque curvulis, guttulis 2-4 majusculis præditis v. spurie tenuiter 3-septatis, 10-20 \times 2-3,5; paraphysibus filiformibus 1 μ . cr.

Hab. ad lignum vetustum locis uidis in Fennia australi ad radices Pini silvestris in Lapponia, ad Salicem in Scotia (KRITH) et Waitaki in Nova Zelandia. — Differt a *Mollisia cinerea* mole ascomatum majore, colore et sporidiis majoribus, a *Moll. discolor* cui sporidia vulgo uniseptata, 10-15 \times 2-3, *Moll. fallace* quæ sporidiis simplicibus, 7-9 \times 1,5-2 est instructa.

123. **Mollisia sordidula** Karst. M. F. I. p. 193. — Gregaria, sessilis, convexa, immarginata, sordide pallida, sicca concaviuscula v. concava, margine elevato, umbrino-pallida v. fuscescens, 0,2-0,3 mm.

lat.; ascis cylindraceo-clavatis, apice attenuatis, 65-75 \approx 9-11, obturaculo minutissimo jodo cœrulecente; sporidiis distichis, fusideo- vel aciculari-elongatis vel bacillaribus, guttulis plerumque 4 majusculis fœtis (vel spurie tenuiter septatis), rectis vel leniter curvulis, 14-30, plerumque 18-25 \approx 3-4; paraphysibus flexuosis 1 μ . cr.

Hab. ad lignum ramorum Pini silvestris decorticatorum in Mustiala Fenniæ.

124. **Mollisia graminea** Karst. M. F. I, p. 199. — Gregaria, convexa, 1456 subimmarginata, sessilis, albida v. livescente-albida, basi fuscescens, sicca concava, pallescens, circiter 1 mm. lat.; ascis fusoideo-clavatis, 70-90 \approx 8-10, apice jodo non cœrulecente; sporidiis conglobatis v. subdistichis, bacillaribus, utrinque attenuatis, guttulis majusculis præditis v. (simulate?) 3-septatis, leniter curvulis v. subrectis, 20-30 \approx 3-4; paraphysibus 2 μ . cr. apice 4 μ . cr.

Hab. ad culmos Poæ nemoralis, Calamagrostidum et Baldingeræ siccis locis uidis dejectos, in Fennia australi. — Ascomata basi fibrillis tenerrimis radiatim serpentibus fuscis cincta. Spermogonia ejus forte sistunt socii obvii pulvinuli gelatinosi, flavescentes (in sicco), minuti, 0,2 mm. lat.; sporulis cylindraceis v. fusoideo-elongatis, vulgo curvulis, incoloribus, simplicibus, 8-10 \approx 1,5. — Var. **caricina** Sacc.: ascis teretiusculis, paraphysatis, 90-100 \approx 7; sporidiis bacillaribus, leniter curvis, 15-21 \approx 2-3, pluriguttatis vel spurie septatis hyalinis. In foliis emarcidis Caricis ripariæ in Gallia.

125. **Mollisia atriella** Cooke Hedw. 1875, p. 83, Disc. U. S. I, p. 297. 1457

— Gregaria, sessilis, ceraceo-mollis, minuta, $\frac{1}{5}$ mm., cupula primum hemisphærica, mox applanata, anguste marginata, glabra, atra, disco atrocinereo-nigrescente; ascis clavato-cylindraceis; sporidiis anguste fusiformibus, 30 \approx 2,5, multinucleatis; paraphysibus filiformibus.

Hab. ad culmos Andropogonis, New Jersey Amer. bor. (ELLIS).

126. **Mollisia advena** Karst. Fung. Spetsb. n. 24. — Sparsa v. subgregaria, erumpenti-superficialis, hemisphærica, sicca subsphæroideo-contracta, madefacta fusco-pallida, disco pallescente, sicca fusco-nigricans v. nigricans, margine integro, excipulo cellulis parenchymaticis fuscis contexto, 0,1-0,2 mm. lat.; ascis oblongato- v. elongato-clavatis, 50-70 \approx 12-18, apice jodo non cœrulecente; sporidiis conglobatis, fusoideo-elongatis v. fusoideo-bacillaribus, gut-

tulatis v. demum spurie tenuiter septatis, 20-36 \approx 3-4; paraphysibus filiformibus, 1,5 μ . cr., apicem versus leviter incrassatis.

Hab. in foliis Eriophori angustifolii, Luzulæ hyperboreæ et Luzulæ arcuatæ β . confusæ putrescentibus ad Adventbay et Nordfjorden Spetsbergiæ.

127. **Mollisia excelsior** Karst. M. F. I, pag. 199, *Belonidium* Phill. 1459 Disc. p. 150. — Gregaria, sessilis, convexa, livido- v. cinereo-albida v. pallida, sicca, concava concolor v. nigricans, disco albido v. pallido, 1-1,2 mm. lat.; ascis fusoideo-clavatis, 120-130 \approx 8-10, obturaculo minutissimo jodo leviter cœrulescente; sporidiis conglobatis, filiformibus, utroque apice attenuatis, guttulis majusculis fœtis v. spurie tenuiter pluriseptatis, 42-50 \approx 3-4; paraphysibus 1,5 μ . cr., apice clavato, 4-6 μ . crasso.

Hab. ad culmos Phragmitis communis et Arundinis vetustos in lacubus in Fennia australi et Scotia. — Sporidia sec. Phill. 50-65 \approx 4-5; cellulæ marginales oblongæ, umbrino-brunneæ. An potius *Belonium* v. *Gorgoniceps*?

C. *Hysteropeziza* Rabenh. Ascoma erumpens, siccum elongatum, atrum, hysterioideum.

128. **Mollisia petiolaris** (A. S.) Sacc. *Hysterium petiolare* Alb. et Schw. Consp. p. 59, *Hysteropeziza erumpens* (Grev.) Rab. F. E. n. 1811 c. ic., *Pyrenopeziza* Rehm Asc. n. 354, *Peziza erumpens* Grev. Crypt. f. 99, Gonn. et Rab. t. V, f. 6, *Trochila petiolaris* (A. S.) Rehm Disc. p. 132. — Parallele sparsa, minuta, initio subcutanea dein erumpens, sessilis, flavidо-cinerea, concava, subcircularis, glabra, sicca oblonga in corticis rimis atratis immersa, subimmarginata, integra; ascis clavatis, apice tunica incrassatis, 36-45 \approx 6; sporidiis clavulatis, hyalinis, obtusis, rarius curvulis, 7 \approx 1,5-2; paraphysibus filiformi-clavulatis.

Hab. in petiolis Aceris, Æsculi, Ailanthi, in caulinis Epilobii (?), in Britannia, Germania, Belgio. — Sicca hysteriformis.

129. **Mollisia abdita** Ellis Torr. bot. Cl. 1882, p. 19. — Orbicularis, pallida, 0,6-0,7 mm. diam., margine angusto denticulato, membranaceo, initio per rimam angustam epidermidis erumpens, demum sicca omnino abdita; ascis sessilibus, tereti-oblongis, 40-45 \approx 4-5; sporidiis distichis, clavato-oblongis, 10 \approx 2-2,5; paraphysibus clavulatis.

Hab. in petiolis dejectis Juglandis regiæ, Newfield N. J.

Amer. bor. — Sicca vix ob rimam angustam epidermidis discernenda.

130. **Mollisia hysteropezizoides** Rehm in Hedw. 1885, pag. 231. — 1462
Ascomatibus sessilibus, primitus orbicularibus, dein lirelliformiter elongatis, medio impressis, atro-fuscis, in statu humido fuscis, labiis tumidis parallele juxtapositis, parenchymatice e cellulis fuscis parvulis contextis, sparsis, disco crasso, flavo; ascis clavatis, apice non incrassatis, 8-sporis, 60-70 = 18; sporidiis oblongis, rectis, hyalinis, 1-cellularibus, haud nucleatis, distichis, 15 = 7; paraphysibus filiformibus, haud clavatis, septatis, hyalinis, 2,5 μ . cr.

Hab. ad pericarpia putrida eorumque pedunculos *Epilobii Fleischeri* juxta glaciem æternam montis Ortler prope Suldental, Tyrol (REHM).

131. **Mollisia clavigera** E. et E. Journ. Myc. 1888, p. 100. — Uda 1463
protrusa, sicca contracta, hysterioidea, disco cinereo v. livido-albo, oblongo, $\frac{1}{4}$ - $\frac{1}{2}$ mm. lato, $\frac{1}{2}$ -1 mm. longo, margine fimbriato, contextu fibroso donato, fibrarum extremitatibus rugosis; ascis tereti-clavatis, sessilibus, in basin stipitiformem angustatis; sporidiis distichis, oblongis, hyalinis, obtusis, 2-3-guttulatis, 12-15 = 4-4,5; paraphysibus clavulatis, rugulosis, septulatis, ascis superantibus.

Hab. in foliis emortuis *Ammodophilæ longifoliae* Sand Coulee Montana Amer. bor. (ANDERSON).

132. **Mollisia stenostoma** B. et C. North Am. Fungi n. 744 et Cooke 1464
Disc. U. S. I, p. 297. — Erumpens, elongata, aterrima, ore angusto; ascis clavatis; sporidiis oblongis, angustis, uniseriatis, binucleatis.

Hab. in culmis *Andropogonis* in Carolina inf. Amer. bor.
 — Habitus (in siccis) minuti *Hysterii*.

PYRENOPEZIZA Fuck. Symb. p. 293 [1869] (Etym. *pyren* granum et *Peziza*), *Mollisia* Karst., Sacc., Phill. ex p. *Urceola* sect. b Quél. Ench. p. 320. — Ascoma duriusculum, minutum, sessile, globosum, demum apertum, extus atrum, verticaliter plicatum v. granulosum. Discus concavus pallidior, dein obscurus. Ascii 8-spori. Sporidia oblonga, clavata cylindraceave, continua. Paraphyses filiformes. — Formis mediis in *Mollisiam* abit.

I. Caulicolæ.

1. **Pyrenopeziza atrata** (Pers.) Fuck. Symb. pag. 294, Sacc. F. ital. 1465 t. 1398, *M. atrata* (Pers.) Karst. M. F, I, p. 200, Pat. t. 92, Pe-

piza atrata Pers. Syn. Fung. p. 669, Fr. Syst. Myc. II, p. 148, *Peziza atratula* Nyl. Obs. p. 61. — Gregaria, prominula, e subsphæroideo concava, demum planiuscula, sicca subsphæroideo- (v. difformiter) contracta, cinerea v. fuscescens, sicca nigricans, margine subconnivente albicante, ruguloso, disco albicante v. cinereo-pallido, 0,1-0,5 mm. lat.; ascis fusoideo-clavatis v. fusoideo-elongatis, 25-40 = 5-6, rarius -8, apice jodo vix tinctis; sporidiis elongatis, 5-8 = 1,5-2, simplicibus; paraphysibus numerosis, filiformibus, gracilescensibus.

Hab. ad caules herbarum variarum, præprimis Spirææ ulmariæ, Cerefolii silvestris, Asparagi officinalis, Rannunculi acris, Heraclei, Cynanchi, Gei, Gentianæ, Ambrosiæ, Eryngii, Loniceræ, Eupatorii, Solidaginis Sarraceniæ aridos sat frequenter in Fennia, Suecia, Germania, Gallia, Italia, Belgio, Britannia, Am. bor. — *Var. punctiformis*: ascomate vere punctiformi, 80-100 μ . diam., nigricante, subsuperficiali; ascis clavulatis, 25 = 8-9, paraphysatis, octosporis; sporidiis botuliformibus vel clavulatis, 8 = 1 $\frac{3}{4}$ -2, guttulatis, hyalinis. In caule *Tam i communis* in Gallia. — *Var. Asparagi* Wint. in Phill. Disc. p. 182: sparsa v. gregaria, globoso-hemisphærica, nigricans, scabra, margine albido, connivente; ascis tereti-clavatis; sporidiis oblongo-fusoideis, 8-10 = 1; paraphysibus filiformibus. In caule *Asparagi* in Germania et Britannia. — *Var. Enanthes* Phill. Disc. p. 182: gregaria, globosa, dein hemisphærica, glabra, levigata, cinerea v. nigricans, margine pallidiori; ascis fusoideis vel clavato-fusoideis; sporidiis oblongo-fusoideis, 8 = 1; paraphysibus parcis. In caule *Enanthes crocatæ* in Britannia. Ascoma 0,3 mm. lat.

2. ***Pyrenopeziza jugosa*** (Phill.) Sacc. *Mollisia jugosa* Phill. et Plow. 1466 Disc. pag. 184. — Conferta, globosa, verticaliter rugosa, scabra, nigra, disco cinereo, ore contracto; ascis clavatis; sporidiis octonis, oblongo-ellipsoideis vel fusoideis, 3-5-guttulatis dein simulate 3-5-septatis, 14-18 = 3-5; paraphysibus numerosis subcoalitis.

Hab. in caulibus emortuis King's Lynn Britanniæ. — Ascoma 2 mm. lat. *P. atratæ* affinis.

3. ***Pyrenopeziza fracta*** (B. et C.) Sacc. *Peziza fracta* B. et C. North 1467 Am. Fungi n. 742 et Disc. U. S. I, p. 296. — Minuta, erumpens, nigra, subglobosa, cito expansa fracta; ascis clavatis; sporidiis biseriatis oblongo-clavatis hyalinis.

Hab. in caule *Hydrangeæ vulgaris* in mont. Virginie Amer. bor. — Ascoma initio ore punctiformi apertum.

4. **Pyrenopeziza Chailletii** (Pers.) Fuck. Symb. myc. p. 294, *Mollisia* 1468
Gill. Disc. c. ic., *Peziza Chailletii* Pers. Myc. Eur. I, p. 288. —
Sessilis; cupulis urceolatis carnosulis gilvis, margine denticulato
nigrescente; ascis oblongo-clavatis, 8-sporis, $60 \approx 5$; sporidiis di-
stichis, cylindraceis, biguttulatis, hyalinis, c. 8μ . long.

Hab. ad caules plantarum Angelicæ silvestris (CHAILLET)
et Chærophylli aurei (MORTHIER), Fœniculi forsan et Gentianæ luteæ (MOUGEOT) in Gallia, Helvetia et Italia. — Juniores suburceolatæ fuscescentes, adultæ magis depresso subventri-
cosæ, aqua irrigatæ subturgidæ et margine dentibus subtriangula-
ribus erectis distinctius ornatæ; in statu sicco concavo-delapsæ,
margine subinflexo. Color subcervinus aut pallido fuscescens. « Cu-
pulæ (in forma italica Fœniculi) erumpentes initio et siccæ clau-
sæ, sphærioidæ, tandem et udæ apertæ extus atræ rugosæ, disco
margineque subcrenulato pallidioribus; asci clavati, apice acutati,
parce paraphysati, octospori; sporidia cylindracea, subinæquilate-
ralia, $14-18 \approx 3-4$, utrinque 1-guttulata, hyalina. Ab *Heterosphæ-*
ria Patella Fr. et *Heteropatella lacera* Fuck. bene diversa ».

5. **Pyrenopeziza Gentianæ** (Pers.) Fuck. Symb. p. 295, *Peziza Gen-* 1469
tianæ Pers. Myc. Eur. I, p. 310, *Mollisia* Pat. Tab. 286. — Erum-
pens, ceracea, concava, umbrino-nigricans, disco submolli griseo-
albido, margine sublacero; ascis oblongis, stipitatis, 8-sporis, $62 \approx 8$;
sporidiis distichis, oblongis, hyalinis, $11 \approx 3$.

Hab. ad caules emortuos Gentianæ luteæ in Vogesia, Jura,
Germania, Helvetia. — Solitaria v. aggregata, etiam disco planiu-
sculo lutescente. Sub epidermide caulis crusta quædam nigrescens
observatur, in qua residet hæc species. Sec. Fuck. crusta hæc fun-
gillo cuidam subcutaneo tribuenda videtur.

6. **Pyrenopeziza Galii** Fuck. Symb. myc. p. 295. — Ut *Pyr. Gen-* 1470
tianæ sed ascis elongatis, stipitatis, 8-sporis, $76 \approx 8$; sporidiis ob-
lique monostichis, obovatis, antice obtusis, basi attenuatis, $10 \approx 5$,
hyalinis.

Hab. in caulis siccis Galii Molluginis ad Neuhof in
Rhenogovia.

7. **Pyrenopeziza Galii-veri** (Karst.) Sacc. *Mollisia Galii-veri* Karst. 1471
M. F. I, p. 203, *Pyren. Molluginis* Rehm. Ascom. n. 257, *Pyr.*
Galii Kunze. — Sparsa v. subgregaria, primum sphæroidea, dein
hemisphærica, tandem explanata, concava, cinerea, disco dilutiore,
margine fimbriato, albicante, sicca nigricans varieque contracta,
rugulosa, 0,5 mm. lat.; ascis fusoideo-clavatis, $45-50 \approx 6-8$, apice

jodo vix tinctis; sporidiis fusoideo-filiformibus, rectis v. subrectis, guttulatis v. spurie pluriseptatis, 12-23 \approx 1-2; paraphysibus sursum leviter incrassatulis.

Hab. ad caules Galii veri siccis Mustiala Fenniae et Germania. — Asci sec. Rehm 42 \approx 9; sporidia 16-18 \approx 2,5.

8. **Pyrenopeziza nigritella** (Phill. et Harkn.) Sacc. *Mollisia nigritella* 1472
Phili. et Harkn. Calif. p. 2. — Minuta sparsa, sessilis, nigricans,
sphaerica dein patellaris, margine serrulato; ascis late clavatis;
sporidiis ovoideis, medio subconstrictis, binucleatis, 15 \approx 6; para-
physibus nullis.

Hab. in caule putri Galii borealis in California. — Habitu-
tus *sphaerioideus*,

9. **Pyrenopeziza Phyteumatis** Fuck. Symb. mycol. App. I, p. 335. — 1473
Cupulis in crista aterrima ortis, gregariis, sessilibus, 1 mm. latis,
primo clausis, demum late apertis explanatisque, extus rugulosis,
nigris, margine valde granulo sublaceratoe, concolori, disco car-
noso cæsio; ascis elongato-clavatis, sessilibus, 8-sporis, 5? \approx 8; spo-
ridiis distichis, cylindraceis, utrinque obtusis, parum curvatis, con-
tinuis, biguttulatis, hyalinis, 9 \approx 3; paraphysibus filiformibus.

Hab. ad caules putres Phyteumatis spicati in Jura Hel-
vetiae (MORTHIER).

10. **Pyrenopeziza Campanulæ** Fuck. Symb. mycol. App. II, p. 59. — 1474
Cupulis dense gregariis, nigris, media magnitudine, sessilibus, ex-
tus margineque crasso rugulosis, humidis apertis, siccis margine in-
voluto clausis, disco griseo, concavo; ascis oblongis, stipitatis, 8-
sporis, 70 \approx 10; sporidiis in asci superiori parte subcylindraceis,
continuis, hyalinis, 12 \approx 3; paraphysibus numerosis, filiformibus.

Hab. in caulis siccis Campanulæ Trachelii, greges sæpe
magnos nigros formans pr. Neuchâtel Helvetiae (MORTHIER). — Sec.
cl. Morthier huc spectat *Phyllachora Campanulæ* (DC.) Fuck.

11. **Pyrenopeziza nigrella** Fuck. Symb. mycol. App. III, pag. 30. — 1475
Cupulis in striis longitudinalibus, atris, quandoque totum caulem
occupantibus, gregariis, usque ad millimetrum latis, sessilibus, di-
sco humido subplanis, nigro-cinereo, margine crenulato, nigro, ex-
tus concoloribus, granulosis, siccis plicato-clausis, atro-olivaceis;
ascis oblongis, 8-sporis, 50 \approx 8; sporidiis distichis, ovato-clavatis,
continuis, 9 \approx 4.

Hab. in caulis siccis Galeopsidis Tetrahit pr. Neuchâ-
tel Helvetiae (MORTHIER).

12. **Pyrenopeziza pusilla** S. et S. Mich. I, p. 423, F. ital. f. 1401. 1476

— Ascomatibus gregariis, initio epidermide subvelatis, sphærioides, dein semierumpentibus, scutellaribus, vix $\frac{1}{4}$ mm. diam., ubique atris, parce hiantibus, contextu excipuli minute parenchymatico, circa marginem parenchymatico olivaceo-fuligineo; ascis subclavatis, breve stipitatis, 55-65 \approx 7, obsolete paraphysatis, octosporis; sporidiis oblique monostichis ovato-oblongis, 8-10 \approx $3\frac{1}{2}$ - $4\frac{1}{2}$, guttulatis, hyalinis.

Hab. in caule emortuo *Menthæ*, Conegliano Ital. bor. — Affinis *P. Gentianæ* sed matrix non nigrificata etc.

13. Pyrenopeziza digitalina (Phill.) Sacc. *Mollisia digitalina* Phill. 1477

Disc. p. 190. — Sparsa v. gregaria, globosa, dein expansa, extus nigra, squamulosa, margine pallido, disco cinereo; ascis clavatis, acutiusculis; sporidiis cylindraceis, rectis curvulisve, 5-7 \approx 1,3; paraphysibus filiformibus.

Hab. in caule *Digitalis purpureæ*, Shrewsbury Britanniæ.

— Ascoma 300-800 μ . diam.

14. Pyrenopeziza Bongardii (Weinm.) Sacc. *Pez. Bongardii* Weinm. 1478

Hym. Gaster. p. 476, *P. globosa* Bong. in Ms. (non Fr.). — Sparsa, sessilis, basi subcontracta, ceterum globosa, demum hemisphærica, glabra, disco subexcavato, albido, margine dentato erecto, extus nigrescens.

Hab. ad caulem plantæ aridæ *Melampyri nemorosi* in Rossia minore (Dr. BONGARD, in herbario Acad. Imp. scient. Petrop.)

— Species pulchella! Ascoma crassiusculum, firmulum satis elevatum, punctiforme, exsiccatum contractum et totum nigrum evadit.

15. Pyrenopeziza Rhinanthei (Karst.) Sacc. *Mollisia Rhinanthei* Karst. 1479

M. F. I, p. 200. — Sparsa, erumpens, sessilis, glabra, sicca nigricans et varie contracta, sæpe subtriquetra, rugulosa, humectata planiuscula, fusca, disco albicante vel pallido, margine pallidiore, 1-1,5 mm. lat.; ascis fusoideo-clavatis, 70 \approx 6-7, obturaculo minutissimo jodo obsoletissime cœrulescente; sporidiis aciculari-filiformibus, recte v. leniter curvulis, guttulatis interdum spurie uniseptatis, 18-25 \approx 2-2,5; paraphysibus filiformibus.

Hab. ad caules *Rhinanthi minoris* exsiccatos in Mustiala Fenniæ.

16. Pyrenopeziza lugubris (De Not.) Sacc. *Peziza lugubris* De Not. 1480

Comm. critt. I, p. 368. — Ascomate madore cupulari-hemisphærico, concavo, in siccis coarctato-sphæriæformi, excipulo crasso, cellulis fuscis stratosis contexto; ascis octosporis; sporidiis oblongatis, exiguis, hyalinis, continuis.

Hab. in caulis siccis Euphrasiæ, Valsesia Italiæ borealis (CARESTIA).

17. **Pyrenopeziza millepunctata** Sacc. F. Ven. Ser. IV, n. 79, F. it. 1481 f. 1397. — Ab affini *P. lugubri* præcipue recedit ascis brevioribus, cylindraceo-oblongis, 45-50 \approx 7-8; sporidiis distichis cylindraceis v. clavulatis, 6-7 \approx 2 $\frac{1}{2}$ -3; paraphysibus bacillaribus.

Hab. in caulum putrescentium basi Senecionis nemorensis et sarracenici in silva montello (Treviso), dense gregaria, et Malmedy (LIBERT). — Ascomata gregaria ex urceolato scutellata, nigricautia, extus rugulosa, vix $\frac{1}{2}$ mm. lat.

18. **Pyrenopeziza aterrima** Rehm Ascom. Ill. n. 512. — Gregaria, 1482 primitus globosa, subclausa, per epidermidem erumpens, dein subglobosa, sicca atra, corrugata, humida patellari-explanata, hemisphærica, margine tenuissimo, irregulariter eroso cincta, disco pallido, fusco-parenchymatico contexta, cellulis marginalibus elongatis, distantibus viridulo-fuscis, circ. 0,3 μ . diam.; ascis clavatis, apice obtuse acuminatis, 8-sporis, 50 \approx 10; sporidiis ellipticis, rectis v. subcurvatis, nucleis 2 apicalibus præditis, hyalinis, 24 \approx 5; paraphysibus filiformibus, hyalinis, haud clavatis.

Hab. ad caules siccis Centaureæ Jaceæ, Regenstauf prope Ratisbonam (DR. REHM).

19. **Pyrenopeziza Carduorum** Rehm Asc. n. 68, Winter in Flora 1872 1483 p. 526. — Cupulis dense gregariis v. confluentibus, sessilibus, extus glabris, papillatis, opaco nigro-fuscis, basi pilis hyalinis, repentibus cinctis, primo globosis, margine incurvo, papillato-venoso fimbriatove, demum deformibus, apertis planis, fere emarginatis, disco plano, incano-glauco, 0,5-1,7 mm. latis; ascis cylindraceis, sessilibus, 8-sporis, 50-65 \approx 7; paraphysibus simplicibus, filiformibus, hyalinis, rectis vel curvatis, 4-8-nucleatis, 12-19 \approx 1,5-2.

Hab. in caulis putridis Carduorum in Franconia.

20. **Pyrenopeziza Artemisiæ** (Lasch) Sacc. *Peziza Artemisiæ* Lasch 1484 in Rabenh. Herb. myc. n. 335. — Erumpens, tenuis, turbinata, e sphæroideo hemisphærico-applanata, scabrosella, subolivaceo-cinerea vel cinerea, disco pallidiore, margine primitus albicante, siccitate nigrescens; ascis cylindraceis; sporidiis linearibus, minutis, hyalinis.

Hab. in caulis emortuis Artemisiæ vulgaris in Germania, Britannia, Fennia et Amer. bor. — Sec. Rehm ascom. n. 66: asci clavati 50 \approx 5; sporidia oblonga, obtusula, rectiuscula, 2-guttata, 7-9 \approx 1,5-2.

21. **Pyrenopeziza subatra** (C. et P.) Sacc. *Peziza subatra* C. et P. 1485

in Peck 28 Rep. St. Mus. p. 66. — Gregaria, erumpens, atra, mollis v. ceracea; cupulis initio hemisphæricis dein apertis, levibus v. leniter rugosis, disco fuligineo, margine pallidiori; ascis subcylindraceis; sporidiis cylindraceis, rectis curvulisve, 2-3-nucleatis, 15 μ . longis.

Hab. ad caules emortuos herbarum h. e. Solidaginis et Neseæ verticillatae in America boreali. — *P. atratæ, Ebuli et sphærioideæ affinis.*

22. **Pyrenopeziza Ebuli** (Fr.) Sacc. *Mollisia Ebuli* Karst. M. F. I, 1486 p. 292, *Pez. depressula* Nyl. Obs. p. 62 (?) — Erumpens, gregaria, primo subsphæroidea, mox urceolata, demum dilatata, siccitate sphæroideo-depressa, cinereo-nigricans, margine albidiore, sicca nigricans, disco cinereo-pallescente, 0,3-0,5 mm. lat.; ascis cylindraceo-clavatis, apice attenuatis, 30-50 \approx 5-7, obturaculo apicali minutissimo jodo leviter cœrulecente; sporidiis clavulatis v. aciculari v. fusoideo-elongatis, guttulatis, 9-12 \approx 1,5-2,5.

Hab. ad caules Sambuci Ebuli et Epilobii in Germania, Fennia et Suecia. — *P. atratæ* affinis, sed cupulis erumpentibus sporidiisque longioribus.

23. **Pyrenopeziza Thalictri** (Peck) Sacc. *Peziza Thalictri* Peck 29 Rep. St. M. p. 55. — Cupulis numerosis, sessilibus per epidermidem erumpentibus minutis, siccis punctiformibus, extus atris, margine plurumque albido v. cinereo et subfimbriato, disco cinereo; ascis oblongis; sporidiis confertis, elongatis, simplicibus vel multinucleatis, 25-30 \approx 5.

Hab. ad basim cauli emortuorum Thalictri cornuti Center, Amer. bor. — Cupula uda expansa, disco conspicuo, extus scabra.

II. Ramicolæ.

24. **Pyrenopeziza aborigena** Speg. Mich. II, p. 259. — Ascomatibus 1488 minutis, 120-150 μ . diam., sparsis, superficialibus, conico-cupuliformibus, extus atris, disco glacuo-cinereo subatro; contextu obsolete celluloso, marginem versus prosenchymatico-radiante; ascis cylindraceo-clavatis, deorsum in pedicellum brevem attenuatis, apice acutiuscule truncato-rotundatis, crassiuscule tunicatis, octosporis, 40-45 \approx 4-5, apice jodi ope obsolete cœrulecentibus; paraphysibus filiformibus obvallatis; sporidiis distichis, cylindraceis, utrinque obtuse acuminatis, non vel 1-2-guttulatis, hyalinis, 5-8 \approx 2- $2\frac{1}{2}$.

Hab. in ramis decorticatis putrescentibus *Rhamni Frangulae* in montanis prope Pieve di Cadore, Ital. bor.

25. **Pyrenopeziza Rubi** (Fr.) Rehm Ascom. n. 416, *Excipula Rubi* 1489
 Fr. Syst. Myc. II, p. 190, Grev. t. 334, *Cenangium Rubi* Duby
C. rugosum Niessl, *Mollisia Rubi* Pat. f. 486. — Ascomatibus erumpentibus, subcoriaceis, sæpius complicatis, cervino-nigricantibus, disco pallidiore, e plano concavo; ascis clavulatis, 45-50 \times 6-7; sporidiis tereti-clavulatis, hyalinis, 7-8 \times 1,5-2; paraphysibus filiformibus.

Hab. in sarmentis emortuis *Rubi fruticosi*, Idæi et cæsii in Germania, Gallia, Britannia, Belgio, Amer. bor. — Var. **tenuior**, *Trochila Rubi* De Not.: ascomatibus subsuperficialibus, tenuioribus, demum totis atris et siccis varie complicatis, excipuli contextu parenchymatico, ad marginem breve prosenchymatico toto fuligineo-atro, cellulæ nonnullis prominentibus et saturationibus; ascis clavato-cylindraceis, 40 \times 7; paraphysibus bacillaribus obval-latis, octosporis; sporidiis distichis oblongo-cylindraceis, 8-10 \times $1\frac{1}{2}$ hyalinis. In sarmentis *Rubi* in Ital. bor.

26. **Pyrenopeziza lacerata** (C. et P.) Sacc. *Peziza lacerata* C. et P. 1490
 in Grevillea I, pag. 6. — Gregaria, globosa, dein expansa, atro-brunnea, margine in dentes subtriangulares, inæquales fisso; disco cinereo, dein nigricante; ascis subcylindraceis; sporidiis tereti-clavatis, 12-13 μ . longis.

Hab. in sarmentis *Rubi odorati* in Mont. Adirondack Americae borealis.

27. **Pyrenopeziza rimicola** (Karst.) Sacc. *Mollisia rimicola* Karsten 1491
 in Rev. Myc. 1885, p. 106. — Subgregaria, superficialis, sphæroidea (saltem siccitate), sublevis, atra, ore integro, rotundo, vix pallidiore, 100-150 μ . lat.; ascis fusoideo-clavatis, 36-40 \times 5-6; sporidiis elongatis v. fusoideo-elongatis, eguttulatis, hyalinis, 4-5 \times 1; paraphysibus numerosis.

Hab. in rimis corticis ramorum aridorum *Ribis alpini* pr. Mustiala Fenniæ.

28. **Pyrenopeziza Vitis** Rehm Ascom. Ill. n. 67 (1871), Wint. in Flora 1492
 1872, p. 526, *Peziza ampelina* Pass. (1874) in Rab. F. E. n. 1706. — Ascomatibus sparsis gregariisve e globoso hemisphæricis, verruculosis, fusco-atris, $\frac{1}{3}$ mm. diam., margine crenulato, contextu parenchymatico fusco; sporidiis obtuse conoideis, unicellularibus, hyalinis, 8 \times 2,5, distichis; ascis clavatis, apice incrassatis, 36-40 \times 6; paraphysibus hyalinis, filiformibus.

Hab. ad corticem trunci *Vitis viniferae* in silva Kislakensi prope Kaposvár (LOJKA) et Parma Ital. bor. (PASS.). — Videtur potius forma *Pyr. atratae*, teste ipso auctore.

29. **Pyrenopeziza Tamaricis** (Roum.) Sacc. Mich. II, p. 536, *Peziza* 1493

Tamaricis Roum. F. Gall. n. 263, *Mollisia Myricariæ* Bres. Rev. Mycol. IV, pag. 212 et 221, *Mollisia Tamaricis* (Roum.) Bres. Trid. p. 42, t. 45, f. 2. — Subcæspitosa, hemisphærica, dein expansa, mutua pressione contorta, fuligineo-tabacina, margine prosenchymatico, dein pallidiore, furfuracea, disco griseo-brunneo, subinde flavidolivaceo tincto, levi 1-2 mm. lato; ascis subfusoideis, 40-50 \approx 5-6; sporidiis tereti-fusoideis, sæpe curvulis, 6-8 \approx 2-2,5, hyalinis; paraphysibus filiformibus.

Hab. in ramis decorticatis *Myricariæ germanicæ*, *Tamaricis narbonensis*, *Robiniae* in agro Tridentino, in Gallia, Germania, Britannia. — Var. **argentina**, cfr. Speg. F. Arg. Pug. IV, n. 228: ascomatibus udis explanatis, undulatis, subpellucidis, glaberrimis, extus olivaceis, disco pallidiore, 0,5-1 mm. diam., siccis contracto-globulosis v. subhysteriformibus, rugulosis, atris, fragilibus; ascis cylindraceo-clavatis, antice obtusiuscula rotundatis, deorsum breve crasseque stipitatis, 35-40 \approx 3,5-5; paraphysibus filiformibus obvallatis; sporidiis oblique monostichis vel distichis, subelliptico-cylindraceis, utrinque obtuse rotundatis ac 1-guttulatis, 7-8 \approx 1,5, hyalinis. In cortice duriore trunci adhuc vivi *Tamaricis africanae* in Argentina.

30. **Pyrenopeziza rugulosa** Fuck. Symb. mycol. pag. 293. — Cupulis 1494

quandoque in massam granulosam nigram confertis, superficialibus, sessilibus, usque ad 1 mm. diametr., globosis, primo punctiformibus, globosis, perforatis, demum magis apertis, margine connivente, extus granuloso, subverticaliter striato-rugosis, nigris, disco atro-fusco; ascis oblongis, 8-sporis, 28-30 \approx 4-5; sporidiis cylindraceis, subrectis, 6 \approx 1,5 \approx 2, hyalinis.

Hab. in ramis decorticatis putridis Carpinii ad Nepfelbach in silva Cœstrich Rhenogoviæ.

III. Foliicolæ.

31. **Pyrenopeziza Eryngii** Fuck. Symb. pag. 294. — Cupulis sparsis, 1495
sessilibus, 2 mm. diam., nigris, margine concolore, denticulato, demum clauso, disco vix pallidiore; ascis oblongis, 8-sporis, sessilibus, 70-74 \approx 10-12; paraphysibus asci longitudine, clavatis; spori-

diis distichis, oblongis, parum curvatis, utrinque obtusis, biguttulatis, 15-16 \approx 4,5-5, hyalinis.

Hab. in foliis putridis *Eryngii campestris* in Rhenogovia et Bavaria.

32. **Pyrenopeziza nigrificans** (Wint.) Rehm Ascom. n. 862, *Niptera* 1496
nigrificans Winter F. Eur. n. 2832. — Cupulis superficialibus, sparsis v. gregariis, humectatis patellæformibus seu fere scutellatis, rarius regulariter orbicularibus, immo plerumque irregulariter angulatis, flexuosis repandisque, difformibus, $\frac{1}{2}$ -2 mm. latis, extus fusco-atris, rivulosis, margine crenato et denticulato, erecto, disco plano v. subconcavo, cinereo; siccis hemisphæricis, margine involuto, fere clausis, contextu parenchymatico; ascis cylindraceis, deorsum parum attenuatis, 8-sporis, 50-70 \approx 6-7; paraphysibus filiformibus, sursum plus minus incrassatis, saepe clavatis, usque 5 μ . crass., hyalinis; sporidiis monostichis vel subdistichis, ellipsoideis seu ellipsoideo-oblongis, continuis, hyalinis, guttulis 2 præditis, 7-8,5 \approx 3,5-4 crassis.

Hab. ad *Adenostyolidis alpinæ* petiolos putridos, Rigi pr. Luzern Helvetiæ (G. WINTER). — Mycelium effusum, substratum longe lateque (etiam profunde) penetrans et nigrificans.

33. **Pyrenopeziza Medicaginis** Fuck. Symb. p. 295. — Cupulis sessilibus, 1 mm. latis, primo globosis, clausis, dein apertis, hemisphæricis, extus atro-fuscis, rugulosis, margine elevato, crenulato, concolore; disco pallide griseo; ascis oblongo-clavatis, 8-sporis, 80 \approx 10; sporidiis distichis, oblongo-ovatis, utrinque uniguttulatis, 10 \approx 5, hyalinis; paraphysibus linearibus, rectis, asci longitudine.

Hab. ad folia et petiolos *Medicaginis sativæ* pr. Cœstrich Rhenogoviæ.

34. **Pyrenopeziza Agrostemmatis** Fuck. Symb. myc. p. 295, tab. VI, 1498 f. 34. — Cupulis superficialibus, sparsis, sessilibus, $\frac{1}{2}$ mm. latis, extus nigris, rugulosis, primo globosis, demum apertis planisque, margine denticulato, disco pallescente, carnosæ; ascis oblongo-clavatis, stipitatis, 8-sporis, 84 \approx 12; paraphysibus elongato-clavatis; sporidiis in asci superiori parte subdistichis, clavatis, curvatis, guttulatis, in parte angustiore obscure (spurie) uniseptatis vel continuis, guttulatis, hyalinis, 20 \approx 5.

Hab. in foliis languidis *Agrostemmatis Githaginis* pr. Hallgarten Rhenogoviæ. — Stat. conidicus, teste Fuck., est *Marsonia Deslastrei* (De Lacr.).

35. **Pyrenopeziza Delavayana** Sacc. *Mollisia nigrella* Pat. Champ. 1499

Chine pag. 1 (nec Fuckel). — Globulosa, nigra, 0,5-1 mm. lata, extus glabra, ore sinuoso, disco cinereo; ascis clavatis, longe stipitatis, 53-60 \approx 10-12; sporidiis ovoideis, hyalinis; paraphysibus filiformibus, ramulosis.

Hab. in foliis radicalibus Campanulæ dimorphanthæ Yunnan Sinarum (DELAVAY).

36. **Pyrenopeziza Plantaginis** Fuck. Symb. mycol. p. 294, *Pez. atrata* 15 form. *foliicola* Desm Crypt. n. 1313. — Cupulis plerumque in macula expallescente, sessilibus, gregariis, minutis, punctiformibus, primo clausis, globosis, demum apertis, extus nigro-furfuraceis, margine involuto, disco concavo, pallide fusco; ascis elliptico-oblongis, 8-sporis, 40-55 \approx 7-9; sporidiis distichis, elongato-lanceolatis, rectis, 12-16 \approx 3-4, hyalinis, 2-guttulatis granulosisque.

Hab. in foliis scapisque emortuis Plantaginis lanceolatae in Germania, Italia, Britannia, Belgio.

37. **Pyrenopeziza nervisequia** (Pers.) Sacc. *Mollisia* Phill. Disc. p. 179, 15 t. VI, f. 33, *P. nervisequia* Pers. M. E. I, p. 308, Fr. Syst. Myc. II, p. 153. — Seriata, nigra, cupulis in subiculo indurato linearis insidentibus, disco concavo, cinereo v. nigro; ascis clavatis; sporidiis oblongo-ellipsoideis vel oblongis, 13 \approx 3; paraphysibus filiformibus.

Hab. in foliis emortuis Plantaginis lanceolatae in Vogesis (MOUGEOT), Batheaston Britanniæ (BROOME). — In nervis foliorum observantur lineæ pariter nigræ, super quas positæ sunt cupulæ parvæ subpresso-flexuosaæ.

38. **Pyrenopeziza acicola** S. et S. Mich. I, p. 423, F. it. f. 1405. — 15 Ascomatibus sparsis erumpenti-superficialibus, totis atris, duriusculis, $\frac{1}{4}$ mm. diam., concaviusculis, contextu obsolete celluloso, ad marginem radiante ochraceo; ascis cylindricis brevissime stipitatis, paraphysatis, 40-45 \approx 6, octosporis; sporidiis oblique monostichiæ bacillari-clavulatis, 7-8 $\frac{1}{2}$ \approx 1-1 $\frac{1}{2}$, curvulis, hyalinis.

Hab. in acubus putridis Pini silvestris, Susegana Ital. bor.

39. **Pyrenopeziza protrusa** (B. et C.) Sacc. *Peziza protrusa* B. et 15 C. North Am. Fungi n. 745 et Cooke Disc. U. S. I, pag. 297. — Erumpens, punctiformis, epidermide hic illic circumdata, ore flexuoso, extus granulata castanea, intus concava alba.

Hab. in pag. inf. foliorum emortuorum Magnoliæ glaucæ in Carolina inf. Amer. bor.

40. **Pyrenopeziza sphæroides** (Pers.) Fuck. Symb. App. I, pag. 335, 15 *Excipula sphæroides* (Pers.) Fr. Syst. Myc. II, p. 191, *Xyloma*

Pers. Syn. pag. 106. — Erumpens, initio globosa, nigricans, dein subexpansa, margine erecto dein collapso crispa, disco aquoso-cinereo, concavo, demum atro; ascis sessilibus, oblongis, 8-sporis, $36 \approx 6$; sporidiis distichis, cylindraceis, continuis, hyalinis, $6 \approx 2$; paraphysibus filiformibus.

Hab. in pag. inf. foliorum putrium *Salicis capreæ*, Oestrich Rhenogoviæ. — Forma **Lychnidis**, cfr. Phill. Disc. p. 187: sporidiis $10-15 \approx 1-1,5$, ascis tereti-fusoideis. In caulis emortuis *Lychnidis dioicae* King's Lynn Britanniæ (PLOWRIGHT).

41. **Pyrenopeziza foliicola** (Karst.) Sacc. Mich. I, pag. 65, *Mollisia foliicola* Karst. Fenn. I, p. 201. *Trochila atrata* * *Tr. foliicola* Karst. Symb. p. 244. — Gregaria, subsphæroidea aut hemisphærica v. concava, pallida v. cinereo-pallida, sicca sphæroideo-contracta, nigricans marginequè albo-flocculoso, circiter 0,1-0,2 mm. lat.; ascis $36-40 \approx 5-6$; sporidiis fusoideo-elongatis, $6-10 \approx 1,5-2$.

Hab. supra folia *Alni* putrescentia circa Mustiala Fenniæ. — Var. **quercina** Sacc.: ascis clavato-oblongis, $20-25 \approx 5$, paraphysatis, octosporis; sporidiis fusoideis, $7-8 \approx 1\frac{1}{2}-2$, hyalinis. In foliis putrescentibns *Quercus pedunculatae* in Ital. bor. et Gallia. Verisimiliter huc spectat *Peziza Rabenhorstii* Awd. in Herb. mycol. n. 222, De Not. Prof. Discom. in Comm. critt. ital. I, p. 368. *P. Massoniana* De Not. in Tassi Fl. Sen. p. 61, forte huc spectat, sed specimina authentica non vidi.

42. **Pyrenopeziza Heteromelis** Phill. et Harkn. Grev. XIII, p. 22. — 1506 Gregaria, erumpens, demum sessilis; cupulis 0,2-0,5 mm. lat., sphæricis dein expansis, leniter scabrosis, atro-brunneis, margine serrato, disco fuligineo flavescente; ascis cylindraceo-clavatis; sporidiis ellipticis, $10 \approx 5$, flavo-tingentibus; paraphysibus filiformibus, ad apicem incrassatis, flavescentibus.

Hab. in pag. inf. foliorum *Heteromelis arbutifoliae* in California (HARKNESS).

43. **Pyrenopeziza betulinola** Fuck. Symb. myc. pag. 294, *Calloria Winteri* Kunze F. sel. n. 283. — Cupulis gregariis in macula ex-pallescenti, sessilibus, $\frac{1}{2}$ mm. latis, primo globosis, demum apertis, margine conniventis, extus nigris, scabris substriatisve, margine parum lacerato, disco concavo pallidiore fusco flavescenteve; ascis oblongis vel clavatis, antice acuminatis, in stipitem attenuatis, 8-sporis, $56 \approx 8$; sporidiis submonostichis, oblongis, utrinque obtusis, hyalinis, guttulatis, $8-10 \approx 3$.

Hab. ad folia putria *Betulæ albæ* in silva Mittelheim Rhe-

nogoviæ. — Var. **rubicola** Rehm Ascom. n. 572: sporidiis clavulatis, uno apice acutioribus, 8-9 \approx 2-2,5; ascis 36 \approx 6-7. In pag. inf. foliorum Rubi fruticosi in Helvetia (WINTER).

44. **Pyrenopeziza emergens** (Phill. et Harkn.) Sacc. *Mollisia emergens* Phill. et Harkn. Calif. p. 2, nec Karst. — Sparsa, erumpens, sicca compressa, nigro-brunnea, margine involuto, striatulo, disco griseo-cyaneo; sporidiis ovoideis, 7-9 \approx 3-4; paraphysibus filiformi-clavulatis.

Hab. in pag. inf. foliorum emortuorum Eucalypti Globuli in California. — Ascoma 0,5 mm. lat.

45. **Pyrenopeziza viburnicola** (B. et Br.) Sacc. *Peziza viburnicola* B. et Br. Ann. N. H. n. 1170 (1866). — Subglobosa, minuta, punctiformis, dein hemisphærica, grisea, extus granulata, margine denticulato, furfuraceo, disco pallidiori; ascis clavatis; sporidiis lanceolatis, 10-12 μ . longis.

Hab. in foliis emortuis Viburni, Shere Surrey Britanniæ. — Cupula extus cellulis dense fasciculatis, subinde in pilos paucos breves flexuosos productis granulata.

IV. Ligni-corticicolæ.

46. **Pyrenopeziza ligni** (Desm.) Sacc. Mich. II, p. 611, F. it. f. 1399, 155 *Moll. ligni* (Desm.) Karst. M. F. II, pag. 204, *Cenangium ligni* Desm. Exs. I, n. 2014, Tul. Carp. III, p. 169, t. XX, f. 5-7, *Trochila* De Not. Disc. p. 15. — Ut plurimum cæspitose aggregata, e sphæroideo concava, sessilis v. subsessilis, fusco-cervina, sicca varie contracta et fusco-nigricans, 0,8-1 mm. lat.; ascis cylindraceo-clavatis, sessilibus, obtusis, 45-55 \approx 4-5,5; sporidiis fusoido-elongatis vel fusoido-oblongatis, distichis, vulgo rectis et eguttulatis, 6-12 \approx 1,5-2,5; paraphysibus filiformibus.

Hab. ad lignum Quercus, Salicis capræ, Coryli, Populi, Castaneæ et Cratægi coccineæ vetustum in Fennia, Germania, Gallia, Italia. — Variat sporidiis et ascis nec non cupulis paullo minoribus. Typus aberrans. Spermogonia modo ascomatibus mixta, nunc ab iis segregata, sphæriiformia, sessilia, sub-sphæroidea, gregaria, fusco-nigra, circiter 0,3 mm. lat.; sporulæ cylindraceæ, rectæ v. leniter curvulæ, simplices, 3-3,5 \approx 1.

47. **Pyrenopeziza lignicola** (Phill.) Sacc. *Mollisia lignicola* Phill. Disc. p. 180. — Gregaria v. sparsa, sessilis, initio globosa dein leniter expansa, extus scabrosa et verticaliter rugulosa, brunnea v. brunnneo-nigra, sicca nigra, margine ob cellulas elongatas inæquales

subfimbriato, disco pallido; ascis tereti-clavatis, $35 \approx 4$; sporidiis cylindraceis v. fusoideis, obtusis, rectis curvulisve, $5-8 \approx 1,5-2$; paraphysibus obsoletis.

Hab. in ligno vetusto fabrefacto Bagley Wood Berkshire Britanniae. — Ascoma 0,5-1 mm. lat.

18. **Pyrenopeziza cariosa** (Peck) Sacc. *Peziza cariosa* Peck 24 Rep. 1512 of the Bot. 1872, p. 95. — Dense gregaria, sessilis, tenuis, applicata, subconcava, glabra, atra, madore atro-brunneo-rufescens, margine distincto, flexuoso vel in individuis majoribus anguloso; sporidiis oblongis, continuis, hyalinis.

Hab. in lignis putridis in montibus Catskill Amer. bor. — Vix 1 mm. lat.

19. **Pyrenopeziza Stilbum** Fuck. Symb. p. 294. — Cupulis ad basim 1513 fungi conidiophori natis, brevissime stipitatis sessilibusque gregariis, pusillis, 74μ . diam., semper subclausis, atris, vertice (margine) candide ciliatis, ciliis cupulam diametro æquantibus; ascis oblongo-clavatis, 8-sporis, $27 \approx 6$; sporidiis oblongo-cylindraceis, hyalinis, utrinque uniguttulatis, continuis, $8 \approx 3$.

Hab. ad lignum putre Carpini Betuli, rarius, Äpfelbach Rhenogoviæ. — Fungus conidiophorus *Stilbum* refert. Hyphis erectis, fertilibus brevioribus, atris, 107μ . altis, apice capitulum brevissime ramosum, conidiophorum, album gerentibus, conidiis cylindraceis, subcurvatis, $4 \approx 2$. — An *Cyathicula*?

V. Graminicolaæ.

50. **Pyrenopeziza Karstenii** Sacc. Mich. II, p. 329, F. ital. fig. 1402, 1514 *Mollisia graminis* Karst. Fenn. I, 195, nec Desm. — Gregaria vel sparsa, erumpens, concava, rugulosa, umbrina, disco umbrino-pallido v. griseo-pallente, margine albo-villosulo-fimbriato, circiter 0,4 mm. lat.; ascis cylindraceo-clavatis, $40-60 \approx 5-6$, obturaculo minutissimo jodo obsolete cœrulecente; sporidiis aciculari- v. fusoideo-elongatis, rectis v. leniter curvulis, guttulatis v. spurie tenuiter 1-septatis, $8-12 \approx 1,5-2$; paraphysibus filiformibus, sursum leviter incrassatis.

Hab. ad culmos Poæ cæsiæ, Agrostidis et Airæ cæspitosæ putrescentes per totam Fenniam et Lapponiam in Gallia, Germania et Groenlandia. — Var. **Caricis** Rehm Ascom. n. 711: erumpens, gregaria, concava, sicca rugulosa, umbrino-fusca, disco pallidiore, $\frac{1}{2}$ mm. diam., contextu fusco parenchymatico, cellulis marginalibus elongatis distantibus, ex hyalino fuscidulis, $15 \approx 5$; ascis

clavulatis, 40 \approx 8; sporidiis acicularibus, 2-guttulatis, 2-3-stichis, 12 \approx 2, hyalinis; paraphysibus clavulatis. Ad folia sicca Caricis in Saxonia (KRIEGER).

51. **Pyrenopeziza stictoidea** Sacc. Michel. II, p. 259, F. ital. f. 1403. 1515
— Ascomatibus seriatim gregariis, innatis epidermide diu velatis, tandem somierumpentibus, urceolato-depressis, nigricantibus, $\frac{1}{5}$ - $\frac{1}{4}$ mm. diam., margine involutis, disco exiguo pallidiore; ascis fusoides vel obclavatis, subsessilibus, 35-38 \approx 6, sursum acutatis; paraphysibus filiformibus apice curvulis, obvallatis, octosporis; sporidiis distichis tereti-fusoideis, 8-10 \approx $1\frac{1}{2}$ - $1\frac{3}{4}$, rectiusculis, 3-guttulatis, hyalinis.

Hab. in foliis emortuis Hierochloes v. affinis graminis in *M. Sol*, Bassano Italiæ bor. (A. Rossi). — Fungillis pluribus cognata species, nempe *Stictidi exiguae* Desm., *St. obvelatae* (Lacr.) De Not., *Trochilæ pusillæ* (Lib.) Speg., quæ revera aptius *Pyrenopezizæ* generi adscribendæ videntur.

52. **Pyrenopeziza fuegiana** Speg. Fung. Fueg. n. 339. — Cupulis superficialibus sparsis v. hinc inde laxe gregariis, e globoso-hemisphæricis, 250-300 μ . diam., disco applanatis v. vix concavis sordide fusco-fuligineis, extus atris glabris, contextu parenchymatico fuligineo; ascis cylindraceis v. subclavatis, antice rotundatis crassiusculeque tunicatis, deorsum breviter attenuato-pedicellatis, 35-40 \approx 10, octosporis; paraphysibus filiformibus densiusculis; sporidiis oblique monostichis, cylindraceo-fusoideis, utrinque acutiusculis, 6-7 \approx 1,5-2, hyalinis, continuis. 1516

Hab. ad folia viva v. languida Poë minutæ cujusdam in silvis prope Ushuvaia Fuegiae.

53. **Pyrenopeziza Arundinariæ** (B. et C.) Sacc. *Peziza Arundinariæ* Berk. et Cooke Disc. Unit. St. I, p. 297. — Applanata, mox sedens, piceo-nigra, subiculo orbiculari, maculiformi insidens, disco pallido; ascis..... 1517

Hab. in culmis Arundinariæ in Carolina Amer. bor.

54. **Pyrenopeziza denigrata** (Kunze) Rehm Ascom. n. 353, *Niptera denigrata* Kunze F. sel. n. 180 (absque diagnosi). — Ascomatibus gregariis, erumpentibus ex urceolato scutellatis, $\frac{1}{3}$ mm. diam., plumbeo-nigricantibus, glabris; ascis oblongis, 40-50 \approx 14; sporidiis elongatis, 2-guttulatis, hyalinis, 15-17 \approx 4-5, distichis; paraphysibus filiformibus. 1518

Hab. ad caules Antherici ramosi in Germania (KUNZE) et culmos Festucæ tenellæ in Amer. bor. (ELLIS).

55. **Pyrenopeziza multipuncta** (Peck) Sacc. *Peziza multipuncta* Peck 1519

33 Rep. pag. 31. — Cupulis numerosis, minutis, 150-250 μ . lat., culmicolis v. raro hypophyllis, brevibus, sessilibus, madore pallide umbrinis, siccis fusco-nigris, ore angusto, hymenio albido; ascis sessilibus, subcylindricis, 40-50 \approx 4-5; sporidiis confertis vel biseriatis, cylindricis, rectis, hyalinis, 12 μ . long.; paraphysibus filiformibus.

Hab. in culmis emortuis Caricis stellulatae, raro in foliorum mortuorum pagina inferiore, Albany Amer. bor. (CH. H. PECK).

56. **Pyrenopeziza cervinula** (Cooke) Sacc. *Peziza cervinula* Cooke 1520

Hedw. 1875, p. 84, Disc. U. S. I, p. 297. — Subgregaria, sessilis, minutissima, $1/10-1/5$ mm., cupula globosa, primum poro aperta dein hemisphærica ad marginem pallidiorem contracta, extus cervinula disco albido; ascis clavatis 40 μ . longis; sporidiis cylindraceis, rectis v. curvulis, hyalinis, 12-14 \approx 2.

Hab. ad culmos Caricis, New Jersey Amer. bor. (ELLIS).

57. **Pyrenopeziza subconica** (Rehm) Sacc. *Mollisia subconica* Rehm 1521

Ascom. Alp. n. 21. — Gregaria, orbicularis, lata basi sessilis, nigra, sicca collapsa et extus subrugulosa, humida subconica, ore albidulo poriformi subdilacerato, disco et excipulo fuscidulo, parenchymatico-fusco-contexta, c. 0,2 mm. diam.; ascis (inevolutis) clavatis, 8-sporis, 36 \approx 6; sporidiis ellipticis, rectis, 1-cellularibus, hyalinis, distichis, 7 \approx 2; paraphysibus filiformibus, hyalinis, haud clavatis.

Hab. ad culmos sicclos Junci Hostii Taschach-Gletscher in valle Piz alpium Tyrolens., alt. c. 2000 metr. (DR. REHM).

58. **Pyrenopeziza corcellensis** Sacc. *Miscell. mycol.* I, pag. 23. — 1522

Ascomatibus laxe gregariis epidermide diu velatis, dein erumpentibus, plano-scutellatis, $1/4$ mm. diam., totis nigricantibus, margine parum elevato, subinæquali; excipuli contextu parenchymatico fusco, ad marginem prosenchymatico subhyalino; ascis fusoideis, brevissime stipitatis, apice obtuse attenuatis, octosporis; paraphysibus bacillaribus apice anguste clavulatis; sporidiis subtristichis, fusiformibus, rectis, curvulisve, acutiusculis, 30-32 \approx 3,5-4, granulosis hyalinis.

Hab. in foliis calamisque Luzulæ albidæ in silvis Corcelles pr. Neuchatel Helvetiae (MORTIER). — Affinis *Trochilæ diminuenti* Karst., sed sporidia longiora et paraphyses vix apice clavatae.

59. **Pyrenopeziza fœcunda** (Phill.) Sacc. *Mollisia fœcunda* Phill. 2523

Disc. p. 189. — Sparsa, minuta, initio epidermide tecta dein erum-

pens, concava, uda fuliginea, margine pallidiore, disco fusco-atro; ascis late clavatis; sporidiis fusoideis, 3-4-nucleatis, 25-30 \approx 4-5; paraphysibus filiformi-clavulatis.

Hab. in calamis putridis *Eleocharidis*, Forres Scotiæ (J. KEITH). — Ascoma $\frac{1}{2}$ mm. lat.

VI. In Cryptogamis.

60. **Pyrenopeziza Arenula** (A. et Schw.) Sacc. *Peziza Arenula* A. 1524

et Schw. Consp. p. 341, Fr. Syst. Myc. II, p. 148, *Mollisia Arenula* (A. et S.) Karst. Rev. p. 136. — Sparsa, sessilis, subsphæroidea, semper conniventi-clausa, perpusilla, sicca, nudo oculo fere invisibilis, glabra, fuliginea, intus albida; ascis 28-30 \approx 7-8; sporidiis fusoideo-elongatis, simplicibus, hyalinis, 6-8 \approx 1,5.

Hab. in frondibus *Pteridis aquilinæ* putridis in Mustiala Fenniæ, in Suecia et Germania. — Arenæ granulo similis.

61. **Pyrenopeziza urnicola** Mout. et Sacc. (sp. nov.). — Ascomatibus 1525

ex urceolato scutellatis, superficialibus, minimis, sparsis, atris, extus minute rugulosis; ascis tereti-oblongis, 60 \approx 8, octosporis; sporidiis fusoideis, utrinque acutis, continua, hyalinis, 2-guttulatis, 12 \approx 2-2,5.

Hab. in capsulis siccis *Polytrichi* prope Liège Belgii. — Ascomata 110-130 μ . lat., saepe in areolis albicantibus nascentia.

62. **Pyrenopeziza thallophila** (Karst.) Sacc. *Moll. thallophila* Karst. 1526

M. F. I, p. 205. — Sparsa v. gregaria, sessilis, subsphæroidea v. hemisphærica, sicca varie contracta, fusca vel umbrina, subfurfurilla, rugulosa, 0,2-0,4 mm. lat.; ascis clavatis, 30-48 \approx 6-7,5, jodo luteolis; sporidiis distichis, subellipsoideis v. oblongato-ellipsoideis, rectis, vulgo guttulis 2 mediocribus præditis, 7-8 \approx 3-3,5; paraphysibus gracilescensibus, apicem versus incrassatis.

Hab. supra thallum apotheciaque *Lecanoræ* subfuscæ, ad corticem *Salicis viminalis* et *Alni incanæ* crescentis in Mustiala Fenniæ. — Externa facie *M. atratam* nonnihil refert, partibus vero internis magis cum præcedente et sequente convenit.

63. **Pyrenopeziza kolaensis** (Karst.) Sacc. *Mollisia kolaensis* Karst. 1527

M. F. I, p. 201. — Aggregata, hemisphærica, subflocculoso-furfurilla, fusco-nigricans (humectata), margine albido, disco pallido, 0,2-0,3 mm. lat.; ascis subfusoideo-elongatis, 25-30 \approx 4-5, apice jodo non cœrulecente; sporidiis elongatis, 5-8 \approx 1,5.

Hab. supra Corticum ad corticem vetustum *Salicis* crescentis, in regione Kolaensi Fenniæ. — Ab auctore ut subspecies *P. atratæ* habita.

TAPÉSIA Pers. M. E. I, p. 220, Fuck. Symb. mycol. p. 300 (Etym. *tapes*). — Ascomata subiculo plus minusve tomentoso, saepe conidiophoro incidentia, minuta, sessilia raro substipitata, ceracea v. duriuscula, urceolato-scutellata, extus glabra furfuracea v. pilosa. Ascii elongati plerumque 8-spori, paraphysati. Sporidia oblonga, oblongo-ovata, cylindraceave, continua, hyalina.

I. *Eu-Tapesia*. Ascoma glabrum v. granulosum.

A. Fructificatio nota.

1. **Tapesia sanguinea** (Pers.) Fuck. Symb. pag. 303, Pat. Tab. anal. 1528 f. 91, *P. sanguinea* Pers. disp. p. 34, Myc. Eur. pag. 273, Nees Syst. f. 271, Fr. Syst. Myc. II, p. 110. — Cupulis concavis, glabris, nigrescentibus, basi tomento brevi sanguineo cinctis; ascis clavatis, antice obtusis, 8-sporis, $56 \approx 8$; sporidiis distichis, oblongo-ovatis, hyalinis, $8 \approx 3$; paraphysibus ramosissimis.

Hab. ad ligna exsiccata duriora Abietis, Fagi, Piri, Populi in Suecia, Germania, Gallia, Helvetia, Belgio, Britannia, Amer. bor. — Cupulae pusillæ, plus minus aggregatae, juniores molles fuscescentes, margine rubello cinctæ, substantia atro-sanguinea, dein induratæ, nigræ, apothecium *Lecideæ* vel *Sphaeriam* collapsam referentes. Subiculum proprie non contextum; sed tomentulum basin cupularum cingit et quando hæ valde confertæ, tomentum in subiculum confluit.

2. **Tapesia atro-sanguinea** Fuck. Symb. pag. 303. — Cupulis primo 1529 clausis, demum apertis subplanis concavisve, 2 mm. latis, breviter crasse atre stipitatis, extus ochraceo-pulverulentis, disco fusco, deinde nigro, margine distincto, candido, e fibrillis longissimis, repentibus, saturate sanguineis, persistentibus, saepe in massam gelatinosam conidiophoram, diaphanam confluentibus ortis; conidiis subglobosis, minutissimis; ascis oblongis, substipitatis, 8-sporis, $24 \approx 6$; sporidiis distichis, cylindraceis, curvatis, hyalinis, $4 \approx 2$.

Hab. in ligno putri, molli Betulæ, Fagi, Cerasi pr. Mappen et Frankensteiner Kopf Rhenogoviæ. Prædicta species.

3. **Tapesia mollisoides** (Schw.) Sacc. Mich. II, p. 140, *Peziza* Schw. 1530 Syn. p. 174. — Ascomatibus per exiguis, $\frac{1}{8}-\frac{1}{7}$ mm. diam., plano-scutellatis levibus sessilibus, fuligineis, disco pallidiore, excipulo parenchymatico fusco, cellulis marginalibus longioribus dilutioribus; ascis clavatis, $50-55 \approx 10$, breve stipitatis apice rotundatis; paraphysibus filiformibus obvallatis, octosporis; sporidiis oblongis vel

subfusoideis, 8-9 \times 3, leniter curvis quandoque plasmathe bipartito, hyalinis.

Hab. in ramis decorticatis New England (SPRAGUE), Pensylvania (SCHW.), Carolina (RAVENEL). — Subiculum atro-fuligineum *Helminthosporium humile* sistens: hyphis filiformibus apice obtusis septulatis, fuligineis; conidiis tereti-oblongis, 15-22 \times 6, 3-septatis, apice rotundatis, fuligineis, acrogenis.

4. **Tapesia minutissima** Fuck. Symb. App. I, p. 48. — Subiculo late effuso, olivaceo-nigro, strigoso; cupulis gregariis, minutissimis, sessilibus, subcarnosis, primo clausis, extus fusco setulosis, disco concavo, fusco-griseo; ascis stipitatis, oblongo-ovatis, 8-sporis, 48 \times 10; sporidiis distichis, oblongo-ovatis, utrinque obtusis, biguttulatis, continuis, hyalinis, 8 \times 4; paraphysibus linearibus, 6-8-guttulatis.

Hab. in disco truncorum Fagi, pr. Östrich Rhenogoviæ. — Forma **meiospora** Sacc. Mich. II, p. 260, F. ital. f. 1433: ascomatibus punctiformibus, 120-150 μ . diam., nigricantibus, intus pallidioribus, setulis fuscis præcipue basi vestitis; ascis clavulatis, 60-70 \times 12-14, paraphysatis, octosporis; sporidiis distichis, oblongo-ovatis, utrinque rotundatis, 10-12 \times 4, 2-guttatis, hyalinis. In ligno Mali in Italia boreali.

5. **Tapesia Gaillardii** Roum. et Pat. Tab. anal. p. 37, f. 90. — Cupulis subsessilibus, concavis, minutis, $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. lat., gregariis, subiculo atro-brunneo, demum subevanescente insertis, flavidis, intus pallidioribus, initio subclausis citrinisque; ascis subclavatis, filiformi-paraphysatis, 8-sporis; sporidiis ovoideis, 2-3-guttulatis, hyalinis, 5-6 \times 2 (ex icono).

Hab. in ligno putri Alençon Galliæ (GAILLARD).

6. **Tapesia Legeriana** (Gill.) Sacc. *Phialea Legeriana* Gill. Disc. p. 111, cum icono. — Subiculo albo, copioso; cupulis sessilibus, subglobosis, dein planis, junioribus orbicularibus, dein sinuosus, extus griseis, 1-2 mm. lat.; disco albido; ascis cylindraceis; sporidiis oblongis.

Hab. in ligno nudo subputrido in silvis Galliæ bor.

7. **Tapesia vincita** (C. et P.) Sacc. *Peziza vincita* C. et P. in Grevillea I, pag. 6. — Gregaria, sessilis, ceraceo-mollis, irregularis, initio subglobosa, dein expansa, margine incurvo glabra, saturate atro-brunnea, fere nigra, disco plano cinereo; ascis subcylindraceis; sporidiis cylindraceis obtusis, rectis curvulive, 12-13 μ . longis.

Hab. in ligno putri Sandlake N. Y. Amer. bor. — Ascomata

pilis delicatis, brunneis, subiculum arachnoideum formantibus inserta.

8. **Tapesia aureliella** (Nyl.) Karst. Rev. monogr. p. 137, *Peziza aureliella* Nyl. Obs. p. 49, *Mollisia* Karst. M. F. I, p. 206. — Subiculo albido v. albido-flavescente furfurello-subtomentoso; cupulis albidis v. pallido-albidis, siccitate pallido-luteis, gyalectoideis, sessilibus, 0,2-0,4 mm. lat., margine mediocri, tenuiter albo-villosulo-fimbriato aut glabrescente; ascis oblongatis, 40-6-7, apice summo solutione jodi cœrulecente; sporidiis oblongatis, continuis, 5-8-2-2,5.

Hab. ad lignum putridum Pini prope Helsingfors Fenniæ. — Filamenta subiculi satis obsoleta, incoloria, circiter 7 μ . crass. et forsitan aliena; paraphyses vix ullæ v. tandem raræ, graciles; ascomata præsertim juvenilia granuliformia. An huc accedit *Peziza aurelia minor* Weinm. in Flora 1832, p. 455?

9. **Tapesia byssina** Fuck. Symb. p. 302, Rehm Ascom. Lojk. n. 33. 1536 — Cupulis sessilibus, explanatis, lobato-plicatis, medio adfixis, 4-6 mm. latis, disco albido, margine elevato, fusco, subtus byssso candida, delicatula sed persistente; ascis linearibus, oblique stipitatis, 8-sporis, 48-50-5-6, apice jodo cœrulecente; sporidiis distichis, cylindraceis, rectis, 8-11-2-3, hyalinis; paraphysibus filiformibus.

Hab. in ligno putrido, udo pr. Estrich Rhenogoviae (Fuck.) et Kaposvar Hung. (LOJKA). — Quoad affinitatem cum *Moll. mela-leuca* cfr. Rehm l. c.

10. **Tapesia atro-fusca** (B. et C.) Sacc. *Peziza atro-fusca* B. et C. 1537 North Am. Fungi n. 732. — Cupulis atro-fuscis, margine inflexo crenulato granulato nitido e subiculo ochroleuco membranaceo oriundis; sporidiis ellipticis binucleatis, 7-10 μ . long.

Hab. in ligno emortuo, Alabama Amer. bor. (PETERS).

11. **Tapesia epitheliohora** (Saut.) Sacc. *Peziza epitheliohora* Saut. 1538 Fl. Salzb. p. 9, Gonner t. 5, f. 3. Wint. Hedw. 1881, p. 131. — Pusilla; ceracea, sicca, sessilis, concava, cinerea, margine albido, disco nigrescente, subiculo thelephoroideo, gilvo; ascis cylindraceis, basim versus parum attenuatis, 35-45-4-5; sporidiis oblongo-fusoideis, inordinate stipatis, 6,5-7-1-1,5, hyalinis.

Hab. in ligno putri prope Mittersill Salzburg et in Fago S. Giovanni d'Andorno Ital. bor. (CESATI). — Ascoma $1\frac{1}{4}$ mm., subiculo flavo griseo, tomentoso insidens.

12. **Tapesia tela** (B. et C.) Sacc. *Peziza tela* B. et C. North Am. 1539 Fungi n. 731. — Subiculo expanso albido contexto, margine byssino; cupulis minutis concavis fuscis.

Hab. in ligno emortuo, in Carolina inf. Amer. bor. (RAVENEL).

— Affinis videtur *T. Bloxami*, sed non confluens.

- 13. **Tapesia culcitella** (C. et E.) Sacc. *Peziza culcitella* C. et E. 1540

Grev. VI, p. 7. — Gregaria; cupulis in strato brunneo nidulanti-
bus, hemisphæricis, demum applanatis, albidis, glabris; ascis cylin-
draceis; sporidiis linearibus, $8 \approx 2$.

Hab. in truncis vetustis quercinis New Jersey Amer. bor.

— Ascoma 1 mm. lat., initio lacteum.

14. **Tapesia fusca** (Pers.) Fuck. Symb. pag. 302, *Pez. fusca* Pers. 1541

Obs. II, p. 29 (adulta), *P. Pruni-avium* Pers. Obs. II, p. 82 (ju-
nior), *Mollisia fusca* Karst. M. F. I, p. 207, *Nipt. livido-fusca*
Fuck.? *Tapesia prunicola* Rehm, Sacc. et veris. etiam Fuck. —
Subiculo e filamentis fuscis dense contexto, late effuso, indetermi-
nato, fusco vel fusco-nigro, interdum parum visibili; ascomatibus
sparsis vel gregariis, primo urceolatis, nigrantibus ore albido,
connivente, dein explanatis, fusco-cinerescentibus aut cinereo-pal-
lescentibus aut subfuscis, disco pallido cinerescente v. albido-pal-
lente, margine tenui albicante v. disco concolore; interdum flexuoso
v. sublobato, basi ciliis radiantibus adfixis, glabris, 1 mm. diam.;
ascis fusoideo-clavatis, $55-68 \approx 6-8$, apice ultimo vix vel obsolete
jodo cœrulecente; sporidiis fusoideo-oblongatis vel fusoideo-elon-
gatis, simplicibus guttulatis, interdum spurie tenuiter uniseptatis
 $9-14 \approx 2-3$.

Hab. ad ligna et cortices arborum frondosarum et acerosarum
e. g. *Pini silvestris* putrescentium in Fennia, Britannia, Sue-
cia, Germania, Italia, Gallia, Hungaria, Belgio. — Forma **Corni**
Rehm Asc. n. 908: cupulis griseis, applanatis, undulatis, disco fla-
vido, margine albido; ascis 60-70 \approx 7-9; sporidiis 9-10 \approx 2,5-3, bi-
guttulatis, subiculi hyphis rectangulariter ramosis, septatis, brun-
neis, 5 μ . cr. In ramis Corni in Germania.

15. **Tapesia Rosæ** (Pers.) Fuck. Symb. mycol. pag. 301 et App. I, 1542

p. 48, *Peziza Rosæ* Pers. Obs. II, p. 28, Fr. Syst. Myc. II, p. 109,
Tymparis obtecta Wallr. var. *pezizæformis* Wallr. Flor. crypt.
II, p. 428, *Mollisia Rosæ* Karst. M. F. I, pag. 208. — Subiculo
late effuso, indeterminato, tenui, subtomentoso, fusco, hyphis brun-
neis 4 μ . cr.; ascomatibus sparsis v. confertis, subtomentosis aut
glabrescentibus, fuscis concavis v. concaviusculis, margine turgido,
siccis difformiter contractis, coriaceis, 1 mm. lat.; ascis cylindra-
ceo-clavatis, apice attenuato, $35-45 \approx 6-7$; sporidiis fusoideo-oblon-
gatis v. fusoideo-elongatis, 7-9 \approx 2-2,5; paraphysibus filiformibus.

Hab. ad caules et ramos Rosæ caninæ, alpinæ, tomentosæ, coriifoliæ aridos in Fennia, Lapponia, Suecia, Germania, Gallia, Britannia, Belgio, Amer. bor. — Variat sporidiis paullo longioribus (var. *tenebrosa* Karst. Monogr. Pez. p. 177), in caulis Rubi odorati.

6. **Tapesia Prunorum** (Fr.) Fuck. Symb. myc. App. I, p. 336, *Peziza Prunorum* Fr. S. v. Sc. p. 352 in nota. — Sessilis, concava, firmula, rufo-brunnea, extus glabra, demum applanata, subiculo strigoso atro, insidens; ascis et sporidiis maturis nondum visis (Fuckel).

Hab. ad cortices Prunorum in Suecia (FRIES) et Helvetia (MORTHIER). — Cupulis calvis a *T. prunicola* differt. «An, observante Frieso, eadem ac *Peziza atriseda* Saut. et *Peziza Pruni-avium* Pers.? Sec. Winter huc spectat *Pez. nitida* Saut. Salzb. pag. 8».

7. **Tapesia scariosa** (B. et C.) Sacc. *Peziza scariosa* B. et C. North Am. Fungi n. 733. — Cupulis tenuibus confluentibus collapsis nigris, margine albido, e subiculo nigro oriundis; ascis clavatis; sporidiis linearis-subfusiformibus.

Hab. in ramis emortuis in uliginosis in Amer. boreali. — Affinis *T. fuscæ*.

18. **Tapesia Torulæ** Fuck. Symb. myc. p. 302. — Cupulis minutis, usque ad 1 mm. latis, sessilibus, concavis, siccis margine albo, squamuloso, connivente, extus atris, disco albescente, fungo conidiophoro insidentibus; ascis oblongis, oblique stipitatis, 8-sporis, 40 \approx 6; sporidiis distichis, oblongis, 8 \approx 2-3.

Hab. in ramis adhuc pendulis, emortuis Salicis capreæ in silva Oestrich Rhenogoviæ. — Status conidicus, teste Fuck., est *Torula stilbospora* Corda subicum formans.

9. **Tapesia cinerella** Rehm Ascom. Alp. n. 16. — Plerumque gregaria, primitus subglobosa, sessilis, dein explanata, orbicularis, sicca concaviuscula, diaphano-cinerea vel flavidula, humida dilatata, plana, margine dilutiore, albido, tenui cincta, cinereo-hyalina, 1 mm. diam., contextu parenchymatico, basi subflavido, sessilis in byssò ex hyphis longissimis, simplicibus, septatis, fuscis, glabris, 3-5 μ . crassis composita; ascis clavatis, 8-sporis, 40-45 \approx 9, apice jodo cœrulecente; sporidiis obtuse ellipticis v. oblongis, rectis v. subcurvatis, 1-cellularibus, interdum nucleolis 2 præditis, hyalinis, distichis, 10-12 \approx 3,5; paraphysibus filiformibus, hyalinis, apice sensim 3 μ . incrassatis.

Hab. ad ramulos putridos Rhododendri ferruginei in alpibus tyrolensis, Peischl-See in monte Arlberg c. 1800 m. alt. (BRITZELMAYR) prope Kühtai (Oetz) c. 6200 (DR. REHM).

20. **Tapesia fusco-atra** (Hazsl.) Sacc. *Pyrenopeziza fusco-atra* Hazsl. 1547

Banat. pag. 58, t. II, fig. 10. — Mycelio hypodermio caules ambiente fusco, cupulis gregatim erumpentibus solitariis v. rarius fasciculatis in punctis mycelii densius textis, primum obovatis, demum patellaribus subsessilibus, fusco-nigris, margine crasso, crenulato albido, disco fusco concavo, $\frac{1}{2}$ mm. lato; ascis oblongo-linearibus, 110-125 μ . longis; paraphysibus paucis, apicibus crassioribus lanceolatis; sporidiis octonis, oblongo-linearibus, hyalinis, primum granulosis demum 2-guttatis, 25 \times 5-7.

Hab. in caulibus Tordylii maximi ad Soborzin Hungariæ.

21. **Tapesia apocrypta** Rehm Ascom. n. 861. — Solitaria, dispersa 1548

in maculis fuscis sessilis, primitus globosa, dein urceolata, ceracea, glabra, fusca vel fusco-nigricans, contextu parenchymatico e cellulis parvulis fuscis composito, versus marginem tenuem pseudoprosenchymerico ibique hyphis septatis, fuscis, obtusis, 4 μ . cr., apice subdistantibus, basi sessilibus, in hyphis plurimis longissimisque, intricatis, vix ramosis, septatis, fuscis, 2 μ . cr. insertis; disco subhyalino, 300 μ . diam.; ascis clavatis, apice obtuse acutatis et subincrassatis, 30-36 \times 4-5, 8-sporis; sporidiis fusiformibus vel oblongo-cuneiformibus, rectis, interdum curvatulis, 1-cellularibus hyalinis, 6-7 \times 1-1,5, distichis; paraphysibus filiformibus, haud clavatis, 2 μ . cr., hyalinis.

Hab. ad folia sicca farcta Saxifragarum juxta glaciem æternam montis Ortler Tyrolis. — Sicca oculo inermi vix visibilis.

22. **Tapesia balsamicola** Peck 34 Rep. p. 51, t. I, f. 14-21. — Su- 1549

biculo tenui adpresso, griseo, 2-4 mm. lato, filamentis dimorphis composito, aliis rugulosis, ramosis, septatis, atro-brunneis, breviramulosis, corpuscula sporomorpha 3-4-lobata, colorata, 15-22 μ . longa lataque gerentibus, aliis delicatis, hyalinis, conidia anguste fusoidea, hyalina gerentibus; cupulis minutis, 300-400 μ . lat., sessilibus, glabris, immarginatis, ceraceis, albidis, subpellucidis; ascis sursum dilatatis, apice obtusis, 40-50 μ . longis; sporidiis subfusoides, stipatis, 15-20 \times 5-7, 3-4-nucleatis; paraphysibus filiformibus.

Hab. in foliis vivis vel languidis Abietis balsameæ Stony Clove Catskill Mount. Amer. bor. — Cfr. Peck l. c.

23. **Tapesia aurea** Fuck. Symb. App. II, p. 60. — Cupulis gregariis, 1550
ceraceis, glabris, distincte stipitatis, in subiculo delicatissimo aureo,

hyphis ramosis contexto, conidiophoro ortis, primo cylindraceis, dein turbinatis, concavis, postremo subplanis, usque ad millimetrum latis, distinete marginatis, margine subrepando, aureis, disco pallidore, stipite pro ratione crasso, cupulam dimidiā aequante, basi obscuriore et subtilissime puberulo; ascis cylindraceo-clavatis, sessilibus, 8-sporis, $32 \approx 4$; sporidiis distichis, cylindraceis, parum curvatis, continuis, hyalinis, $6 \approx 2$, conidiis cylindraceis, rectis, endosporis duplo majoribus.

Hab. in acubus putridis *Pini silvestris*, rarius, pr. Eberbach Rhenogoviæ. — Subiculum serius evanescit, cupulæ nigrescunt.

24. **Tapesia strobilicola** Rehm Ascom. n. 761, ut var. — Sessilis, 1551 primitus globosa, dein urceolata, denique plana, crasse marginata extus fusca, disco flavo v. flavidо-pallido, 1-2 mm. lat.; subiculo late expanso, fusco, hyphis copiosis, longis, ramulosis, septatis, brunneis, $4-5 \mu$. cr.; ascis clavatis, 8-sporis, $45 \approx 6$; sporidiis distichis oblongo-ellipsoideis, obtusiusculis, rectis curvulisve, hyalinis, $6-8 \approx 2,5$; paraphysibus filiformibus.

Hab. in strobilis Pinorum in Germania. — Excipulum parenchymaticum, margine cellulæ subprosenchymaticis prominulæ. Cl. Rehm habet fungillum tamquam varietatem *Pezizæ melaleucæ* Fr., sed hæc subiculo caret et *Mollisia* potius videtur. Cfr. suo loco.

25. **Tapesia earina** Ellis Bull. Torr. 1881, p. 124. — Tenuis, molliuscula, mycelio laxo, sordide albo, gossypino, orbiculari, 2 mm. lato, fibroso contexto inserta; disco lateritio-rubro (sicco); ascis cylindraceis, $120-125 \approx 12$; sporidiis monostichis, ellipsoideis, hyalinis, saepius uninucleatis, $15 \approx 7$; paraphysibus filiformi-clavulatis, obtusis.

Hab. in culmis foliisque graminum lignoque putri sub nive in « Pleasant Valley » Utah Amer. bor. (HARKNESS).

26. **Tapesia mutabilis** (B. et Br.) Sacc. Phill. Disc. p. 278, *Pez. mutabilis* B. et Br. Ann. N. H. n. 564. — Minuta e macula seu subiculo minuto radiatim fusco-villoso erumpens, primum hemisphærica extus fusca, demum expansa concava, pallide albida glabrescens; sporidiis elongatis curvulis, $13-18 \approx 2-4$; paraphysibus filiformibus.

Hab. in foliis Airæ cæspitosæ, Derry Hill, Wilts Britanice. — Sicca Pyr. atratam in mentem revocat. Ascoma 200-300 μ . lat.

27. **Tapesia evilescens** (Karst.) Rev. mon. pag. 137, *Mollisia evile-* 1554

scens Karst. M. F. I, p. 208. — Subiculo tenui, effuso, cupulis planiusculis, fusco-cinerescentibus, margine albicante, disco cinerescente vel pallido, siccis difformiter contractis et obscurioribus 0,5-1 mm. lat.; ascis fusoideo-clavatis 45-5; sporidiis fusoideo-elongatis, guttulatis, 6-10-1-1,5.

Hab. ad culmos graminum putrescentes in Fennia australi.

B. Fructificatio adhuc incognita.

28. **Tapesia megaloma** (Schw.) Sacc. *Peziza megaloma* Schw. Syn. 1555 p. 175. — Subiculo late effuso, atro, contexto ex filis crassiusculis; cupulis majusculis, elevatis, subrotundis, ceraceis ex lutescente albidis, margine crasso involuto subrepando, glabris aut subfurfuraceis, disco cinereo.

Hab. in lignis putridis, passim, Bethlehem Amer. bor.

29. **Tapesia adhærens** (Wallr.) Sacc. *Pez. adhærens* Wallr. Fl. crypt. 1556 n. 2544. — Ascomatibus sessilibus sparsis carnosulis glaberrimis luteo-fulvis, e basi inde fila parca arachnoidea emittentibus, primum conniventibus dein urceolatis, margine inflexo concolore cinctis, demum deplanatis immarginatis.

Hab. ad ligna carie exesa salicina in Thuringia rarius.

30. **Tapesia luteola** (Fr.) Sacc. *Pez. luteola* Fr. S. M. II, p. 110. — 1557 Sessilis, sparsa, glabra, concava, luteola, basi tomento brevi albido insidens.

Hab. ad ligna abietina in Suecia et Germania. — *Tap. sanguinea* multo major et magis ceracea, mollier sed subicum ejusdem indolis, villus modo cupularum basin cingens. Cupulæ solitariæ, regulares, integerrimæ, luteo-flavæ.

31. **Tapesia mortuaria** (Ces.) Sacc. *Peziza mortuaria* Ces. in Kl. 1558 Herb. myc. II, n. 18 et in Bot. Zeit. 1855, pag. 596. — Subiculo tenui, nigro; cupulis minutis, sessilibus aggregatis, junioribus cinereo-virentibus cito nigris, semper glabris.

Hab. ad ligna? pr. Vercelli Italiæ superioris (CESATI).

32. **Tapesia atriseda** Sauter in Flora 1845, p. 133 (*Peziza*). — Ses- 1559 silis, cupulis glabris, planis, albidis, subiculo lato atro-tomentoso insidentibus.

Hab. in ligno udo vetusto, quod subinde obtegit, in agro urbis Stiriae. — Cupulæ 600 μ . ad 1 mm. latæ.

33. **Tapesia byssiseda** (Fr.) Sacc. *Pez. byssiseda* Fr. Obs. II, p. 309, 1560 Syst. Myc. II, p. 107. — Sessilis, planiuscula, testacea, extus mar- gineque granulosa, subiculo granuloso-villoso argillaceo.

Hab. ad ligna putrida, saepe humectata, Pin i in Suecia. — A reliquis admodum diversa. Subiculum latissime effusum, crustaceum, cohærens, magis a furfure conglomerato, quam e tomento contex-tum. Cupulæ subiculo fere immersæ, submembranaceæ, persistentes, juniores magis concavæ.

34. **Tapesia lateritia** (Pers.) Sacc. *Pez. lateritia* Pers. M. E. I, p. 1561 273, Fr. Syst. Myc. II, p. 110. — Sessilis, cupulis plano-concavis glabris nigris, subiculo dense lateritio immersis.

Hab. circa Parisios ad ligna rarissime. — Rhacodium refert ob tomentum crassum, cui cupulæ majusculæ sat profunde immersæ observantur. — Forma **Wallrothii**, cfr. Wallr. Crypt. n. 2555: asco-matibus sessilibus carnosis majusculis sparsis, extus fusco-furfura-ceis rugulosis, primum hemisphæricis dein deplanatis patellaribus, discum nigrum rugosum planum, margine reclinato conformi cinctum recludentibus stratoque hypostromatico fibroso-membranaceo flavo-lateritio insidentibus. Ad truncos fagineos putredine exesos nemoris roslebiens. (Wallr.).

35. **Tapesia dædalea** (Schw.) Sacc. *Peziza dædalea* Schw. Syn. p. 1562 174. — Crustacea, insidens subiculo pulverulento crustaceo, crassi-sculo, secedente, expanso in ligno et cortice, cupulis dense gregariis, majusculis sessilibus dædaleis, i. e. varie tortis et flexuosis, invicem adpressis, cum subiculo concolore, cinereo-pulverulentis; disco fuligineo-cinereo.

Hab. in cortice Lauri æstivalis, rara, Bethlehem Ameri-cæ borealis.

36. **Tapesia pruinata** (Schw.) *P. pruinata* Schw. Car. n. 1216, Fr. 1563 Syst. Myc. II, p. 109, *Pez. conspersa* Chaillet, nec DC. — Ses-silis, dense aggregata, membranacea, subclausa, cinereo-pulveru-lenta.

Hab. longe lateque profusa in corticibus emortuis Vitis, Li-quidambaris, Corni, Myricæ in Carolina sup. et inf. Amer. bor. et Gallia (?). — Subiculo tenuissimo insident cupulæ complanatæ, concavæ, subclausæ, minutæ, nigræ, omnino pulvere farina-ceo cinereo tam dense rectæ, ut sphærias credas; madidæ aperiuntur, ore inflexo, disco nigricante. Asci a Berkeleyo nulli inventi; an eadem ac *Arthonia confluens*?

37. **Tapesia conspersa** (Pers.) Sacc. *Thelebolus hirsutus* Dec. fl. fr. 1564 2, p. 272, *P. conspersa* Pers. Myc. Eur. p. 271 (ex syn.), Fr. S. M. II, p. 108. — Crustacea, sessilis, conferta, urceolata, fusco-ni-gra, pulvere albido obtecta, subiculo concolori.

Hab. ad cortices arborum prope Neufchatel, raro (KUNZE). Ill. Fries in indice S. M. III, p. 129 ad *T. pruinata* referre videtur.

— Primo obtutu Thelephoram frequentissime papillosam (*T. granulosam*) v. potius Lichenem refert subiculo late effuso, indeterminato, furfuraceo. Cupulæ innatæ, confertæ, subglobosæ, ore conivente, nigricantes, sed undique furfure tenui albido obrutæ; humectatæ vero magis apertæ, fuscescentes, ore albido. *Pyrenoteis* habitu proxima, sed vita non perennis neque cupulæ corneaæ quare vix Lichen.

38. **Tapesia decipiens** Wallr. Fl. crypt. n. 2487. — Ascomatibus sessilibus glaberrimis sparsis minutissimis rufo-fuscis, primum subglobosis dein planiusculis, margine concolore obtuso cinctis. 1565

Hab. ad caudices Pruni domesticæ, Helminthoporii hyphis in crustam furfuraceo-stygiam congestis insidens simulque stratum hypostromaticum filamentosum tenerimum mentiens.

39. **Tapesia griseo-pulveracea** (Schw.) Sacc. *Peziza griseo-pulveracea* Schw. Syn. p. 173. — Affinis *T. subiculata* sed subiculo araneoso multo tenuiori griseo, cupulis minoribus, extus nigris, margine tumidulo non lobato griseo-pulverulento; disco ochraceo etiam pulverulento. 1566

Hab. in ramis dejectis, Bethlehem Amer. bor.

40. **Tapesia subiculata** (Schw.) Sacc. *Peziza subiculata* Schw. Syn. 1567 p. 174. — Subiculo crasso, tomentoso, effuso, cinereo-fusco, densissime intertexto, cupulis glabris, marginatis et fibris quasi radiantibus, albis, sub nudo margine radiatim expansis, subiculo affixis, majusculis, in ambitu mire lobato-undulatis, 6 mm. lat., disco ochraceo.

Hab. passim in ramis dejectis, Bethlehem Amer. bor.

41. **Tapesia chlorotica** (Fr.) Fuck. Symb. p. 303, *P. chlorotica* Fr. 1568 Syst. Myc. II, p. 110. — Sessilis; cupulis planiusculis glabris rufescenti-nigris, subiculo tenui villoso viridi; ascis elongato-clavatis, stipitatis, 8-sporis; sporidii s subdistichis, oblongis, hyalinis, 5-6-guttulatis, 9-3.

Hab. ad ramos decorticatos dejectos virore obductos in Suecia, in ramis putridis Carpinii pr. Oestrich Rhenogoviæ. — Cupulæ minutæ, plane sessiles, gregariæ, præsertim disco rufescentes. Subiculum tenue, villosum, præcipue cupularum basin cingit.

42. **Tapesia? Bloxami** (B. et Br.) Sacc. *Peziza Bloxami* B. et Br. 1569 in Ann. Nat. Hist. n. 566. — Dense gregaria, mycelio albo inserta,

concava, pallide cervina, extus farinacea, disco concolore; ascis.... non visis.

Hab. in ramis dejectis pr. Twycross et Barmouth Britanniæ et in Carolina superiore Amer. bor. — Sec. Phillips, Disc. p. 408, asci nulli inventi, hinc species dubia.

43. **Tapesia Johnstonii** (Berk.) Phill. Disc p. 282, *Peziza Johnstonii* 1570 Berk. in Ann. Nat. Hist. n. 313. — Sessilis, globosa v. subturbinata, demum aperta v. rufa, sericeo-nitidula, subiculo lato, atro-brunneo, grumoso inserta; ascis.... non visis.

Hab. in ramis dejectis Berwick Britanniæ. — Ascoma 1 mm. lat., initio brunneum et pulvereum.

44. **Tapesia Messerschmidii** (Weinm.) Sacc. *Peziza Messerschmidii* 1571 Weinm. Hymen. Gaster p. 453. — Conferta, substipitata, carnosoceracea, extus cum stipite aureo-fulva, granulosa, intus dilutior, margine albida; stipite brevi.

Hab. in acubus Pini silvestris in Rossia. — Minuta sed pulchella et distincta species. Subiculum tenue, araneosum, primo aureo-sulvum, dein pallidius, irregulariter effusum; ascomata regularia, primo subglobosa, mox obconica cum stipite confluentia. Tota primitus aureo-fulvo, deinde margo albescit et discus dilutior evadit; adulta iterum toto anno fulva.

45. **Tapesia herpotricha** (Berk.) Sacc. *Peziza herpotricha* Berk. Dec. 1572 n. 304, Hook. Journ. 1851, p. 16, t. I, fig. 2. — Cupulis convexis, pallido-rufis demum obscurioribus glabris immarginatis, fibris ramosis repentinibus insidentibus.

Hab. in foliis viventibus, Caripi Brasiliæ (SPRUCE).

46. **Tapesia epitephra** (Berk.) Sacc. *Peziza epitephra* Berk. Fl. Tasn. 1573 pag. 275. — Minuta, alba, hemisphærica v. subglobosa, insigniter cava, floccis crispis fuscis intertextis oriunda.

Hab. in pagina inferiore foliorum in Tasmania (ARCHER). — Ascii ignoti, an ergo potius Solenia v. Cyphella.

II. *Eriopezia* Sacc. Ascoma pilosum v. puberulum.

A. Fructificatio nota.

47. **Tapesia cæsia** (Pers.) Fuck. Symb. mycol. p. 301, Sacc. F. it. 1574 f. 1431, *P. lichenoides* Pers. ic. et descr. pag. 29, t. 7, f. 1, *P. cæsia* Pers. Syn. pag. 657, Nees Syst. fig. 272, Dittm. in Sturm. Deutschl. fl. III, t. 31. — Sessilis, applanata, villosa, albida, basi pilis longis in subiculum intertextis, disco subtremellosa cæsio; ascis anguste lanceolato-oblongis, stipitatis, 8-sporis, 32-40 × 4-5;

sporidiis subdistichis, cylindraceis, curvatis, continuis, hyalinis, 3,5-4 \approx 1,5-2; paraphysibus filiformibus.

Hab. ad ligna dejecta quercina faginea etc. in Suecia, Germania, Gallia, Belgio, Britannia, Carolina. — Variat sec. Fuck.: sporidia paullo majora, 8 \approx 3. Cupulæ confertæ, juniores concavæ, dein explanatæ, plures in membranam villosam ob pila basin cingentia intertexta.

48. **Tapesia Chavetiæ** (Lib.) Fuck. Symb. myc. p. 301, *Peziza Chavetiæ* Lib. Exs. p. 26. — Gregaria, sessilis, minuta, membranacea, hemisphærica, concava, albo-tomentosa, basi pilis longis, in subiculum arachnoideum, album abeuntibus; disco subtremeloso, brunneo-atro; ascis oblongo-clavatis, 8-sporis, 24 \approx 4; sporidiis distichis, cylindraceis, curvatis, 5 \approx 2.

Hab. ad ligna putrescentia in Germania, Belgio, Gallia et Britannia. — Teste Phillipis Disc. p. 277 non satis differt a *Tapesia cæsia*.

49. **Tapesia erratica** (Fr.) Fuck. Symb. myc. p. 301, *Peziza erratica* Fr. Syst. Myc. II, p. 108. — Sessilis, granuliformis villosa, alba, basi pilis longis liberis cincta, disco ceraceo-atro, initio subpunctiformi dein magis dilatato; ascis et sporidiis ut in *Tapesia Chavetiæ* (teste Fuckel).

Hab. ad truncos dejectos Quercus in Suecia et Rhenogovia. — *Tapesia cæsia* peraffinis, differt subiculo densiore et cupulis magis applanatis sessilibus: siccitate flavescit.

50. **Tapesia fusco-umbrina** (Fr.) Sacc. *Tap. variicolor* v. *fusco-umbrina* Fuck. Symb. p. 302, Sacc. Mich. I, p. 65, F. ital. f. 1432, *P. variecolor* C, Fr. Syst. Myc. II, p. 100. — Cupulis scutellariibus, sessilibus, plerumque tapeto conidiophoro (*Tricholeconii fusci* Corda), insidentibus, extus fusco-griseis villosulis, præsertim ad marginem pilis simplicibus, septatis, fuligineis, 120-130 \approx 3, sursum asperulis, pallidis (hyphis analogis) ornatis; disco flavido; ascis cylindraceo-clavatis, 70 \approx 6-6 $\frac{1}{2}$, p. s. 50 long.; paraphysibus (spuriis?) filiformibus obvallatis, octosporis; sporidiis distichis, fusoideis, curvulis, 8-10 \approx 1 $\frac{1}{2}$ -2, nucleo medio bipartito ideoque spurie 1-septata, hyalina.

Hab. ad lignum populinum etc. putre in Suecia, Germania, Italia. — Quoad st. conid. cfr. Michelia l. c.

51. **Tapesia prolifica** Ellis Torr. Cl. 1881, p. 73. — Gregaria, sessilis, minuta, pilis cinereis, septatis, rugulosis, patulis dense obsessa, disco cupulato, atro, margine integro; ascis clavatis, p. sp.

25 μ . long.; sporidiis distichis, cylindraceis, curvulis, obtusis, subhyalinis, 7-8 \times 2,5.

Hab. in disco ligni quercini sicci, New Jersey Amer. bor. —

Pili sub microsc. brunnei, oculo inermi cinerei.

52. **Tapesia Rehmiana** Sacc. Bomm. et Rouss. Contrib. Mycol. Belgique 1579 II. p. 3 et in Rehm Ascom. n. 860. — Sessilis, minima, 175-226 μ . diam., non superans olivacea, margine pilis brevibus, hyalinis, fasciculatis, discum nonnihil superantibus instructa; disco subconcavo, olivaceo-livido; ascis subsessilibus, clavatis, 8-sporis, 30-40 \times 5-6, filiformi-paraphysatis; sporidiis oblique monostichis, ellipticis, interdum subcurvatis, hyalinis, 2-guttatis, 6-7 \times 3.

Hab. ad truncos decorticatos, putridos querneos, Britsfort Belgii (BOMMER et ROUSSEAU). — Subiculum, in quo ascomata sparsa vel conferta nidulantur, crassum, filamentis dematioideis erectis, continuis, obtusis, brunneis, 300 \times 6 formatum.

53. **Tapesia vinacea** (Rabenh.) Sacc. *Peziza vinacea* Rabenh. Hedw. 1580 1878, p. 31. — Siccando fuscescens, cupularis, sessilis, primitus basi byssio candida postea evanida cincta, cupula patelliformis 2-3 mm. lata (rarius supra), extus puberula subnuda, margine recto, sicco plerumque involuto, subtilissime crenulato; ascis cylindrico-tenuioribus, angustis, jodo non tinctis; sporidiis monostichis, ellipticis, hyalinis, 12-15 \times 8-9, episporio hyalino, tenui; paraphysibus numerosis, crassiusculis, apice vix incrassatis, non coloratis.

Hab. in vinaceis dejectis et putrescentibus in horto bot. Neapolitano (CESATI). — An potius *Pyronema* v. *Lachnea*?

54. **Tapesia prunicola** Fuck. Symb. p. 302, *Tympanis obtexta* Wallr. 1581 Cryp. p. 428 (pr. p.). — Cupulis duplo majoribus quam in *T. Rossæ*, aterrimis et villo magis atro-fusco; ascis oblongis, stipitatis, 8-sporis, 54 \times 5; sporidiis monostichis, cylindraceis, utrinque obtusis, rectis, 10 \times 2, hyalinis; paraphysibus filiformibus, non clavatis.

Hab. in ramis corticatis Pruni spinosæ pr. Oestrich Rheno-goviæ. — *Tap. prunicola* Rehm, Sacc. (præcipue in Alno) a *T. fusca* distingui nequit; an et typus Fuckelii qui inter villosas enumeratur?

55. **Tapesia eriobasis** (Berk.) Phill. Disc. p. 278, *Peziza eriobasis* 1582 Berk. Ann. Nat. Hist. n. 312. — Gregaria sed plerumque discreta, orbicularis, tomentosa, areola rotunda minuta nivea, gossypina sæpe confluente inserta, viva alba, disco in sicco flavido; ascis subclavatis v. obtuse fusoideis; sporidiis oblongis, 5-6 μ . longis.

Hab. in pariete interiori levi corticis in Britannia. — A *Tapesia cæsia* et *Chaveliae* differt cupulis majoribus, disco pallido etc.

56. **Tapesia pileo-crocata** (Cr.) Sacc. *Peziza pileo-crocata* Cr. Fin. 1583 p. 51. — Ascomate mycelio arachnoideo, albo inserto, 1-2 cm. diam., hemisphærico, margine undulato, pilis fasciculatis luteis tecto; disco albescente; ascis clavatis, octosporis; sporidiis oblongis; paraphysibus capillaribus, ascis longioribus.

Hab. in cortice ramuli emortui Betulæ atque ad basim caulis Angelicæ, Finistère Galliæ.

57. **Tapesia Corni** Fuck. Symb. p. 301. — *T. Rosæ* affinis; cupulis 1584 gregariis, sessilibus, villo olivaceo insidentibus, siccis clausis, globosis, humectatis apertis, 2 mm. latis, extus verticaliter plicatis (ut octogonæ appareant), brevissime olivaceo-pilosus, ore connivente, margine candide ciliato, disco concavo, pallide cinereo; ascis oblique stipitatis, oblongis, 8-sporis, 48 × 8; sporidiis monostichis, oblongo-ovatis, 10 × 5, hyalinis.

Hab. ad ramos siccos Corni albæ, raro, Reichartshausen Rhenogoviæ.

58. **Tapesia albo-viridis** (Gill.) Sacc. *Lachnea albo-viridis* Gill. Disc. 1585 p. 91. — Ascomate sessili, primo clauso, dein aperto, parum ventricoso, pilis tenuissimis fasciculatis tecto, citrino v. luteo-aureo v. fulvo, 4-5 mm. diam., subiculo tenui albescente, fere membra naceo; ascis cylindraceis; sporidiis ellipsoideis; paraphysibus filiformibus.

Hab. in foliis delapsis et in pericarpiis *Hederæ* et *Quercus* in Gallia.

B. Fructificatio adhuc ignota.

59. **Tapesia discincola** (Schw.) Sacc. *Peziza discincola* Schw. Syn. 1586 p. 174. — Minuta aggregata, in subiculo tenuissimo, albescenti; cupulis turbinatis, substipitatis, laxe stipatis, pallide ferrugineis, extus omnino tectis, pilis albido-cinereis, pulverulentis, primo conniventibus, demum irregulariter expansis; disco pallido.

Hab. in disco cæsi trunci et ad ligna Bethlehem Amer. bor.
— *Habitus Soleniae anomala*.

60. **Tapesia arachnoidea** (Schw.) Sacc. *Peziza arachnoidea* Schw. 1587 Syn. p. 174. — Subiculo ex filis arachnoideis longe lateque expansis tenerrimis, intertextis, subcandido-cinereis; cupulis pallido-aquosis, extus villoso-albis, minutissimis, siccitate globoso-contractis, fere imperceptibilibus.

Hab. in lignis dejectis udis, Bethlehem Amer. bor.

61. **Tapesia scutelliformis** (Wallr.) Sacc. *Pez. scutelliformis* Wallr. 1588

Fl. crypt. n. 2554. — Ascomatibus sessilibus, submembranaceis tenacellis, sparsis, extus tegmine furfuraceo terso fusco-nigrescente nitidulo vestitis, e basi inde telam arachnoideam exserentibus sensimque in stratum stipato-intertextum membranaceum spadiceum nitidum subcrescentibus, primum globosis vertice hiantibus, dein reclinando discum planum stramineo-pallentem, margine inflexo flexuoso cinctum recludentibus, majusculis, adpressis, scutellaribus.

Hab. ad ligna abietina mausolei cujusdam in agro Northusano (WALLR.). — Stratum hypostrom. dein membranulæ continuæ instar cubilia undique obducit et siccando integrum secedit. Ascomata *P. melaxantham* s. *xanthomelam* haud male referunt.

62. **Tapesia albo-olivacea** (Gill.) Sacc. *Lachnea albo-olivacea* Gill. 1589

Disc. p. 92. — Ascomate sessili, applanato, extus tomentoso et albicante, 1-1 $\frac{1}{2}$ mm. diam., margine prominulo albo, primo regulari, dein undulato; disco primo olivaceo, dein brunneo-olivaceo, subiculo albo copioso.

Hab. in fragmentis ligneis, locis humidis Galliæ super.

63. **Tapesia candido-fulva** (Schw.) Sacc. *Peziza candido-fulva* Schw. 1590

Syn. p. 174. — Subiculo araneoso, satis denso, longe lateque effuso, candidissimo, cupulis eo sparse insertis, 2-4 mm. lat., elegansissime fulvis, extus pilis brevibus subfasciculatis fusco-fulvis ornatis.

Hab. rara sed per pulchra in cortice, Bethlehem et Carol. inf. Amer. bor. — Affinitatem cum *T. fulvo-cana* habet sed differt subiculo. Sec. Berk. habitus *Belonidii Aureliæ*, sed sporidia non visa.

64. **Tapesia stipata** (Fr.) Sacc. *P. stipata* Fr. Syst. Myc. II, p. 106, 1591

Summa p. 352, nec Pers. — Confertissima, confluens, sessilis, villosa, griseo-fusca, margine tenui, disco urceolato nigrescente.

Hab. ad ligna putrida Fagi, Tiliæ in Suecia. — Subicum fere nullum, sed cupulæ adeo dense aggregatæ, ut primo obtutu *Polypori* speciem omnino referant. Cupulæ oblongæ, leviter ventricosæ, etiam basi villosæ.

65. **Tapesia fibrillosa** (Wallr.) Sacc. *Pez. fibrillosa* Wallr. Fl. Crypt. 1592

n. 2541, *Peziza ferruginea* Pers. Myc. I, 274, tab. 1, f. 7,8 nec Schumach. — Ascomatibus sessilibus, hemisphæricis, vellere furfuraceo tenui ferrugineo vestitis, e basi telam hypostromaticam stygiam teneram exserentibus, demum vertice in discum concavum dilutiorem, marginibus inflexis cinctum reclusis.

Hab. ad ramulos quercinos vallum Hercyniæ infer. udarum.

66. **Tapesia erysiphoides** Rabenh. DC. Fl. p. 357, n. 2752. — Gre. 1593

garia, sessilis, primo clausa, villosa, subobovata, obtusa, demum aperta, depressa, 2-5 mm. lata, profunde cupulata, ex olivaceo-viridi brunneola, margine involuto, nigrescente; disco brunneo, demum subsulcato; subiculo araneoso molli, albido, radiatim extenso.

Hab. ad lapides arenosas in Helvetia saxonica, sociâ *Schistostegâ osmundacea*.

DESMAZIERELLA Lib. Ann. Sc. nat. 1829, p. 82 c. ic., Phill. Disc.

p. 283 (Etym. a mycologo acuto et sollerti J. B. DESMAZIÈRES). —

Ascoma sessile v. subsessile, cupulatum, disco paraphysium apicibus emergentibus atris hirtello. Asci cylindracei, octospori. Sporidia ellipsoidea, hyalina.

1. **Desmazierella acicola** Lib. Crypt. Ard. n. 24, Phill. Disc. p. 283, 1594 t. 8, f. 51. — Sparsa, sessilis, turbinata, dein plana, extus nigra et tomento nigro byssideo obsessa, margine setulis longis, nigris ciliato; disco atro-brunneo, paraphysibus exilientibus hirtello; substantia pallide cinerea; ascis cylindraceis; sporidiis ellipsoideis, bигuttatis, 15-8, paraphysibus in fasciculos coalescentibus, sursum discretis, apice acuto nigricante.

Hab. in acubus emortuis Pini silvestris in Arduennis, Germania et Britannia. — Ascoma 5-6 mm. lat., initio chætomiiforme.

PIROTTÆA Sacc. Mich. I, p. 424 [1878] (Etym. a cl. prof. R. PIROTTA, botanico ital.). — Ascomata superficialia v. erumpenti-superficialia, sessilia, scutellata, subcarbonacea, nigra, setulis rigidis plerumque septatis vestita, disco levi. Asci octospori, paraphysati. Sporidia oblongata, continua hyalina. — Habitus *Pyrenopezizæ* sed ascoma setigerum.

A. *Eu-Pirottæa*. Sporidia breviuscula.

1. **Pirottæa veneta** Sacc. et Sp. Mich. I, p. 424, F. it. f. 1395, — Asco. 1595 matibus subsuperficialibus, scutellatis, sessilibus, $\frac{1}{6}$ - $\frac{1}{5}$ mm. diam., aterrimis, in sicco subclausis, setulis cuspidatis, 40-50 x 5-7 parce septatis, obscure fuligineis (ad marginem crebrioribus) vestitis; excipuli contextu distinete parenchymatico, atro, cellulis facile secendentibus; ascis cylindraceo-clavatis, 40-45 x 8, apice obtusiusculis, subsessilibus; paraphysibus bacillaribus obvallatis, octosporis, apice jodo cœruleacente; sporidiis cylindraceis, utrinque rotundatis, 14-15 x 2-2 $\frac{1}{2}$, 2-guttulatis, hyalinis.

Hab. gregatim in caule, rarius foliis emortuis Hellebori viridis et Adenostyli dis, Conegliano Ital. bor., in Tyrol. et Belgio.

2. **Pirottæa gallica** Sacc. Mich. II, p. 82 et 59, F. ital. f. 1394. — 1596
 Ascomatibus gregariis, superficialibus, aterrimis, $\frac{1}{5}$ mm. diam., concavo-scutellaribus, sessilibus, siccis semiclausis, ubique setulosis; peritheci contextu parenchymatico, rigidulo, fuligineo, cellulis superficialibus setulisque cuspidatis 2-3-septatis, $30 \approx 5$, obscurioribus facileque secedentibus; ascis cylindraceo-clavatis, $35-45 \approx 5-7$, breve stipitatis jodii ope flavescentibus; paraphysibus filiformibus obval-latis, octosporis; sporidiis cylindraceo-clavulatis, $7-9 \approx 1\frac{1}{2}$, minute 2-guttulatis, hyalinis.

Hab. in caulibus emortuis Cichorii Intybi, Aconiti, Adenostyli dis, Senecionis in Gallia, Italia, Belgio, Allgau, Tyroli. — A *P. veneta* S. et S. differt præcipue sporidiis dimidio minoribus etc.

3. **Pirottæa Bresadolæ** Sacc. Mich. II, p. 536, F. ital. fig. 1393, *Mollisia cæsiella* Bresad. F. trid. II, p. 25, t. XXX, f. 1 (icon justo pallidior). — Erumpenti-superficialis, fusca, disco expallenti, sed non vere cæsio; contextu excipuli fuligineo-atro, parenchymatico, setulis marginalibus acicularibus, $50-60 \approx 5$, sæpe 1-septatis fuligineis, subinde fasciculatis; ascis tereti-clavatis, breve stipitatis, $100 \approx 10-12$, filiformi-paraphysatis, octosporis; sporidiis distichis, tereti-clavulatis, $23-27 \approx 3-3,5$, leniter curvis, 3-4-guttatis, hyalinis.

Hab. in caulibus herbarum majorum e. g. *Cirsii palustris* in agro tridentino (BRESADOLA). — A *Pir. veneta* differt ascomatibus majoribus, suberumpentibus, ascis sporidiisque multo amplioribus, cellulis excipuli non secedentibus, sed ob contextum fuligineo-atrum setulasque evidentissimas vera *Pirottæa*.

4. **Pirottæa cembricola** Rehm Ascomyc. fasc. XVI, p. 6, n. 763. — 1598
 Ascomatibus gregariis, primitus orbicularibus, dein explanatis, patellaribus, sessilibus, $0,5-1,5$ mm. diam., siccis difformibus, subglobosis, urceolatis, disco flavidulo, humidis diaphanis, subfimbriatis, contextu parenchymatico fusco, margine (interdum etiam versus basim) fimbriis septatis, fuscidulis, obtusis, $36-40 \approx 5-6$ crebris obsessis; ascis clavatis, basi lata, sessilibus, 8-sporis, $40 \approx 7$; sporidiis distichis, continuis hyalinis, oblongis, obtusis, rectis v. subcurvatis, $8-9 \approx 2,5-3$; paraphysibus filiformibus haud clavatis, obtusis, ascos superantibus, $0,3 \mu$. crassis; hyphis basalibus longis septatis, simplicibus, fuscis 3μ . crassis..

Hab. ad ramos decorticatos putridos Pini Cembræ supra Sulden alpium tyrolensium. — Hyphæ basales longæ, septatæ, simplices, 3 μ . cr. fuscae.

5. **Piottæa frondicola** (E. et E.) Sacc. *Lachnella frondicola* E. et 1599
E. Jour. Myc. 1888, p. 99. — Sparsa, sessilis, cupuliformis, nigra,
sicca fere clausa, $1\frac{1}{4}$ - $1\frac{1}{3}$ mm. diam., disco pallido, extus marginæ
que pilis brevibus, brunneis, obtusis, septatis, sursum parum ru-
gosis, 35±3, tecta; ascis oblongo-lanceolatis, sessilibus, utrinque
subacutis, 30-35±6; paraphysibus filiformibus; sporidiis distichis,
clavato-oblongis v. subfusoideis, hyalinis, nucleatis, 8-12±1,5.

Hab. in frondibus delapsis Osmundæ? Newfield Amer. bor.

6. **Piottæa mimatensis** Pass. et Roum. Rev. Mycol. 1885, p. 172. — 1600
Gregaria, minuta, atra, superficialis, scutellata, concava, sessilis,
extus aculeata, aculeis facile secedentibus, simplicibus, fuligineis,
28-45 μ . longis; sporidiis acicularibus hyalinis; paraphysibus fili-
formibus, ascos paullo superantibus.

Hab. in foliis semivirgentibus Pini silvestris in territorio
Mimatensi «Fôret de Mercoire» Galliæ (PROST).

7. **Piottæa venturioides** Sacc. et Rom. n. sp. — Sparsa sessilis v. bre- 1601
vissime stipitata, uda aperta, urceolata v. subsphæroidea, minuta,
150-180 μ . diam., undique nigro-setulosa, setis rectis subulatis fusco-
opacis, 50-270±4-6, primo obtutu ob setulas subfusca, verum al-
bida, disco hyalino pallido, margine albo minutissime crenulato,
sicca clausa sphæroidea nigra; ascis tereti-oblongis v. subclavatis,
30-42±5-8; sporidiis acicularibus, hyalinis, continuis, 6-11±1,5-
1,7; paraphysibus filiformibus, 1 μ . cras.

Hab. in foliis putridis dejectis Vaccinii uliginosi in Ne-
ricia Sueciæ. — Habitus *Venturiae Myrtilli*.

8. **Piottæa uliginosa** Karst. Symb. Myc. Fenn. XXIII, p. 5. — Sparsa,
planiuscula, setulis strictis, articulatis, nigris, 100-180±6, obsessa,
minima, contextu minuto, parenchymatico hyalino; ascis cylindra-
ceis, apice leviter attenuatis, 34-42±5; sporidiis.... non satis evo-
luti visis; paraphysibus filiformibus, parcis, 1 μ . cr.

Hab. in foliis putrescentibus Calamagrostidis lanceola-
tæ, Mustiala Fenniæ. — Contextu hyalino a typo desciscit.

9. **Piottæa erubescens** Rehm Ascom. n. 412. — Primitus globosa, 1602
dein urceolata, gregaria, sessilis, extus fuscorubra, ore albido, sicca
globoso-contracta, extus rugulosa, humida urceolato-explanata, di-
sco concavo, pallide flavidulo, contextu parenchymatico fusco, ad
marginem prosenchymatico, cellulis elongatis, fere filiformibus, p.

p. hyalinis et fuscis, distantibus; ascis clavatis, 8-sporis, 75 ± 8 ; sporidiis oblongis, obtusis, nucleolis 2 præditis, rectis, hyalinis, plerumque 1-stichis, $12-15 \pm 4,5$; paraphysibus lanceolatis, acutis ascos superantibus, c. 5. μ . cr.

Hab. ad caules emortuos Eupatorii cannabini prope Augustam Vindelicorum seu Augsburg (BRITZELMAYR).

B. Mancinia Sacc. *) Sporidia prælonga, filiformia.

10. **Pirottæa strigosa** (Karst.) Rev. mon. p. 138, *Lachnum strigosum* Karst. M. F. I. pag. 183. — Sparsa v. subgregaria, sessilis, adnata, sphæroideo-applanata, nigra, strigosa, disco rufescente, 0,5-0,7 mm. lat.; ascis cylindraceo-clavatis, $60-74 \pm 8$, jodo non cœrulencentibus; sporidiis fusoideo-filiformibus, guttulatis, rectis, 30-42 $\pm 1,5$; paraphysibus 1,5-2 μ . cr., sursum 2-3 μ . cr., apice acutiusculo.

Hab. in caulis Achilleæ Millefolii aridis prope Mustiala Fennia. — Pili nigri (sub microsc. fusi), $120-250 \pm 6-8$, apice attenuati. Spermogonia forsitan ejus sistit *Dinemasp. strigosum*. Vergit ad Mollisiam. *P. Vecti* proxima.

11. **Pirottæa Vectis** (B. et Br.) Phill. Disc. p. 284, t. 8, f. 52, *Peziza Vectis* Berk. et Br. Ann. Nat. Hist. n. 957. — Minuta, sparsa, sessilis, subglobosa, subdepressa dein hemisphærica, extus obscure brunnea v. subnigra, pilis brevibus brunneis rigidis obsessa, disco concavo pallido; ascis tereti-clavatis; sporidiis e filiformi fusoideis curvulisve vermicularibus, $25-30 \pm 4$, (spurie) 1-3-septatis; paraphysibus filiformibus.

Hab. in caulis emortuis Centaureæ nigræ, Ryde in ins. Wight (Vectis) Britanniæ (BLOXAM). — Ascoma 300 μ . lat. Setulæ plerumque marginales, granulis polygoniis brunneis intermixtæ.

C. Species fructificatione ignota.

12. **Pirottæa? aterrima** (Lasch) Sacc. *Peziza aterrima* Lasch in Klotzsch Herb. Myc. n. 336 et in Rabenh. Krypt. Fl. I, p. 359 nec Rehm, nec *Peziza corvina* Fr. Syst. Myc. II, p. 97. — Gregaria, sessilis, globoso-hemisphærica, obtuse marginata, nigra, setulis patulis? extus vestita.

Hab. in acubus Pinorum putridis prope Driesen Germaniæ.

13. **Pirottæa? tentaculata** (Wallr.) Sacc. *Pez. tentaculata* Wallr. 1607

*) Collaboratori optimo, prof. VINC. MANCINI dicatum subgenus.

Fl. crypt. n. 2463. — Ascomatibus sessilibus parvis subrotundis atris carnosso-membranaceis, primum conniventibus, dein marginem obtusum nitidulum inflexum reclinantibus discumque planum opacum concolorem recludentibus, patellaribus, extus margineque setis brevibus stipatis tentaculato-porrectis nigris sparsis parce instructis.

Hab. ad virgas salicinas decorticatas Thuringiæ rarius: — *Lecanid. atrati* habitum exacte reddit, extus tamen setis brevibus stipatis, tentaculorum instar porrectis munita certo dignoscitur.

14. **Pirottæa? corvina** (Pers.) Sacc. *P. corvina* Pers. S. M. I, p. 248, 1608

Fr. Syst. Myc. II, p. 97. — Sessilis, minuta, subdepressa, externe pilosa, utrinque atra.

Hab. ad ligna exsiccata indurata, passim hieme, sed videtur perennis. — Gregaria, subiculo maculæformi nigricante plerumque insidens, e ligno suberumpens, primo punctiformis, dein late aperta. Non cum simili *Cœlosphæria exili* commutanda; potius *Excipulæ* species.

15. **Pirottæa? Hydrangeæ** (Schw.) Sacc. *P. Hydrangeæ* Schw. Carol. 1609 n. 1219, Fr. Syst. Myc. II, p. 109. — Sessilis, subglobosa, pilis nigris longis involuta, cupulis coriaceis cinereo-albis, extus nigris pilosis.

Hab. rario ad truncos emortuos Hydrangeæ in Carolina Amer. bor. — Primum clausæ et involutæ, deinde aperiuntur cupulæ, margine infexo; etiam interdum clausæ manent, ore tamen conspicuo.

LACHNELLULA Karst. Medd. af Soc. pro Fauna et Fl. Fenn. 11, pag. 138 [1884] (Etym. demin. *Lachnellæ*). — Ascoma sessile vel substipitatum minutum, cupulatum vel scutellare, ceraceum, extus pilosum. Asci cylindracei, octospori, paraphysati. Sporidia minuta globosa, hyalina. — Est *Trichopeziza* v. *Lachnella* sphærospora.

1. **Lachnellula chrysophthalma** (Pers.) Karst. l. c. et Rev. monogr. 1610 p. 132, *Peziza chrysophthalma* Pers. Myc. Eur. I, pag. 259 nec Ger., *Helotium* Karst. M. F. I, p. 155, *Pitya suecica* Fuck. Symb. App. III, p. 32, *P. calycina* β. *Abietis* Fr. Syst. Myc. II, p. 91, pr. p. Karst. Pez. et Asc. p. 16 pr. p. *Pez. pulchella* b. *flavococcinea* Alb. et Schw. Conspl. Fung. nisk. p. 323? — Gregaria v. sparsa substipitata, albo-villosa vel tomentosa, explanata, sicca concava, disco aurantiaco-luteo v. aurantiaco, 2-4 mm. lat., excipulo e fibrillis ramosis, laxis, hyalinis composito; pilis asperulis, hyalinis, simplicibus, obtusis, 40-4; stipite brevi vel brevissimo,

in cupulam dilatato; ascis subcylindraceis, 50-60 = 6-7; sporidiis monostichis, sphæroideis, 4-5 μ . diam.; paraphysibus gracilibus.

Hab. ad corticem et lignum Abietis excelsæ, Pini Cembrae, P. pumilionis, Laricis in Fennia, Lapponia rossica, Germania, Hungaria, Helvetia.

2. **Lachnellula calycina** (Vuill.) Sacc. *Trichoscypha calycina* (Schum. e. p.) Vuillemin Bull. Soc. Bot. Franc: 1888, LXX, *Pez. calycina* Schum. e. p. sec. Vuill. — Ascomatibus sparsis v. cæspitosis, 0,8-2 mm. diam., udis cupulatis, siccis vel in juventute contractis, basi in stipitem brevem (subinde nullum) desinentibus, extus albo-tomentosis, disco plano v. leviter concavo, pallide aurantiaco; ascis elongato-clavatis, 35-40 = 3, octosporis; sporidiis sphæroideis, monostichis, 2 μ . diam., levibus; paraphysibus filiformibus septatis.

Hab. in cortice vivo Pini silvestris, P. Strobi, rarius Abietum in Vogesis.

3. **Lachnellula theioidea** (C. et E.) Sacc. *Peziza theioidea* C. et E. Grev. VII, p. 7. — Sparsa, gilva; cupulis subglobosis, dein hemisphæricis, leniter tomentosis, pulvere sulphureo ubique conspersis; ascis cylindraceis; sporidiis globosis, 3 μ . diam.

Hab. in ramis decorticatis Rhois venenatæ, New Jersey Amer. bor. — Matrix subinde granulis sulphureis, a cupulis delapsis, conspersa.

4. **Lachnellula hysterigena** (B. et Br.) *P. hysterigena* B. et Br. 1613 Fung. of Ceyl. n. 947. — E cupuliformi applanata, margine inflexo, extus albida, furfuraceo-tomentosa; hymenio cinereo; ascis clavatis; sporidiis minutis globosis, 3,7-4 μ . diam.

Hab. parasitica in Hysteriis, Central province Ceylon.

5. **Lachnellula cyphelloides** (E. et E.) Sacc. *Peziza cyphelloides* E. et E. Journ. Mycol. I, p. 151 (1885). — Subsessilis, minuta, $\frac{1}{4}$ - $\frac{1}{3}$ mm. lat., mollis, alba, pubescens, margine incurvo, contextu tenui, celluloso; ascis teretibus, fasciculatis, 24-30 = 1,5 p. sporif. 15 μ . long., octosporis; sporidiis globosis 1 μ ; paraphysibus nullis.

Hab. in aculeis Hydnæ (membranacei?) Newfield N. J. Amer. bor. — Socium adest *Penicillium glaucum*, cuius conidia illa *Lachnellulæ* imitantur. Habitus *Cyphellæ*, sed, teste Ellisio, asci genuini adsunt.

LACHNELLA Fr. Summa p. 365 (1849), Fuck. Symb. p. 280 (Etym. demin. a *Lachnea*) *Peziza* sect. *Dasyse.* et *Fibrina* Fr. e. p. — Ascoma sessile v. brevissime stipitatum, urceolato-scutellare, extus

villosum, ceraceo-coriaceum. Asci elongati, 4-8-spori. Sporidia te reti-oblonga v. ovoidea, continua, hyalina. Paraphyses filiformes. — A *Trichopeziza* præcipue differt ascomate coriacello et paraphysibus sæpius filiformibus.

1. **Lachnella flammea** (A. et S.) Fr. Summa p. 365, *P. flammea* A. 1615 et S. p. 319, t. 11, f. 7, Pers. Myc. Eur. p. 252, Fr. Syst. Myc. II, p. 96, *Helotium* Karst. — Sessilis, subglobosa, firma, strigoso-villosa, unicolor flammeo-rubiginosa, pilis simplicibus, fuscis, obtusis, sublevibus circ. 4-5 μ . crass.; ascis clavatis, 75-80 \approx 9-10, 8-sporis; sporidiis oblongis, obtusis, rectis v. subcurvatis, 1-cellularibus, hyalinis, 10-15 \approx 3- $3\frac{1}{2}$, distichis; paraphysibus filiformibus, apice haud clavatis.

Hab. ad ramos vetustos induratos nudos Piri Mali, Carpinii, Liquidambaris, Salicis in Suecia, Fennia, Germania, Gallia, Britannia, Italia, Hungaria, Amer. bor., Ceylon. — Contextus excipuli prosenchymaticus; pili sæpe ob crystalla minuta rugulosi, 3-4 μ . cr. Matrix circa fungum rufo-brunnea. Proximis utique cognata loco et substantia; etiam caro interna nigra. Primo tota clausa, globosa, subrugosa, dein patens, cupularis, hemisphærica, regularis, 2-3 mm. æquans. Hymenium vivide rubiginosum.

2. **Lachnella barbata** (Kunze) Fr. Summa p. 365, *Peziza* Kunze in 1616 Fr. Syst. Myc. II, p. 99. — Sessilis, globoso-applanata, connivens, firma, rugoso-tomentosa, ferruginea, disco pallido, pilis tereti-clavatis, septatis, fuligineis, 3,6-5 μ . cr.; ascis cylindraceis, 80-90 \approx 5-7, octosporis; sporidiis oblique monostichis, tereti-fusoideis, 8-10 \approx 2-2,5, triguttatis v. spurie biseptatis.

Hab. ad ramos Lonicerae Xylostei et L. nigræ, in Helvetia, Italia, Germania, Suecia, Gallia, Britannia. — Gregaria, adnata, nec e cortice erumpens, persistens, 2 mm. vix lata, sicca rugoso-crispa, connivens, sed humectata quasi rimose dehiscit. To-mentum longum, ferrugineum, sæpius longitudinaliter adpressum, ad marginem barbatum, densum. A simili *L. flammea* differt sub-stantia receptaculi alba, hymenio crassiore. — * **L. pellita** Pers. Myc. Eur. p. 264: cupulis subglobosis, 3 mm. lat., dense tomentosis e griseo dilute gilvis; sporidiis 12-18 \approx 2,5, 3-4-guttatis. In ramulis Lonicerae Caprifolii in Vogesia et Britannia. — * **L. rhodophæa** Mich. I, p. 66, F. ital. f. 1373: habitu et colore extus *L. barba-tæ*, at disco non albido sed amœne roseo; ascis cylindraceo-clavatis, 90 \approx 5, octosporis; sporidiis cylindraceis curvulis, utrinque

obtusiusculis, 12-13 \approx 3, hyalinis. In lignis putrescentibus Ulmi etc. Conegliano, Selva, Ital. bor. Ad *Lachn. flammeam* quoque vergit, sed extus non rufa.

3. **Lachnella Loniceræ** (A. S.) Fuck. Symb. p. 280, *P. Loniceræ* A. 1617 S. p. 328, t. 11, f. 8, Fr. Syst. Myc. II, p. 115, *Lachn. Periclymeni* Fuck. Symb. p. 280. — Sessilis, ceracea, subglobosa, ex albo sensim fusco-nigra, extus subpilosa, ore connivente crenato; ascis linearibus, 8-sporis, 60 \approx 8; sporidiis subdistichis, cylindraceis sublanceolatisve, continuis, hyalinis, 8-10 \approx 2.

Hab. ad ramos sicclos Loniceræ Periclymeni rarius Corni in Suecia, Germania. — Sparsa, minuta, persistens; sicca 1 mm. lata, triquetra, fusco-nigra, margine albidior; humectata vero explanata, 2 mm. lata cæsio-albida. Subiculum deest, sec. Fries et Fuck. adest vero, etsi evanidum, sec. Alb. et Schw. Cupulæ sec. Fuck. gregariæ, seminis Erucæ magnitudine, strigoso-pilosæ, sordidæ, margine connivente, subclauso, disco sordido.

4. **Lachnella Berberidis** (Pers.) Fuck. Symb. p. 280, *P. Berberidis* 1618 Pers. Syn. p. 649, Fr. Syst. Myc. II, pag. 99. — Sparsa, sessilis, rufo-fuscescens, hirta, basi setis strigosis subradianibus cincta; ascis clavatis, 8-sporis; sporidiis cylindraceis, curvatis, continuis, hyalinis, 8 \approx 2 (Fuck.).

Hab. ad ramos sicclos Berberidis in Germania, Belgio, Italia. — Cl. Rehm hoc dicit dubitanter *L. Periclymeni* Fuck. Substantia sicca. Cupulæ minutæ, basi villis longioribus, subiculum tamen non efficientibus, matrici adnatæ.

5. **Lachnella corticalis** (Pers.) Fr. Summa p. 365, Pat. Tab. fig. 71, 1619 *P. oorticalis* Pers. Obs. I, pag. 28, 112, *P. granulifor.*? Schum. Saell. p. 435, *P. urceolata* Rutstr. diss. pag. 19, *Helot. corticale* Karst. M. F. I, p. 159. — Sessilis, subglobosa, firma, flocculosotomentosa, e cinereo rufescens; pilis simplicibus, septatis, glabris, fuscis, c. 3 μ . cr.; ascis clavatis, apice incrassatis, 75 \approx 6, 8-sporis, jodi cpe apice cærulecente; sporidiis ellipticis, subacutis, rectis, hyalinis, distichis, 12-18 \approx 3-4; paraphy sibus filiformibus.

Hab. supra cortices duriores sicclos arborum adhuc vegetarum Populi, Quercus, Alni etc. in Suecia, Germania, Fennia, Gallia, Italia, Belgio, America bor., Cuba. — Gregaria, persistens, superficialis; humida disco aperto subrufescens et hemisphærica: sicca clausa subdifformis albido-grisea. Species facillime dignoscenda ob amœnitatem colorationis; disco roseo, purpureo v. cin-

nabarino-aurantiaco; excipulo vivide cæruleo vel cinereo-cæruleo; magn. $\frac{1}{2}$ ad $1\frac{1}{2}$ mm. diam.

6. **Lachnella canescens** Cooke Phill. Disc. p. 259. — Gregaria vel 1620 conferta, sessilis, hemisphærica, firma, extus pilis flexuosis, septatis brunneis obsessa; disco fuligineo-brunneo; ascis clavatis; sporidiis fusoideis, denuque simulate 1-septatis, 9-12 \pm 2; paraphysibus filiformibus.

Hab. in lignis nudis pr. Hereford Britanniæ (RENNY). — Ascoma $\frac{1}{2}$ mm. lat. A *L. corticali* differt sporidiis fusoideis, pilis evidentius septatis.

7. **Lachnella tricolor** (Sow.) Sacc. *P. tricolor* Sow. t. 369, f. 6, Fr. 1621 Syst. Myc. II, p. 134, *P. Godroniana* Mont. Cent. III, n. 46, Syll. Crypt. n. 623, *P. episcopalis* L. Duf. in Desmaz. Ann. 2, XIX, p. 367. — Subsessilis vel brevissime stipitata, primum ore clauso breviter turbinata aut clavæformis, tandem aperta, margine inflexo hemisphærica, extus villo amœne violaceo vestita, carne nigrescente; disco ceraceo carneo; stipite obtuso albido; ascis clavatis, 60 \pm 6-7; sporidiis oblongo-cylindraceis, 12-15 \pm 2-3, 4-guttulatis.

Hab. ad corticem in truncis vivis Quercuum prope Nancy, Bruyères Galliæ et in Britannia. — Ascoma 1 mm. lat.

8. **Lachnella papillaris** (Bull.) Phill. Disc. p. 257, *Peziza papillaris* 1622 Bull. Champ. p. 244, t. 467, f. 1, Fr. Syst. Myc. II, p. 102, *Hel. papillare* Karst. M. F. I, p. 160, *P. granuliformis* A. et S. p. 312. — Gregaria, planiuscula, sicca subsphaeroideo-contracta, granuliformis, albida, villosa-hirta, margine granulato, 0,8 mm. lat., pilis marginalibus, 50 \pm 2,5, filiformibus, sursum asperulis, dilute ochraceis; ascis cylindraceo-clavatis, 55-65 \pm 6-7, obturaculo jodo haud vel obsoletissime cærulescente; sporidiis vulgo distichis, fusoideo-oblongatis v. fusoideo-elongatis, 2-guttulatis, interdum spurie tenuiter uniseptatis, 10-15 \pm 2-3; paraphysibus gracilibus.

Hab. ad lignum putridum pineum, populinum in Fennia meridionali, Lapponia, Germania, Britannia, Belgio. — Forma major cupulis 1,5 mm. lat., supra folia putrida *Populi* et *Juniperi* prope Mustiala lecta (KARST.).

9. **Lachnella spadicea** (Pers.) Phill. Disc. p. 258, *P. spadicea* Pers. 1623 Myc. Eur. p. 250. Fr. Syst. Myc. II, p. 98. — Sessilis, subgloboso-hemisphærica, demum expansa, tomento denso subcompacto spadiceo vestita, ore connivente, disco pallido; ascis tereti-clavatis octosporis; sporidiis elliptico-fusoideis, 8 \pm 2; paraphysibus filiformibus.

Hab. ad ramos *Populi tremulæ*, *Tiliæ*, *Ulicis* in Gallia,

Britannia, Germania. — Majuscula, sparsa, substantia sicca. Cupulæ juniores orbiculares, nitide badiæ; adultæ expallentes opaca, $\frac{1}{2}$ mm. lat. Pili breves, septati, intus granulosi.

10. **Lachnella Robergei** (Desm.) Sacc. *Peziza Robergei* Desm. 11 Not. 1624 p. 365. — Erumpens, sessilis, exigua, sparsa, villoso-hirta, junior subglobosa, dein magis hemisphærica, unicolor testacea; ascis subclavatis, 40 μ . longis, octosporis; sporidiis ovoideis.

Hab. in ramis Lonicera rūm prope Caen Galliæ. — Ascoma 1-2 mm. diam., extus pilis brevibus rufulis, margine albescens tectum.

11. **Lachnella cedrina** (Cooke) Sacc. *Peziza cedrina* Cooke et Ger. 1625 in Cooke Disc. Unit. St. II, p. 294, II, p. 22, Hedw. 1875. p. 81. — Sparsa, brunneo-picea, extus fibroso-rugosa; ascomate globoso, cupuliformi, 1 mm. diam.; margine contracto; disco vix pallidiori; ascis cylindraceis; sporidiis ovoideis, 20 \approx 10; paraphysibus clavatis, apice leniter curvis.

Hab. in ramis Juniperi virginianæ, New York Amer. bor. (GERARD).

12. **Lachnella macrochæta** Speg. in Thüm Pilz. d. Weinst. pag. 78, 1626 Mich. I, p. 428, F. ital. f. 1374. — Ascomatibus spadiceo-fuscis cupulatis, sessilibus, $\frac{1}{2}$ - $\frac{3}{4}$ mm. lat., setulis filiformibus, flexuosis, 100-200 \approx 3-4, septatis olivaceo-fuligineis vestitis; ascis cylindraceis, 50-55 \approx 4 $\frac{1}{2}$ - $5\frac{1}{4}$, apice rotundatis, paraphysatis, octosporis; sporidiis fusoideis, utrinque acutiusculis, 7-10 \approx 1 $\frac{1}{2}$ - $2\frac{1}{2}$, 2-guttulatis, hyalinis.

Hab. in sarmenis emotis Vitis viniferæ, Conegliano Italia boreali.

13. **Lachnella? testaceo-rufa** Speg. Fung. Fueg. n. 325. — Cupulis 1627 sparsis v. hinc inde subgregariis, in rimulis fere absconditis, uds explanatis, 1-2 mm. diam., siccis difformi-contractulis, coriacellis, duriusculis, sordide ebsureque testaceis, extus primo parce rufo-puberulis, pilis minutis brevibus continuis flexuosis apice obtusis basi coalescentibus, 20-30 \approx 3 pallide fulvis vestitis, per ætatem glabratis; contextu fulvescenti-ferrugineo parum distincto, subpros-enchymatico, disco non vel vix pallidore plano, margine acutiusculo, subrepandulo; ascis cylindraceo-subclavulatis, apice obtuse rotundatis, deorsum longiuscule attenuatis, 75-80 \approx 5, octosporis; paraphysibus longis, densiusculis, obvallatis; sporidiis recte vel oblique monostichis, elliptico-elongatis vel subnavicularibus, utrinque acutato-obtusatis, 8-10 \approx 3-4, hyalinis.

Hab. vulgata ad rimulas ramulorum emortuorum v. subviventium

Chiliotrichi a melloidis Fuegiæ in Staten island.

14. **Lachnella fraxinicola** (B. et Br.) Phill. Disc. p. 275, *Peziza fra-* ¹⁶²⁸
xinicola B. et Br. Ann. N. H. n. 1160, t. IV, f. 21. — Sparsa v.
 aggregata; ascomatibus externe pallide cervinis, furfuraceis, intus
 brunneis, primo clausis, dein irregulariter apertis, demum orbicu-
 laribus; disco vix depresso; ascis clavatis; sporidiis monostichis,
 ellipsoideis, 10-12 = 7-8.

Hab. in ramulis fraxineis Northamptonshire Britanniæ. —

Ascoma 1 mm. lat. *L. rufo-olivaceæ* affinis et pariter firma.

15. **Lachnella siparia** (B. et Br.) Phill. Disc. p. 276, *Peziza siparia* ¹⁶²⁹
 B. et Br. Ann. N. H. n. 772. — Ascomatibus subsessilibus, extus
 primo subglobosis, dein cyathiformibus furfuraceis ochraceis, disco
 primo ochraceo, dein brunneo; sporidiis linear-i-oblongis, 10-12 =
 2-3, saepe utrinque nucleatis; paraphysibus filiformibus.

Hab. in ramis decorticatis Ulmi Elmhurst in Britannia. —

Subinde adest stratum flocculosum. Ascoma 2-3 mm. lat.

16. **Lachnella Campanula** Ellis Bull. Torr. 1881, p. 73. — Pulvinata, ¹⁶³⁰
 sessilis, circiter 2,5 mm. alta et aperta circiter 2,5 mm. lata, pilis
 tenuibus, niveis, crispis tecta, disco pallide rufo v. carneo, cam-
 panulata, sicca contracta atque densa fimbria pilorum ciliata;
 ascis sessilibus 25 = 5; paraphysibus non visis; sporidiis distichis,
 oblongis, uno apice oblique apiculatis, hyalinis.

Hab. in ramis Fraxini, Utah Amer. bor. (HARKNESS). — Uda
Soleniam in mentem revocans.

17. **Lachnella rubiginosa** Pass. et Beltr. Fung. Sic. n. 18. — Cupulis ¹⁶³¹
 sparsis, solitariis, sessilibus in sicco hemisphærico-contractis, vil-
 loso-tomentosis, rubiginosis, margine pallidiore; ascis clavatis, basi
 attenuatis, breviter stipitatis, 8-sporis, 70-75 = 7-7½; sporidiis ob-
 longo-fusiformibus, distichis, rectis, continuis, opaco-hyalinis, in-
 tens granulosis, 12½ = 3-4; paraphysibus filiformibus strictis, ascos
 subæquantibus v. excedentibus; pilis exterioribus setiformibus lon-
 gissimis integris.

Hab. in ligno denudato Oleæ europææ, Monostalla Siciliæ.

18. **Lachnella cenangoides** Ellis Bull. Torr. Cl. 1881, pag. 123. — ¹⁶³²
 Sessilis, urceolata, dein expansa, 750 mm. diam., extus margineque
 pilis brunneis, rugulosis, septatis tecta; disco sordide albo; ascis cy-
 lindraceis, 65 = 6, octosporis; paraphysibus crassis, sursum septatis,
 apice vix inflatis, obtusis; sporidiis fusiformibus, hyalinis, retis v.
 vix curvis, plerumque uno apice acutis, 15-17 μ . longis.

Hab. in caulibus herbaceis, Utah Amer. bor.

19. **Lachnella Meleagris** Ellis Bull. Torr. Cl. 1881, p. 123. — Ses-¹⁶³³ silis inter fibras ligni, bruneo-purpurea, primo sphærica, dein expansa, 750μ . diam., extus margineque pilis purpureis, obsolete septatis, minute rugosis sursum attenuatis pallidioribus (marginalibus fere hyalinis) tecta; disco sordide albescente; ascis sessilibus, oblongo-cylindraceis, $5-7 \mu$. longis; paraphysibus crassis, 2-3-septatis atque sursum abrupte acuminatis; sporidiis simplicibus, hyalinis, oblongis, $7 \approx 2,5$, in parte superiore asci collectis.

Hab. in ligno senescente, Utah Amer. bor. (HARKNESS).

20. **Lachnella fusco-brunnea** Rehm in Thüm. Pilzfl. Sib. V, p. 17. — Cupulis sparsis, interdum gregariis, sessilibus, globosis subdensis, humiditate vix apertis, strigosis, fusco-brunneis, disco (ut videtur) pallidiore, contextu prosenchymatico, pilis longis septatis, filiformibus fusco-purpureis 3μ . crassis obsessis, circiter 1 mm. diam.; ascis cylindraceo-clavatis, octosporis, $90 \approx 6$, apice jodo cœrulecente; sporidiis subfusiformibus, unicellularibus, hyalinis, distichis, plerumque rectis, subobtusis, $15 \approx 3$; paraphysibus filiformibus, hyalinis.

Hab. in caulibus subputridis Aconiti Lycoctoni, in silvis alpinis pr. montem Kerlygan Sibiriæ. — Ad *Pirottæam* accedit.

21. **Lachnella albido-fusca** Sacc. Misc. myc. I, p. 9. — Ascomatibus gregariis, sessilibus, scutellatis, coriaceo-duriusculis, $1\frac{1}{2}-3\frac{1}{4}$ mm. diam., rufo-atris, breve villosis, setis filiformibus, septatis, fuligineis, levibus, 4μ . cr.; disco concavo, albido, margine leniter involuto; ascis tereti-clavulatis, deorsum attenuato-stipitatis, $50-55 \approx 7-8$, octosporis; sporidiis anguste fusoideis $7-9 \approx 1,5$, distichis, rectis, hyalinis.

Hab. in sarmentis corticatis Rubi idæi, Malmedy (LIBERT).

— *Lach.* barbatæ et *L.* flammeæ subaffinis.

22. **Lachnella escharoides** (B. et Br.) Sacc. *Peziza escharoides* B. et Br. Ann. N. H. n. 1322. t. XIX, f. 15. — Ascomatibus sessilibus, rugosis, rugis e floccis albescentibus, griseo-olivaceis, formatis, margine pallido, floccis delicatis fimbriato; disco cinereo; sporidiis fusiformibus, sœpe binucleatis, 10μ . longis.

Hab. in sarmentis Rubi fruticosi in Britannia. — Ascomata primo clausa, globosa, fere 1 mm. diam., pilis delapsis nigra.

23. **Lachnella lactea** Quél. in Grev. 1879, p. 117 et 12 Suppl. p. 13, t. VII, f. 12. — Ascomate ovoideo-sphærico, 1-2 mm. diam., firmo, pilis granulosis, sericeis, hyalinis vestito; hymenio concavo,

pallide flavescente; sporidiis acicularibus, 15 μ . longis, (spurie?) multiseptatis.

Hab. gregaria in plantis majoribus exsiccatis in collibus Jura.

— *Habitus Cyphellæ villosæ.*

24. **Lachnella velutarioides** Speg. F. Fueg. n. 323. — Hinc inde in 1638
rimulis densiuscule gregaria, sessilis, uda patellari-explanata 1-2
mm. diam. saepe e mutua pressione angulosa, sicca contractula,
marginè subinvoluta, extus pulchre fulvo- v. rufesceni-tabacina,
vilosula, pilis cylindraceis sursum non attenuatis apice tamen acu-
tatis basi non incrassatis sed coalescentibus, 200-250 \approx 4-5 non v.
parce subnoduloso-septatis hyalinis substantia sordide fulvescente
granuloso-incrustatis vestita; disco pallescenti-carneo; ascis cylin-
draceis, apice obtuse rotundatis, deorsum breviter attenuato-pedi-
cellatis, 70-80 \approx 7-8, octosporis; paraphysibus filiformibus vix lon-
gioribus parce obvallatis; sporidiis oblique monostichis, cylindra-
ceis vel allantoideis, utrinque obtuse rotundatis, 11-16 \approx 4, conti-
nuis, hyalinis.

Hab. ad ramulos vivos vel emortuos Chiliostrichi amel-
loidis et Escalloniæ serratæ in Staten island Fuegiæ. —
Contextus excipuli carnosulus fere indistinctus sordidus; jodi vis
nulla.

25. **Lachnella rufo-olivacea** (A. et S.) Sacc. *P. rufo-olivac.* A. et S. 1639
p. 320, t. XI, f. 4, Fr. Syst. Myc. II, p. 99, *Velutaria rufo-oli-
vacea* Fuck. Symb. pag. 300. — Sessilis, applanata, extus villo-
so-pulverulenta sordide rufo-ferruginea, disco virescenti-olivaceo;
ascis tereti-fusoideis, 144-150 \approx 10-14; sporidiis ellipsoideis, mo-
nostichis, 10-14 \approx 6-7, 1-2-nucleatis, hyalinis; paraphysibus apice
clavatis viridi-brunneolis.

Hab. in ramis emortuis humidis Rubi fruticosi, coryli-
folii, cæsii, idæi, rarius in ramis Rosæ, Eucalypti, Ligustræ,
Sambuci, Viburni, Aceris in Suecia, Germania, Italia,
Gallia, Belgio, Britannia, Amer. bor. — Sparsa, orbicularis, adnata,
disco patente plano, primo e viridi-olivaceo, demum nigrescente.
Substantia carnosæ-ceracea, molliuscula, sed sat persistens.

26. **Lachnella diplocarpa** (Curr.) Phill. Disc. p. 232, tab. VII, f. 43, 1640
Peziza diplocarpa Curr. Linn. Trans. XXIV, pag. 153, tab. 25,
f. 30-33. — Ascomatibus planiusculis, obconico-stipitatis, externe
vinoso-brunneis, pilis densis brevibus vestitis, stipite quoque pi-
loso, marginè vix inflexo, fimbriato, disco ceraceo, subglaucio, gri-
seo-olivaceo; sporidiis utrinque nucleatis, rectis, acutiusculis, 10-

12-3; paraphysibus filiformibus apice incrassatis, 2-4-septatis acuminatisque.

Hab. ad terram (?), Joydens Wood, Dartford Britanniæ. — Afinis *L. rufo-olivaceæ*. Ascoma 3 mm. lat. Pili breves, septati, leves, rufi.

27. **Lachnella æruginosa** Sacc. et Sp. Mich. I, p. 443, F. ital. f. 1439 1641 (*Velutaria*). — Ascomatibus sparsis aggregatisque, scutellaribus subsessilibus, $\frac{1}{2}$ - $\frac{2}{3}$ mm. diam., extus breve ochraceo-puberulis, disco concaviusculo concolore, dein fusco-olivaceo; ascis cylindraceis, mediocriter sensim stipitatis, basi ima nodulosis, apice tunica integra rotundatis, 125-130-12; paraphysibus bacillaribus, septulatis, apice clavulâ fusoideâ, 15-22-5-8, crasse chlorino-guttatâ terminatis obvallatis, octosporis; sporidiis monostichis, ellipsoideis, 12-14-7-8, 2-guttulatis nubilosisque, initio hyalinis tandem fulmosis.

Hab. in sarmenis corticatis *Vitis viniferæ*, Belluno Italiæ borealis.

Species imperfecte cognitæ.

28. **Lachnella incarnescens** (Schw.) Sacc. *Peziza incarnescens* Schw. 1642 Syn. p. 173. — Sessilis, gregaria, 2-5 mm. lata, orbicularis, extus nigro-umbrina, pilosula strigosa, margine dilutiori, subclausa, sicca madefacta, expansa, disco eleganter incarnato.

Hab. ad ligna decorticata varia, Bethlehem Amer. bor.

29. **Lachnella cinnabarina** (Schw.) Sacc. *Peziza cinnabarina* Schw. 1643 Syn. p. 173. — Gregaria vix 4 mm. diam. exedens, sessilis, globosa, strigoso-pulverulenta, unicolor, cinnabarina, basi latiuscula nigrescente, rarius disco aperto tota pulvere cinnabarino copiosius tecta.

Hab. ad lignum Liquidambaris, Salem Amer. bor.

30. **Lachnella cinereo-fusca** (Schw.) Sacc. *Pez. cinereo-fusca* Schw. 1644 Car. p. 93, Fr. Syst. II, p. II, f. 97. — Sessilis, applanata, firma, farinaceo-hirta, cinereo-virens, disco fusco-badio.

Hab. in ramis denudatis Cercidis, Gleditschiæ, Vitis vulpinæ Amer. bor. — Gregaria, minuta, 1 mm. lata. Cupulæ, ut in præcedentibus, margine involuto saepius clausæ.

31. **Lachnella atro-fuscata** (Schw.) Sacc. *Peziza atro-fuscata* Schw. 1645 Syn. p. 174. — Sparsa, majuscula (pro ratione) diametro 2 mm. æquante, cupula sessili, extus statu siciori atra, rugosa orbicu-