

paraphysatis, amplis, crasse tunicatis, clavatis, rectis v. gibbis, 65-112 \approx 15-25, 4-8-sporis; sporidiis cymbiformibus, subinde oblongo-ellipticis, granulosis et guttulatis, 20-27 \approx 10-12, utrinque in appendiculam hyalinam 6-7 μ . long. productis.

Hab. in ramis siccis Corni sanguineæ, Vigheffio prope Parma Ital. bor.

TRABUTIA Sacc. et Roum. — Sacc. Syll. I, p. 449.

1. **Trabutia Bauhiniæ** Wint. in Hedw. 1885, I. — Stromate tenui, 2437
sæpe obsoleto, atro, crustaceo; peritheciis gregariis, sæpe stellatim vel radiatim dispositis, non raro concrescentibus, semiimmersis, globosis, vertice subconico protuberantibus, atris, nitidiusculis, parte superiori carbonaceis, atris, parte inferiori (immersis) membranaceis, pallidis, 210-240 μ . diam.; ascis inflato-oblongis, subsessilibus, utrinque attenuatis, membrana tenuissima, apice parum incrassata præditis, 8-sporis, 60-70 \approx 17-22; sporidiis inordinatis, oblongis, utrinque parum attenuatis, sed rotundatis, inæquilateribus, unicellularibus, hyalinis, 16-17,5 \approx 5.

Hab. ad folia viva Bauhiniæ Vahlii in horto botanico Calcuttæ Indiæ (S. KURZ).

2. **Trabutia crotonicola** Rehm in Hedw. 1890, p. 160. — Peritheciis sparsis, epiphyllis, singulis v. 2-3 connato-confertis, primitus in foliorum parenchymate flavo-fuscescente maculato immersis, dein sessilibus, hemisphæricis, carbonaceo-nigris, nitentibus, demum poro minimo pertusis, 0,3-0,5 mm. diam.; ascis clavato-cylindraceis, apice rotundatis, 8-sporis, 90-100 \approx 12-15; paraphysibus filiformibus, teneris, septatis, longissimis, 5 μ . lat.; sporidiis oblongis obtusis, rectis v. subcurvatis, continuis, hyalinis, minute guttulatis, 15-17 \approx 7-8, in ascorum apice sæpe distichis, plerumque monostichis.

Hab. ad folia viva Crotonis floribundi, Rio de Janeiro Brasiliæ (E. ULE).

3. **Trabutia Eucalypti** Cooke et Mass. Grevill. XVII, p. 43. — Epiphylla; stromate coriaceo, suborbiculari, 3 mm. diam., convexo-ruguloso, atro, nitido; peritheciis in stromate innatis, protuberantibus, ostiolo minuto pertusis; ascis cylindrico-clavatis; sporidiis elliptico-lanceolatis, continuis, hyalinis, 30 \approx 8-9.

Hab. in foliis Eucalypti viminalis β . manniferæ in Tasmania Oceaniæ.

DITOPELLA De Not. — Sacc. Syll. I, p. 450.

1. **Ditopella microscopica** Karst. in Rev. mon. p. 71, *Apiosporium microscopicum* Karst. Symb. ad Myc. Fenn. VI, p. 42, Sacc. Syll. I, pag. 132. — Perithecia sparsa, epidermide innata, sphæroidea, subastoma, levia, atra, 30-40 μ . diam.
2. **Ditopella fusispora** De Not. — Syll. I, p. 450. — Forma in Belgio lecta ad Alnum glutinosam (Cfr. Bomm. et Rouss. Contrib. Myc. Belg. II, p. 21) præbet ascis 16-sporos, 60-84 \approx 16-24; sporidia cylindracea vulgo recta, simulate 1-septata, 17-24 \approx 4-5, utrinque rotundata et ciliata.

TRICHOSPHÆRIA Fuck. — Sacc. Syll. I, p. 453.

1. **Trichosphæria Punctillum** Rehm. — Sacc. Syll. I, p. 452. — *T. 2440 **pachyspora** Sacc. Rouss. et Bomm. in Misc. Myc. I, pag. 17. — Peritheciis $\frac{1}{4}$ mm. diam.; setulis septatis, atro-fuscis, 150-200 \approx 8; ascis tereti-fusoideis, 30-80 \approx 10; sporidiis ellipsoideis, utrinque acutiusculis 10-12 \approx 7, granulosis, hyalinis v. dilutissime flavidis.
Hab. in lignis putridis prope Bruxelles in Belgio. — A typo recedit sporidiis paullo brevioribus et crassioribus.
2. **Trichosphæria Rostrupii** Berl. et Vogl. Add. Syll. p. 405, *Trichosphæria* sp. Rostrup. Unders. Svamp. Rhizoct. p. 19, t. I, f. 6-8. — Peritheciis purpureo-brunneis, pseudo-parenchymaticis, orbicularibus, setis rigidis, concoloribus, radiantibus, basi septatis, apice acutis, 100 \approx 5 ornatis; ascis oblongis, sursum acutis, basi in pedicellum breve desinentibus, 45-50 \approx 8-10, octosporis, aparaphysatis; sporidiis inordinate distichis, allantoideis, utrinque uniguttatis, rectis vel curvulis, 14-18 \approx 5.
Hab. in radicibus Ligustri in Islandia.
3. **Trichosphæria pachnostoma** (B. et C.) Cooke Grev. XV, pag. 80, 2442 *Sphæria pachnostoma* B. et C. — Peritheciis lanosis, 1 mm. diam.; ascis clavatis; sporidiis lanceolatis, continuis, hyalinis, 30 \approx 5.
Hab. in ligno in Venezuela.
4. **Trichosphæria pilosa** (Pers.) Fuck. ***nitidula** Sacc. Mich. II, p. 605. 2443 — A typo differt setulis bacillaribus tantum verticeque peritheciis nudato nitidulo; setæ 200-260 \approx 5-6, cuspidatæ fuscæ; ascis tereti-clavatis 54-58 \approx 3-4; sporidiis submonostichis, oblongatis 7-8 \approx 2,5-3, 2-3-guttulatis, hyalinis.
Hab. in sarmenis corticatis Rubi fruticosi prope Rouen Galliæ (LETENDRE).

5. **Trichosphæria corynephora** (Cooke) Sacc., *Byssosphaeria corynephora* Cooke Grev. XI, p. 109. — Effusa, aterrima; hyphis repentibus, septatis, ramulosis, ramulis assurgentibus; conidiis clavatis, infra longe attenuatis, fuscis, multiseptatis (9-11) superne truncatis 150 \approx 18; peritheciis subglobosis, depressis, superne glabris, ad basim tomentoso-intertextis; ascis cylindraceo-clavatis; sporidiis lanceolatis, continuis, hyalinis, 20 \approx 4.

Hab. in ramis Ostryæ virginæ in Amer. bor.

6. **Trichosphæria Elisæ-Mariæ** Sacc. et Pat. Rev. mycol. V, p. 25, 2445 t. XXXVI, f. 5. — Peritheciis hinc inde gregariis superficialibus, globosis, exiguis, $\frac{1}{6}$ mm. diam., atris subcarbonaceis, setulis filiformi-cuspidatis, septulatis fuligineis rectiusculis vestitis; ascis tereti-clavatis, stipitatis, paraphysatis, 90-100 \approx 7-8, octosporis; sporidiis oblique monostichis, breve fusoideis, rectis v. inæquilateribus, 15-18 \approx 5-5,5, continuis, guttulatis 5-7 saepius uniseriatis, hyalinis. — Adsunt hinc inde cæspituli conidiophori ab *Acrothecio simplici* (conidiis fusoideis, 3-septatis fuligineis subcapitatis) formati.

Hab. in foliis siccis Epilobii spicati pr. Bruxelles Belgii.

Legerunt cl. Dominæ ELISA BOMMER et MARIA ROUSSEAU.

7. **Trichosphæria Hariotiana** Karst. et Har. Rev. Myc. 1890, Oct. — 2446 Peritheciis sparsis v. confertim gregariis, erumpentibus, superficialibus, submembranaceis, lentiformibus v. subsphæroideis, atris, pilis simplicibus, concoloribus, rigidis, continuis, strictis vel leviter flexuosis, divergentibus, 50-90 μ . long., vestitis, basi vulgo macula fuliginea cinctis, circ. 0,2 mm. lat.; ascis fusoideo-elongatis, subsessilibus, curvulis, circ. 60 \approx 10; sporidiis octonis, ellipsoideo-oblongatis, utrinque leviter attenuatis, continuis, intus granulosis, rectis, hyalinis v. chlorino-hyalinis, 14-17 \approx 6-7; paraphysibus nullis.

Hab. ad Orchidaceam quandam cum *Glæosporio pallido* insul. Madagascar in Calidariis musei parisiensis.

8. **Trichosphæria? barbicincta** (E. et E.) Sacc., *Byssosphaeria* E. et 2447 E. Journ. Myc. 1888, p. 63. — Peritheciis sparsis, depresso-globosis, circ. $\frac{1}{4}$ mm. diam., ostiolo papilliformi, distincto, nigro, micante præditis; ascis anguste clavato-cylindraceis, circ. 75 \approx 6, copiose paraphysatis; sporidiis distichis, oblongis, 3-4-nucleatis, hyalinis, 9-12 \approx 2,5-3, utrinque subobtusis.

Hab. in Diatrype Tremellophora vetusta et cortice adiacente Magnoliæ, Newfield, New Jersey Amer. bor. — Matrix pilis erectis, opacis, fragilibus, 250 μ . long. tecta.

9. **Trichosphæria lichenum** Karst. et Har. Rev. Myc. 1890 Oct. — 2448

Peritheciis superficialibus, sphæroideis vel sphæroideo-conoideis, confertim gregariis, carbonaceis, atris, setulis strictis divergentibus, continuis, acutis, concoloribus, 60-90 μ . longis, sat parce vestitis, 0,1 mm. v. paullo ultra diam.; ascis fusoideis, curvulis, circ. 45 \pm 6-7; sporidiis octonis, di- vel oblique monostichis, fusoideo-elongatis, hyalinis, 6-9 \pm 2.

Hab. in thallo Peltigeræ caninæ, Fontainebleau Galliæ (TULASNE).

TRICHOSPHÆRELLA Bomm. Rouss. Sacc. (Etym. a *Trichosphæria*, cui affine genus). — Perithecia superficialia, globoso-ovoidea, membranaceo-coriacea, atra, setulis vestita. Asci tereti-clavati, 16-spori, paraphysati. Sporidia ellipsoidea v. globulosa, continua, hyalina. — Est *Trichosphæria* pleiospora.

1. **Trichosphærella decipiens** B. R. S. sp. n. — Peritheciis ovoideis, 90- 2449
120 μ . diam., superficialibus, sparsis, opace nigro-brunneis, vertice setuli rigidis, acuminatis, continuis, parcis (7-10) coronatis; ascis sessilibus, tereti-clavatis vel anguste fusoideis, 16-sporis, 30-36 \pm 6-7; paraphysibus parum distinctis; sporidiis distichis, ellipsoideis v. subglobosis, 4-5 \pm 2,5, continua, hyalinis, vulgo 2-guttulatis.

Hab. in ramis decorticatis Fagi silvaticæ, Poix Galliæ.

SCORTECHINIA Sacc. (Etym. a sollertiss. botanico italo Rev. BENED. SCORTECHINI, Australiæ et Malesiæ exploratore), *Trichosphæria* Sacc. ex p. Syll. I, p. 454, *Sphæria* Mont. — Perithecia in subiculo late effuso, compacto, pannoso fusco, fere penitus immersa, minutissima, levia, epapillata, tandem pertusa, vix carbonacea. Subiculis hyphæ subdichotomæ ramosæ, filiformes, fuligineæ, continuæ, ubique spinulis, apice acute 2-3-dentatis exasperatæ. Asci clavati, tenerrimi, longe stipitati, apice obtusi, octospori, a paraphysati. Sporidia disticha, ex oblongo breve fusoidea, utrinque acuta, hyalina.

1. **Scortechinia Culcitella** (B. et Rav.) Speg. F. Guar. II, pag. 27, 2450
Sphæria Culcitella B. et R. — Syll. II, p. 402 — *Scortechinia acanthostroma* Sacc. et Berl. Misc. Myc. II, n. 22, nec *Sphæria acanthostroma* Mont. seu *Trichosphæria acanthostroma* Sacc. Syll. I, p. 454. — Asci p. sp. 35-40 \pm 9, stipite 35-40 \pm 2,5; sporidia 10-11 \pm 3,5-4,5, biguttata: adsunt quoque conidia hyphis inspersa obovato-oblonga, 12-13 \pm 5-6, binucleata, pallide lutescentia.

Hab. ad cortices arborum, Burleigh Head Australiæ et in Brasilia, Venezuela et Cuba. — Sec. Speg. l. c. *Trich. acanthostroma* (Mont.) Sacc. Syll. I, p. 454 est species omnino diversa sub *Trichosphaeriæ* titulo servanda et cum *Sphæria foveolata* B. et C., Sacc. Syll. I, p. 271 (sub *Rosellinia*) identica. Viget hæc quoque in Brasilia.

WALLROTHIELLA Sacc. Syll. I, p. 455.

1. **Wallrothiella eunotiaspora** (Cooke et Hark.) Berl. et Vogl. Add. 2451 Syll. p. 68, *Sphæria eunotiaspora* Cooke et Harkness in Grevillea XIII, p. 18. — Superficialis, gregaria; peritheciis obpiriformibus, $\frac{1}{2}$ mm. diam., atris, subnitidis, levibus, fragilibus; ascis clavatis; sporidiis ellipticis, medio inflatis, biseriatis, continuis, 30-35 \times 12-14, plasmate granuloso.

Hab. in *Acacia* quadam decorticata ex Australia, in California (HARKNESS). — Socia adest *Diplodia* quædam.

2. **Wallrothiella Fendleri** B. et C. in Herb. Berk. 9628, Grevill. XV, 2452 p. 82. — Peritheciis densissime gregariis, globosis, atris, demum depresso-arcuatis, utrinque acutis, multinucleatis, hyalinis, 50 μ . long.

Hab. in ligno, ubi?, an in America centrali?

3. **Wallrothiella Rhododendri** (Ces.) Sacc. *Sphæria* Klotz. Herb. Myc. 2453 n. 1836, Grev. XIX, pag. 72. — Peritheciis perfectis, subglobosis, ostiolum minutum quidem sed distinctum papillæforme præbentibus; ascis cito evanescentibus; sporidiis ovoido-oblongis, hyalinis, simplicibus.

Hab. in ramis Rhododendri in Ital. bor.

4. **Wallrothiella Salicis** Har. et Br. Rev. Myc. 1891, p. 15. — Peritheciis superficialibus, sparsis vel approximatis, irregulariter globosis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., brunneis, depresso-rugosis, supra sæpe crateriformibus; ascis cylindraceis, sessilibus, 8-sporis, 60-80 \times 10-12; paraphysibus cohærentibus; sporidiis distichis vel oblique monostichis, ovoides, hyalinis, 14-16 \times 8.

Hab. in ramulis decorticatis Salicis, Loudun, Vienne Galliæ.

BOTRYOSPHÆRIA Ces et De Not. — Sacc. Syll. I, p. 456.

1. **Botryosphaeria Quercuum** (Schw.) Sacc. Syll. I, p. 456, var. **Carpinii** Sacc. Misc. Myc. II, pag. 21, n. 92. — Acervulis pulvinatis, minutis, aterrimis; ascis clavatis, crasse tunicatis, 120-140 \times 28-32,

- 606 Pyrenomyceteæ, Sphæriaceæ, Hyalosporæ, Botryosphæria.
crassiuscule paraphysatis, octosporis; sporidiis distichis, ovoideis,
28-32 \approx 14-18, intus granulosis, hyalinis.
Hab. in ramis Carpini, Troyes Galliæ (BRIARD).
2. **Botryosphæria minor** Ell. et Ev. Journ. Myc. 1888, p. 65. — Pe- 2456
ritheciis minutis, 150-170 μ . diam., intus albis, plerumque 2-6 in
stromate imperfecto gregariis, erumpentibus; ascis 75 \approx 20; spori-
diis confertim distichis, subellipsoideis, utrinque angustatis, 14-16
 \approx 6-7, flavidо-hyalinis.
Hab. ad Sesbaniam, Louisiana Amer. bor. (LANGLOIS). — A
B. fuliginosa minutie præcipue differt.
3. **Botryosphæria Ficus** (Cooke) Sacc. *Melogramma Ficus* Cooke Gre- 2457
villea XI, p. 108. — Erumpens; peritheciis minimis, obtusis, atris,
opacis, in cæspitulos lineasque erumpentibus; ascis clavatis; spo-
ridiis ellipticis, lanceolatis, continuis, hyalinis, 25 \approx 10.
Hab. in cortice Fici Caricæ, Aiken Carolina inf. Amer. bor.
(RAVENEL).
4. **Botryosphæria Sumachi** (Schw.) Cooke Grev. *Sphæria Sumachi*
Schw. — Syll. II, n. 4220. — Sporidia majuscula ellipsoidea,
hyalina.
5. **Botryosphæria inflata** Cooke et Mass. Grevill. XVII, pag. 42. — 2458
Peritheciis cortice interiore nidulantibus, demum rimoso-erumpen-
tibus, papillatis, glabris, atris, contextu coriaceo, rimis arcte con-
niventibus, graphideis, flexuosis; ascis clavatis, 8-sporis; sporidiis
distichis, ellipticis, utrinque obtusis, medio inflatis, continuis, hya-
linis, 33-35 \approx 10.
Hab. in cortice quodam, Habgalla Ceylon Asiæ austr.
6. **Botryosphæria hypoxylloidea** Cooke in Grevillea XII, p. 102. — 2459
Mox superficialis, hypoxylloidea; peritheciis minimis, in pustulas
orbiculares coacervatis, convexis, rugosis, atris, opacis, confluen-
tibus; ascis cylindrico-clavatis, octosporis; sporidiis ellipticis, con-
tinuis, hyalinis, 12 \approx 5.
Hab. in ramis in Australia.
7. **Botryosphæria Callicarpæ** Cooke in Grevillea XIII, pag. 101. — 2460
Longitudinaliter erumpens; stromate elongato; peritheciis parvu-
lis confluentibus, atris; ascis clavatis; sporidiis continuis (imma-
turis visis).
Hab. in cortice Callicarpæ americanæ, Darien Georgiæ.
8. **Botryosphæria Berengeriana** De Not. — Sacc. Syll. I, p. 457. * **B. pachyspora** Sacc. Misc. Myc. II, pag. 22, n. 31. — A typo recedit

sporidiis crassioribus, nempe 20-32 \approx 14-18, et ascis angustioribus, nempe 120-180 \approx 28-32.

Hab. in ramis emortuis *Carpini*, Troyes Galliæ (BRIARD).

9. **Botryosphæria melachroa** Berk. et Curt. in Herb. Berk. Cooke 2462 in Grevillea XIII, p. 101. — Stromate longitudinaliter effuso, innato, erumpente; peritheciis ovatis, confertis, subdiscretis, atris, ad apicem rotundatis, paucis; ascis clavato-cylindricis; sporidiis ellipticis, continuis, 20 \approx 8, hyalinis.

Hab. in *Cratægo cordato* in Pennsylvania Amer. bor.

10. **Botryosphæria mutila** (Schw.) Cooke in Grevillea, *Sphæria mu-* 2463 *tila* Schw. Amer. bor. n. 1439, sec. spec. — Stromate innato; peritheciis in series flexuosa erumpentibus subconfluentibus, globosis, superiore parte prominente, inæquali, rugosa, intus nucleo albido farctis; ascis clavatis, octosporis; sporidiis ellipticis, continuis, hyalinis, 20-22 \approx 9,

Hab. in *Populo* etc. in America boreali. — Verisimiliter hoc spectat *Botryosphæria Berengeriana*.

11. **Botryosphæria Syringæ** (Schw.) Cooke in Grevillea *Sphæria Sy-* 2464 *ringæ* Schw. Amer. bor. n. 1667. — Ascis elongato-clavatis, octosporis; sporidiis arcte ellipticis, continuis, hyalinis, 26-28 \approx 8.

Hab. in ramis *Syringæ* in Amer. bor. (ELLIS).

12. **Botryosphæria Araliæ** Curt. Cat. 143, Cooke in Grevillea XIII, 2465 pag. 101. — Stromatibus innato-erumpentibus, corticolis, subpulvinatis, nigris, plerumque lineato-aggregatis, 1 mm. diam., albo-farctis, peridermate cinctis; peritheciis vix distinctis, obtusiusculis, opacis; ascis clavatis, octosporis; sporidiis ellipticis, continuis, hyalinis, 18 \approx 8.

Hab. in cortice *Araliæ spinosæ* in Carolina Amer. bor.

13. **Botryosphæria Viburni** Cooke in Grevillea XIII, p. 102. — Stromatibus erumpentibus, gregariis, minimis $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. diam., corticolis, depresso pulvinatis, atro-fuscis, superficie tuberculosis, nucleo albo farctis; peritheciis 6-10, subimmersis, vertice convexis, pertusis; ascis clavatis; sporidiis arcte ellipticis, continuis, flavescentibus, intus granulosis, 18-21 \approx 8.

Hab. in ramis *Viburni Opuli* in America boreali.

14. **Botryosphæria subconnata** (Schw.) Cooke in Grev. XIII, p. 101, *Sphæria subconnata* Schw. Am. bor. n. 1443, Sacc. Syll. II, p. 393, *Winteria valsarioides* Rehm in Thüm. Myc. Univ. n. 2166. — Specimina Berkeleyana et Schweinitziana perfecte congruunt.

15. **Botryosphæria abrupta** Berk. et Curt. in Herb. Berk. Cooke in 2467

Grevillea XIII, p. 101. — Stromatibus erumpentibus, epidermide rimosa arcte cinctis, subdiscoideis; peritheciis globosis, semiimmersis, atris, demum confluentibus, truncato-obtusis, abruptis, intus albis; ascis clavatis, octosporis; sporidiis elliptico-fusiformibus, utrinque subattenuatis, obtusis, continuis, hyalinis, flavescentibus, intus granulosis, 22-24 = 10.

Hab. in *Cyrilla* in *Carolina Americæ*.

16. **Botryosphaeria imperspicua** Passer. Diagn. Fung. nuovi I, n. 7. 2468

— Peritheciis paucis, minutis, papillatis, in acervulos parvulos pulsuliformes vel lineares vix prominulos, saepe tectos, aggregatis, fuscis; ascis paraphysatis, oblongo-clavatis, attenuato-stipitatis, 8-sporis, 175 = 15; sporidiis continuis, rhombeis, intus granulosis et guttulatis, hyalinis, 20-25 = 10.

Hab. in caulis siccis *Euphorbiæ Characiæ* pr. Lacum Bolsena in prov. Roma.

17. **Botryosphaeria Hypericorum** Cooke in Grevillea XIII, p. 102. — 2469

Subelliptica, planiuscula, atra; peritheciis parvulis, connatis, albo-farctis, epidermide fissa arcte cinctis; ascis clavatis, octosporis; sporidiis anguste ellipticis, continuis, hyalinis, circiter 20 = 5, immaturis.

Hab. in caule *Hyperici* proliferi in America boreali. — Ad **Botryosphaeriam** forte pertinent *Hypoxylon Pseudotubulinam* Ces. (1379), *Hypoxylon palmigenum* B. et C. (1435); sporidiis enim hyalinis et habitu dothideaceo pollent.

GIBELLIA Sacc. in Misc. Myc. II, p. 12 (Etym. a præclaro botanico prof. JOSEPHO GIBELLI de metagenesi Pyrenomycetum deque pathologia mycogena optime merito). — Stromata subcutaneo-erumpentia, e basi definita depresso-pulvinata, atra, intus pallidiora, superficie ostiolis vix prominulis punctulata. Perithecia globulosa, monosticha. Asci tereti-fusoidei, sessiles, spuriæ paraphysati, lumine apice bifoveolati, octospori. Sporidia globoso-ellipsoidea, continua, hyalina. — *Dothideaceas*, præsertim *Mazzantiam*, fere imitatur, sed perithecia vera adsunt. Inter *Sphæriaceas hyalosporas*, inter *Botryosphaeriam* et *Cryptosporellam* locandum.

1. **Gibellia dothideoides** Sacc. et Berl. Misc. Myc. II, n. 23. — Stro- 2470 matibus e basi elliptica depresso-pulvinatis, laxe gregariis, 2 = 1-1,5 mm., epidermide secedente demum nudatis, atris, intus pallidioribus, ostiolis leviter prominulis, remotiusculis, punctiformibus; peritheciis globulosis, monostichis, $\frac{1}{3}$ - $\frac{1}{3}$ mm. diam., in quoque stro-

mate 6-10; ascis tereti-fusoideis, sessilibus, apice obtusis lumine-que bifoveolatis, 90-100 \approx 20-25; paraphysibus spuriis, guttulatis obvallatis, octosporis; sporidiis distichis vel oblique monostichis e globoso-ellipsoideis, 18-20 \approx 12-14, bi-pluriguttulatis, hyalinis.

Hab. in ramis corticatis lignosis ignotæ speciei in M. Moon Australiæ (SCORTECHINI). — A *Bagnisiella Tamaricis* et *B. Ilicis*, quæ *Botryosphaeriam* propius spectant, omnino diversa.

CRYPTOSPORELLA Sacc. Syll. I. p. 466.

1. **Cryptosporella divergens** (Schw.) Berl. et Vogl. Add. Syll. p. 70, 2471
Sphæria Schw. Syn. Car. 123 et Syn. Amer. bor. n. 1393, *Valsa divergens* Cooke in Grevillea XIII, p. 40. — Stromate tumidiu-sculo; acervulis peritheciorum 5-10 mm. diam. æquantibus, sub-orbicularibus, epidermide cinctis; peritheiis atris vel cinerescen-tibus, ostiolis teretibus, longissimis, perithecio triplo longioribus, divergentibus; ascis octosporis; sporidiis subellipticis, hyalinis, 7 \approx 2, endochromate granuloso farctis.
Hab. in ramis dejectis Liquidambaris in Amer. bor.
2. **Cryptosporella paucispora** (Peck) Berl. et Vogl. Add. Syll. p. 70, 2472
Valsa Peck in 33 Rep. St. Mus. 1883, p. 33. — Pustulis epider-mide vix elevata, demum rupta tectis; peritheiis 2-5, cortice in-teriori insidentibus; ostiolis brevibus, nigris, discum minutum, pal-lidum perforantibus, raro leniter radiato-sulcatis; ascis brevibus, 62-70 μ . longis, subcylindraceis, tetrasporis; sporidiis simplicibus, monostichis, subhyalinis, ovato-ellipsoideis, 15-20 \approx 10-11.
Hab. in ramulis emortuis Alni, North Greenbush Amer. bor.
3. **Cryptosporella Lentiginis** Rehm in Ellis Amer. Nat. 1883, Febr. 2473 p. 195. — Peritheiis globosis $\frac{1}{2}$ mill. diam., membranaceis, ple-rumque 3-4 aggregatis, stromate corticali basique lignum penetrante; ostiolis brevibus, epidermidem pustulatim elevatam perforanti-bus; ascis clavato-cylindraceis, 45 \approx 7-8; sporidiis biseriatis, cylin-draceis, hyalinis, rectis curvulisve, 2-3-nucleatis, 12 \approx 2-2.5.
Hab. in ramis Viburni Lentiginis, Decorah Iowa Amer. bor. (E. W. HOLWAY). — Substantia corticis ex stromate nigrificata.

4. **Cryptosporella farinosa** (Ellis) Sacc. *Valsa farinosa* Ellis Torr. 2474 Bot. Cl. 1882, p. 99. — Stromate corticali; peritheiis paucis 2-4 pallidis, 350 μ . diam. corticem pustulatim elevantibus; disco tu-berculiformi, flavidо-albo, furfuraceo; ostiolis latiusculis, olivaceo-nigris, ovoideis v. conicis, dein dilabentibus, foveolaque relinquentibus.

tibus; ascis clavato-cylindraceis, obtusis; paraphysibus indistinctis; sporidiis 1-2-stichis, acute ellipticis, subhyalinis, 1-nucleatis, 15-35.

Hab. in corticibus, New Jersey Amer. bor. — Species ostiolis mire ludibunda.

5. **Cryptospora Ilicis** H. Fab. *Spher. Vaucl.* II, p. 34, f. 3. — Peritheciis modo sparsis, solitariis, cortici simul ac ligno infossis, peridermio nigrificato et pustulatum inflato tectis, modo 2-10 in pulvinulos valseos 1-1 mm. $\frac{1}{2}$ latos, epidermide rupta cinctos, absque stromate irregulariter aggregatis, globosis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. latis; ostiolis atris nitidis, papillulatis, sparse e vertice pulvinuli emergentibus; ascis aparaphysatis, cylindraceis, brevissime stipitatis, fere sessilibus, 8-sporis, 150-160 = 8-9; sporidiis oblique monostichis, hyalinis, fusoides, utrinque arcte attenuatis et longe acuminatis 1-magni-guttatis et granulosis, 30-34 = 6-8.

Hab. ad ramos et ramulos deciduos *Quercus Ilicis*, in umbrosis Vaucluse Galliæ.

6. **Cryptospora Ampelopsis** (Ellis) Sacc. *Valsa Ampelopsis* Ellis 2476 Bull. Torr. Bot. Club. 1852, pag. 112. — Peritheciis paucis ligno insidentibus stromateque distincto parentibus; ostiolis cylindraceis subacutis, apicibus in discum ellipticum planum collectis, corticem dein secedentem rimose perforantibus; ascis clavato-cylindraceis; sporidiis biseriatis ellipticis subacutis, 1-nucleatis, v. 2-3-nucleolatis, 17 = 6-7, hyalinis.

Hab. in ramis Ampelopsis quinquefoliæ, New Jersey Amer. bor.

DIAPORTHOPSIS H. Fab. *Sphér. Vaucl.* II, p. 35, f. 3 (Etym. *Diaporthe* et *ops*, aspectus). — A genere *Diaporthe* (cujus sporidia didyma sunt) differt sporidiis absolute continuis; cetera eadem. — Huc spectant: *Diaporthopsis nigrella* (Niessl) H. Fab. l. c. (= *Diaporthe nigrella* Syll. I, pag. 648).

INZENGÆA Borzì Ein neu. Ascom. (Etym. a claro mycologo siculo G. INZENGA). — Perithecia superficialia, globoso-obovata, poro ampio pertusa, majuscula. Ascii primum globosi, dein stellati, octospori. Sporidia stellata, uninucleata rubentia, hyalina. Status conidicus *Coremium Borzianum* Sacc. — Ob sporidia et ascos eximie stellata potissimumque ob fabricam peritheiorum genus hoc familiolam propriam constituere forte meretur.

1. **Inzengæa erythrospora** Borzi Ioz. ein. neu. Ascomyc. tab. XIX, 2477 fig. 9-10, *I. asterospora* Borzi F. parass. oliv. p. 1. — Perithe-

ciis superficialibus, verticaliter globoso-obovatis, subiculo myceliali insidentibus, poro ampio pertusis, $\frac{1}{3}$ - $\frac{3}{4}$ mm. altis; ascis globosis, primum tunica levi instructis demum crasse muricatis stellatisque, 17 μ . diam., octosporis; sporidiis stellatis, crasse uninucleatis, 8 μ . diam., hyalinis.

Hab. in fructibus putridis Oleæ europææ, S. Filippo Sicilæ. — Status conidicus *Coremium Borzianum* Sacc. — Sec. Cooke (Grev. XVI, p. 116), *Thecospora bifida* Hark. — Syll. IV, p. 679 — est hujus speciei status imperfectus. *Emericella variecolor* B. et Br. — Syll. VII, 1, pag. 154 — est secundum analysim cl. Pat. in Bull. Soc. Myc. 1891, pag. 45, tab. IV, fig. 6-12, fungus ex eodem genere et vix differt perithecio villoso, flavido 1-3 mm. alt. lignicolo; ascis compresso-stellatis, 12-15 μ . diam. 6-8 μ . cr.; sporidiis 8-10-dentatis, 3-4 μ . diam. (sine dentibus), rubris. — Nomen *Emericella* multo anterius sed ejusdem diagnosis a Berkeleyo data omnino falsa.

HORMOSPHÆRIA Lév. in Ann. Sc. Nat. 1863, p. 297 (Etym. *hormos* monile et *Sphæria*). — Perithecia carnosa, grumosa, unilocularia, sporangiis 3-4 repleta. Sporangia membranacea, tenuia, subglobosa, sporidiis concatenatis farcta.

1. **Hormosphæria tessellata** Lév. in Ann. Sc. Nat. 1863, pag. 297. — 2478
Peritheciis gregariis, discretis, erumpentibus, globosis, tessellato-rimosis, maculis albidis, orbicularibus vel irregularibus, ambitu limitato, primo albis, nitoris argentei, demum nigris, opacis, paginam superiorem foliorum occupantibus insidentibus, ostiolis (quando conspicuis) papillatis; sporangiis membrana tenuissima anhista, plus minusve globosis, in perithecii cavitate liberis, pallide viridis; sporidiis globosis, concatenatis, levibus, hyalinis.

Hab. ad folia Thibaudiae floribundæ, Bogota, Paramo San Fortunio Novæ Granat. (LINDIG). — « Genus valde paradoxum; an inter *Algas Nostochineas* locandum? »

Sect. 4. Hyalodidymæ.

SPHÆRELLA Ces. et De Not. — Sacc. Syll. I, p. 476.

I. Herbicolæ.

1. **Sphærella fusispora** Fuck. in Reis. nach Nordpolarmeer et Oud. 2479
Contr. Fl. Myc. Now. Semlja pag. 151, tab. II, fig. 4. — Perithe-

ciis in gregibus laxis sub epidermide nidulantibus, demum totis liberis, minutis, aterrimitis, globosis, ostiolo papillæformi, truncato, perforato; ascis fasciculatis, oblongis, sessilibus, 8-sporis, $60 \approx 20$; sporidiis oblique imbricato-distichis, fusiformibus, subrectis, utrinque obtusiusculis, inæqualiter didymis, ad septum parum constrictis, loculis 1-2-guttulatis, $24 \approx 6$, hyalinis.

Hab. in foliis et caulinis exsiccatis Ranunculi pygmaei, Nowaja Semlja seu Nova Zambla.

2. **Sphærella nivalis** Oudem. Contr. à la Flore myc. de Nowaja Semlja p. 11, t. I, f. 8. — Peritheciis minutis, primitus infra epidermidem absconditis, postremo vertice prominulis, poro pertusis, quoad structuram tenerrimis; ascis cylindraceis, brevissime pedicellatis, $90 \approx 9$, 8-sporis; sporidiis monostichis, hyalinis, oblongis, 1-septatis, loculo superiore subampliore, $11-12 \approx 4-5$.

Hab. in Ranunculo nivali var. sulphureo, Nowaja Semlja.

3. **Sphærella septorioides** Peck 32 Rep. p. 52. — Maculis paucis, orbicularibus, angulosis v. irregularibus, albis; peritheciis numerosis, minutis, hypophyllis, nigris; ascis oblongis; sporidiis confertis, 1-septatis, hyalinis, $12,5 \mu$., loculis subæqualibus.

Hab. in foliis viventibus Thalictri dioici, Central Bridge Amer. bor.

4. **Sphærella Thalictri** Ell. et Ev. in Journ. of Myc. 1885, p. 44. — 2482 Peritheciis hypophyllis, 90μ . diam., contextu crasse parenchymatico, sursum late pertusis, hemisphæricis, in siccitate complanatis, sparsis in maculis parvis, 2-3 mm. diam., rotundis, margine obscurō cinctis; ascis sessilibus, $36 \approx 7$; sporidiis stipatis, ovato oblongis, granulosis, nucleatis (demum uniseptatis?), $12-16 \approx 3-4\frac{1}{2}$.

Hab. in foliis Thalictri dioici, Parsippany Americæ borealis (W. TRELEASE).

5. **Sphærella Allescherii** Sacc. *Sphærella Thalictri* Allesch. Süd-Bay. 2483 II, Nachtr. p. 60. — Peritheciis maculas nigras non efformantibus, gregariis vel sparsis, initio epidermide tectis, dein superficiibus, globosis, apice pertusis; ascis cæspitosis, subcylindraceis, circ. 50-60 ≈ 8 , aparaphysatis; sporidiis oblique distichis, clavatis, sæpe fusoideis, 2-locularibus, uno loculo sæpius latiore, utrinque obtusiusculis, circ. $12-15 \approx 2-2,5$, hyalinis.

Hab. in caulinis siccis et putridis Thalictri aquilegifo-lii pr. München Bavariæ. — *S. caulincola* affinis.

6. **Sphærella Vitalbæ** Passer. Diagn. F. nuovi I, n. 8. — Peritheciis 2484

crebre sparsis, punctiformibus, atris; ascis brevibus, basi ventricosis, breviter constricto-stipitatis, 62 ± 20 , 8-sporis; sporidiis subtristichis, apice altero obtusiore, $20-25 \pm 7-7,5$, hyalinis.

Hab. in sarmenis aridis Clematidis Vitalbæ, Fornovo Parma.

7. **Sphærella vitalbina** Passer. Diagn. F. nuovi III, n. 6. — Peritheciis sparsis, minutis, tectis, globoso-depressis, atris; ascis variis, elongato-clavatis vel inferne inflatis aut ovatis et gibbis, $30-75 \pm 10-15$, 8-sporis; sporidiis distichis vel congregatis, navicularibus vel cuneatis, medio septatis, leniter constrictis, $12,5 \pm 2,5-3$.

Hab. in sarmenis aridis Clematidis Vitalbæ, Sala prov. Parma. — A *S. Vitalbæ* differt præcipue sporidiis multo minoribus et etiam ascorum forma varia.

8. **Sphærella applanata** Ell. et Ev. Journ. Myc. 1888, p. 98. — Sparisa, punctiformis, applanata, epidermide pellucida tectis; ascis oblongis, $40-50 \pm 15$; sporidiis biseriatis oblongo-ellipsoideis vel piriformibus, 1-septatis, hyalinis, $18-20 \pm 6-8$.

Hab. in caulis emortuis Clematidis ligusticifoliæ, Sand Coulee Mont. Amer. bor. (ANDERSON).

9. **Sphærella Papaveris** Fuck. in Heuglin Reisen Nordpol. vol. III 2487 et Oudem. Contr. Fl. Myc. Nov. Seml. p. 6, tab. II, f. 2. — Peritheciis in foliorum nigrescentium pagina superiore et inferiore gregariis, et fere foliorum totam superficiem occupantibus, minutissimis, aterrimis, conicis; ascis fasciculatis, oblongis, 8-sporis, 56 ± 24 ; sporidiis congregatis, oblongo-clavatis, utrinque obtusis, rectis, inæqualiter didymis, $22 \pm 8-10$, hyalinis.

Hab. in foliis exsiccatis Papaveris nudicaulis in insula Nowaja Semlja seu Nova Zembla.

10. **Sphærella pachyasca** Rost. Fungi Groenl. p. 552. — Peritheciis sparsis vel gregariis, foliicolis vel cladogenis; ascis crasse ovoideo-oblongatis, inæquilateralibus, $40-50 \pm 15-24$, vulgo apice tunicatis; sporidiis congregatis, conoideo- vel ovoideo-oblongatis, $16-20 \pm 5-6$.

Hab. in foliis caulisque herbarum plurimarum, e. g. Epilobii, Drabæ, Arabidis, Ranunculi, Thalictri, Saxifragæ, Pyrolæ, Pleurogynæ, Campanulæ, Thymi, Stenhammariae, Plantaginis, Diapensiæ in Groenlandia. — Respondet *Sph. Tassianæ*.

11. **Sphærella Viciæ** Schroet. Nord. Pilze p. 10. — Peritheciis 300 μ . 2489 lat., immersis, ostiolo nigro, brevi, verruciformi prominente; ascis

in quoque perithecio ultra 20, fasciculatis, cylindraceo-fusoideis, breve stipitatis, $55-65 \pm 8-11$; sporidiis distichis vel oblique monostichis, fusoideis, $13-15 \pm 4-5$, bilocularibus, medio subconstrictis, utrinque angustato-rotundatis, hyalinis.

Hab. in caulis vetustis Viciæ Craccæ, Mageerö, Schlucht in Nordcap Norvegiæ.

12. **Sphærella sordidula** Spieg. Fungi Guar. Pug. I, n. 217. — Pe- 2490
ritheciis hinc inde sparsis vel paucis laxe aggregatis, matrice circum circa pallescente vel albicante (ceterum sordide grisea) insidentibus, tectis, vix prominulis, ostiolo minuto, epidermidem perforante, exerto donatis, udis globulosis, siccis cupulato-collapsis, $80-100 \mu$. diam., atris, membranaceis, contextu parenchymatico, pellucide fuligineo, ostiolo subcarbonaceo, opaco; ascis ovatis v. obclavatis, crassis, apice obtuse rotundatis, basi abrupte, minute brevissimeque stipitatis, $30-35 \pm 12-15$, aparaphysatis, 8-sporis; sporidiis di-tri-stichis, elliptico-elongatis, utrinque rotundatis, $12-14 \pm 4-5$, medio 1-septatis, leniter constrictis, loculis 2-guttulatis, æqualibus v. supero vix majore, hyalinis.

Hab. ad siliquas exsiccatas adhuc pendulas Cassiæ speciei cujusdam in dumetis prope Guarapí Bras.

13. **Sphærella baptisiæcola** (Cooke) Sacc., *Sphær. baptisiæcola* Cooke 2491
Grev. XI, pag. 110. — Tecta, punctiformis, caulincola; peritheciis tenuibus, minimis vix prominulis, tectis; ascis clavatis; sporidiis biserialibus, sublanceolatis, uniseptatis, hyalinis, $15 \pm 3,5$.

Hab. in caulis Baptisiæ leucanthæ, Aiken Carolina inf. Amer. bor. — Incertum an huc, an ad *Didymallam* pertineat species, nam Cookeus, more solito, de paraphysibus silet.

14. **Sphærella granulata** Ell. et Ev. in Journ. of Myc. 1886, p. 102. 2492
— Dense gregaria, maculas definite limitatas occupans; peritheciis minutis, $\frac{1}{6}$ mm. diam., epidermide radiatim erupta et perithecia ostendente, non nigrescente tectis, atris, ostiolo papilliformi sub prominulo donatis, nucleo albo farctis; ascis subcylindraceis, circiter 70 \pm 7, subsessilibus, aparaphysatis, octosporis; sporidiis distichis, fusiformibus, vix curvulis, $20-23 \pm 3-3,5$, hyalinis, primo granulosis, continuis, deinde uniseptatis.

Hab. in caulis emortuis Baptisiæ tinctoriæ, Newfield Amer. bor., socia *Sphærella baptisiæcola*.

15. **Sphærella leucophæa** Ellis et Kellerm. in Journ. of Myc. 1886, 2493
p. 3. — Peritheciis punctiformibus, minutis, $70-100 \mu$. diam., in parenchymate foliorum immersis, ostiolis perforatis, longiusculis;

ascis oblongis, sessilibus vel breviter pedicellatis, $75-80 \approx 7-8$; paraphysibus nullis; sporidiis distichis, fusiformibus, 1-septatis, septo leniter constrictis, $18-20 \approx 3\frac{1}{2}$, hyalinis.

Hab. in foliis emortuis Baptisiæ leucophææ, Manhattan, Kansas Amer. bor. — Certe a *Sph. baptisiæcola* distincta.

16. **Sphærella Astragali** Currey Linn. Trans. XXII, n. 362, Journ. of Bot. 1883, p. 136. — Peritheciis sparsis, membranaceis, tectis, globosis, atrofuscis, minutis, hic illic gregariis; ascis clavatis; sporidiis subfusiformibus, uniseptatis, hyalinis, $15-18 \approx 4$.

Hab. in caulis Astragali, in America arctica.

17. **Sphærella pinodes** (Berk. et Blox.) Niessl, Sacc. Syll. I, p. 514, Winter in Hedwigia 1885, pag. 20. — Perithecia numerosissima, dense sparsa, poro pertusa, tenuissime membranacea, sub microscopio fusca, $100-140 \mu$. diam.; ascis oblongo-cylindracei, sessiles vel breviter stipitati, saepe basi parum incrassati, $58-62 \approx 12$; sporidia subdisticha, oblonga, utrinque parum attenuata, saepe inæquilateralia, $14-16 \approx 5$.

18. **Sphærella calycicola** Pass. in Erb. Critt. It. Ser. II, n. 1462. — 2495 Peritheciis solitariis v. gregariis, subglobosis, atris; ascis ovato-piriformibus, apice rotundatis, basi brevissime abrupteque stipitatis, $37-38 \approx 22$; sporidiis triserialibus, oblongo-spathulatis, prope medium septatis, non constrictis, hyalinis, $20 \approx 5$.

Hab. ad dentes calycinios aridos Trifolii angustifolii prope Parma Ital. bor. (PASSERINI).

19. **Sphærella consociata** Rehm in Hedwigia 1885, p. 238. — Pe- 2496 ritheciis plerumque gregariis, subimmersis vel sessilibus, globosis, in papillulam acutam protractis, haud perspicue perforatis, atris, glabris, $0,2-2,5$ mm. diam., in parenchyma fuscum contextis; ascis ellipticis vel pseudo-piriformibus, latis, basi sessilibus, crassis, 8-sporis, $50-60 \approx 18-21$; sporidiis 3-4-stichis, clavatis, subobtusis, 2-cellularibus, cellulis æqualiter longis, medio haud constrictis, rectis, hyalinis, $18-20 \approx 5$; paraphysibus nullis.

Hab. ad caules sicclos Trifolii pallescentis ad glaciem aeternam montis Ortler prope Sulden in Tirolia (DR. REHM).

20. **Sphærella nerviseda** Speg. — Sacc. Syll. I, p. 504 — var. **microspora** Sacc. Misc. Myc. II, pag. 22. — Sporidiis clavulatis, quam in typo brevioribus, $10-12 \approx 4,5$ in ascis clavulatis, $50 \approx 10$, oblique monostichis.

Hab. in foliis emortuis Orobi verni, sociis spermogoniis

Septoria orobicola et *Phyllosticta orobella* pr. Neuchâtel Galliæ (MORTHIER).

21. **Sphærella melanococca** (Lév.) Berl. et Sacc. Add. Syll. p. 406. 2498

Sphæria melanococca Lév. in Ann. Sc. Nat. 1863, p. 296. — Peritheciis epiphyllis, gregariis, discretis, erumpentibus, superficialibus, globosis, depresso-levibus, nigris, opacis, intus albis, ostiolis papillatis; paraphysibus raris, linearibus, continuis; ascis clavato-cylindraceis, sporidiis octonis, ovato-elongatis, medio constrictis, bilocularibus, hyalinis.

Hab. ad folia *Stylosanthis guyanensis* in Nova Grana-ta (LINDIG).

22. **Sphærella Sesbaniae** Ell. et Ev. Journ. Myc. 1888, pag. 98. — 2499

Dense gregaria, erumpens; peritheciis 80-100 μ . diam. pertusis; ascis 35-40 \approx 7-8, aparaphysatis; sporidiis distichis, oblongo-cylindraceis, 1-septatis, haud constrictis, rectis, obtusulis, 10-12 \approx 3,5-4.

Hab. in caule emortuo Sesbaniae macrocarpæ, Louisiana Amer. bor. (LANGLOIS). — Socium adest *Macrosporium* hyphis septatis tortuosis, conidiis oblongis, 2-3-septatis, muriformibus, opacis. Accedit ad *S. granulatum*, sed sporidia multo minora.

23. **Sphærella Desmodii** Winter in Journ. of Myc. I, p. 121. — Peri- 2500

theциis epiphyllis, in maculis magnis, valde irregularibus, arescendo griseis fuscidulisve, margine obscuriori, determinato, varie flexuoso et sinuato circumdati, gregariis, minutis, hemisphæricis, poro simplici pertusis, atris, 70-90 μ . diam.; ascis oblongo-cylindraceis, sursum parum attenuatis, vertice rotundatis, sessilibus, 8-sporis, 35-40 \approx 6-7; sporidiis distichis, cymbiformibus vel fusoideis, sæpe plus minusve curvatis, medio uniseptatis, sed non constrictis, hyalinis, 10,2-12 \approx 2,5-3,5.

Hab. ad folia viva Desmodii canescentis, Missouri.

24. **Sphærella Linhartiana** Niessl in Linhart Fungi Hungar. n. 461. 2501

— Peritheciis numerosissimis, dense disseminatis, epidermide innatis, globosis, ostiolo punctiformi, atris, 100-130 μ . diam.; ascis fasciculatis, clavatis, 45-60 \approx 9-12 et interdum ultra latis, 8-sporis; sporidiis distichis, oblongo-fusiformibus, sæpe curvatis, utrinque obtusis, medio uniseptatis sed vix constrictis, demum guttulis quaternis, hyalinis, 10-12 \approx 3-4.

Hab. in caulis siccis Meliloti albi in Hungaria. — *Sph. leptasca* affinis sed ascis amplis, sporidiis distichis distinguenda.

25. **Sphærella nubigena** Speg. Rev. Myc. 14, pag. 79. — Peritheciis 2502

hinc hinde laxe gregariis, minutis, 100-110 μ . diam., subepidermi-

cis, atris, membranaceis; contextu parenchymatico, olivaceo; ascis cylindraceo-clavatis, brevissime stipitatis, $55-60 \approx 8-10$, octosporis, aparaphysatis; sporidiis distichis ovoideo-subclavulatis, utrinque rotundatis, $15-16 \approx 4$, medio 1-septatis, non constrictis hyalinis.

Hab. in foliis emortuis Drabæ aizoidis in montibus prope Saint-Nigier Galliæ.

26. **Sphærella Passeriniana** Sacc. *Sphærella Cruciferarum* (Desm.?) 2503

Pass. in Erb. critt. it. Ser. II, n. 1279. — Peritheciis punctiformibus, minutissimis, sparsis, atris, epidermide velatis, interdum ostiolo acutiusculo erumpentibus, $100-150 \mu$. diam.; ascis ovatis sessilibus $42-45 \approx 20$, 8-sporis; sporidiis 2-3-serialiter stipatis ovali-elongatis, utrinque obtusis, medio septatis, hyalinis, $15 \approx 5$.

Hab. in caulis et pedunculis aridis Rapistri rugosi atque Erysimi officinalis, Vigheffio, Parma Ital. bor. (PASSERINI). — *Sphærella Cruciferarum* in Rabh. Fungi eur. n. 1852 habitu similis, sed ascii multo minores et sporidia angustiora, fusiformia, apicibus acutis $2\frac{1}{2} \mu$. lata, ideo species præsens diversa et aliter nuncupanda.

27. **Sphærella napicola** Fautrey Rev. Myc. 1890, p. 64. — Maculis 2504

magnis, indeterminatis, nigris; peritheciis numerosis, in maculis gregariis, extus sparsis, rotundatis, nigris; ascis ventricosis; sporidiis hyalinis, 1-septatis, 4-guttatis, loculo supero longiore et latiore, $18-28 \approx 4-6$.

Hab. in caulis siccis Brassicæ Napi var. oleiferæ, Noidan Côte-d'Or Galliæ.

28. **Sphærella pyrenaica** Speg. Rev. myc. 14, p. 78. — Peritheciis am- 2505 phigenis, globoso-lenticularibus, $90-100 \mu$. d., innato-erumpentibus, atris, submembranaceis, contextu parenchymatico, fuligineo; ascis cylindraceo-clavatis, breve stipitatis, $40-45 \approx 8$, apice crassiuscule tunicatis, octosporis, aparaphysatis; sporidiis distichis, cylindraceo-ellipticis, utrinque obtusiuscule rotundatis, $15 \approx 3$, medio 1-pseudo-septatis, non constrictis, hyalinis.

Hab. in foliis emortuis putrescentibus Petrocallidis pyrenaicæ in montibus prope Luchon Galliæ.

29. **Sphærella Cyparissiæ** Passer. Diagn. Fung. nuovi I, n. 29. — 2506

Peritheciis sparsis, epidermide tectis, dein vertice acutiusculo denudatis, subglobosis, atris; ascis obovatis, sessilibus, 8-sporis, $15-17 \approx 7,5-8$; sporidiis subtristicho-stipitatis, oblongo-cuneatis, loculo altero breviore et angustiore, $9-10 \approx 3$, perfecte hyalinis.

Hab. in caule putri Euphorbiæ Cyparissiæ, Vigheffio

- 618 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Sphærella.
 prope Parma. — *Sphærellæ Salicorniæ* affinis, sed ascis minoribus et sporidiorum forma et colore, satis diversa videtur.
30. **Sphærella Tithymali** Passer. Diagn. Funghi nuovi I, n. 30. — 2507
 Peritheciis crebre sparsis, innato-erumpentibus, atris, ostiolo obtuso; ascis cylindricis, 8-sporis, $40-50 \approx 7,5-8$; sporidiis distichis, fusiformibus, medio septatis, hyalinis, $12-15 \approx 2,5-3$, intra ascos tantum visis.
Hab. in caule fracido Euphorbiæ Cyprissiæ, Fornovo in Prov. Parmensi.
31. **Sphærella Thesii** Schroet. Hedw. 1890, p. 59. — Peritheciis epi- 2508
 dermide tectis, extus perspicuis, gregariis, depressis, $250-300 \mu$. lat., ostiolo plano, rotundo, pariete molli, nigro-brunneo; ascis in quoque perithecio numerosis, cylindraceis, breviter stipitatis, apice rotundatis, $40-50 \approx 8-10$, octosporis; sporidiis distichis, fusoideis, deorsum magis attenuatis, utrinque rotundatis, medio non constrictis, $12-13 \approx 4-5$, hyalinis.
Hab. in Thesio intermedio in Torpcider prope Belgrad Serbiæ.
32. **Sphærella tabifica** Prill. et Delacr. Bull. Soc. Myc. 1891, p. 24, 2509
 t. III, f. II. — Peritheciis rotundis, fuscis, erumpentibus, 150μ . lat., poro papillato 35μ . lat. præditis; ascis oblongo-clavatis, obtuse stipitatis, $60 \approx 12$, aparaphysatis; sporidiis hyalinis, didymis, loculo superiore ovato-rotundato, inferiore angustiore, leviter attenuato, septo unico infra medium constrictis, distichis, granulato-guttulatis, $21 \approx 7,5$.
Hab. in petiolis late albo-maculatis Betæ vulgaris cultæ, socia *Phyllosticta tabifica* ejus verisimiliter pycnidio in Mondoubleau, Loir-et-Cher Galliæ.
33. **Sphærella parasitica** Fautrey Rev. Myc. 1890, p. 64. — Ascis 2510
 cylindraceis, 4-6-8-sporis, $45-60 \approx 4-6$, aparaphysatis; sporidiis distichis, inæquilateralibus, 1-septatis, pluriguttatis, $20-25 \approx 5-7$.
Hab. in areis Pucciniæ Malvacearum ad folia Alceæ roseæ ad Noidan Côte-d'Or Galliæ.
34. **Sphærella Saxifragæ** Pass. Diagn. F. nuovi I, n. 19 (1887, 9 Jan.). 2511
 — Peritheciis epiphyllis, sparsis vel subgregariis, globosis, velatis, atris; ascis oblongo-clavatis, $50 \approx 15$, 8-sporis; sporidiis subdistichis vel inordinatis, oblongo-spathulatis, medio septatis, non constrictis, hyalinis, $22,5 \approx 5$, endoplasmate opaco granuloso.
Hab. in foliis emortuis Saxifragæ muscoidis in Colle del Gries Pedemontii.

35. **Sphærella olenjana** Sacc. *S. Saxifragæ* Karst. Symb. XIX. p. 87 2512
 (1887, 5 Mart.) nec Pass. — Peritheciis epiphyllis, raro hypophyllis,
 sparsis aut maculæ nigrescenti gregatiū insidentibus, erumpenti-
 superficialibus, sphæroideo-conoideis, poro pertusis, atris, exiguis;
 ascis sessilibus, oblique ovoideo-elongatis, 40-66 \times 15-18, 8-sporis;
 sporidiis ovoideo-elongatis, medio septatis, haud constrictis, chlori-
 no-hyalinis, 24-28 \times 6-7.

Hab. in foliis languescentibus *Saxifragæ rivularis* prope
 Olenji maris albi in Fennia.

36. **Sphærella ushvaiensis** Speg. F. Fueg. n. 204. — Maculis nullis, 2513
 peritheciis sparsis, epidermide velatis, lenticularibus, 90-100 μ . d.,
 atris glabris coriaceis, opacis, contextu tantum in parte infera distin-
 cte majusculeque parenchymatico fuligineo, ostiolo minuto pertusis;
 ascis cylindraceo-obclavatis, antice obtuse rotundatis vel subtrun-
 cato-rotundatis subcrassiuscule tunicatis, non vel vix 1-foveolatis,
 postice attenuato-rotundatis, 28-32 \times 5, stipite breviusculo crassiu-
 sculoque suffultis, octosporis, aparaphysatis; sporidiis cylindraceo-
 subclavulatis, utrinque acutis sed obtuse rotundatis, 7-9 \times 2, medio
 1-septatis leniter constrictis, loculis minute 2-guttulatis, supero
 vix crassiore, hyalinis.

Hab. in caulis emortuis aridis *Galii antarctici* prope
 Ushuvaia Fuegiæ.

37. **Sphærella Resedæ** Pass. Diagn. Fungi Nuovi III, n. 7. — Peri- 2514
 theciis gregariis matricem infuscantibus, globoso-conoideis promi-
 nulis; ascis clavatis vel obclavatis aut gibbis, basi ventricosis, 45-
 62 \times 15-20, 8-sporis; sporidiis bi-tristichis, oblongis vel oblongo-
 cuneatis, medio septatis non constrictis hyalinis, 15-18 \times 5.

Hab. in caulis putridis *Resedæ luteolæ*, Vigheffio prope
 Parma. — Fructificatione ad *S. carpogenam* accedit, sed peri-
 theciis crassioribus infuscantibus distinguenda.

38. **Sphærella asterinoides** Ell. et Ev. Journ. Myc. 1888, p. 98. — 2515
 Peritheciis sparsis, lenticularibus, latiuscule pertusis, prominulis,
 tectis, basi matrice immersis, mycelio ramoso brunneo plus minus
 distincte cinctis; ascis aparaphysatis, tereti-clavatis, basi breviter
 angustatis, 80-90 \times 18-20; sporidiis stipatis, acute ellipsoideis, 1-
 septatis constrictisque, cellula sup. crassiori, fumose hyalinis, 22-
 28 \times 8-10.

Hab. in caule emortuo *Dipsaci*, Clyde N. Y. Amer. bor. (O.
 F. Cook).

39. **Sphærella xanthicola** Cooke et Harkn. in Grevillea XIV, p. 8. 2516

— Peritheciis punctiformibus, prominulis, atris, in maculas ellipticas congestis, initio epidermide tenui velatis; ascis obclavatis, sessilibus, octosporis; sporidiis ellipticis, uniseptatis, vix constrictis, utrinque rotundatis, hyalinis, 12-15 \approx 5-6.

Hab. in caulis Xanthii in California (HARKNESS).

40. **Sphærella nebulosa** (Pers.) Sacc. Syll. I, pag. 515, var. **Hieracii** 2517

Sacc. et Briard in Rev. Myc. 1885. — Peritheciis numerosis, minutis $1\frac{1}{7}$ - $1\frac{1}{6}$ mm. diam., in maculis griseis sparsis vel gregariis; ostiolis prominulis; ascis sessilibus, octosporis, 40-52 \approx 10-12; sporidiis oblongis, subclavæformibus, apice parum attenuatis, uniseptatis, loculo superiore majore, 16-20 \approx 4-4,5, hyalinis.

Hab. in caulis siccis Hieracii, Troyes Galliæ (BRIARD). —

Var. **Scrophulariæ** Sacc. et Briard in Revue Myc. 1885: Ascis cylindraceo-clavatis, breviter stipitatis, octosporis, 40-48 \approx 8-10; sporidiis distichis, oblongo-fusiformibus, obtusiusculis, 1-septatis, loculis inæqualibus, superiore majore, 12-15 \approx 4,4-5, minute guttulatis, hyalinis. In caulis exsiccatis Scrophulariæ aquatice, Troyes Galliæ (BRIARD). — Var. **Euphorbiæ** Sacc. et Briard in Revue Mycol. 1885: Ascis oblongis, subirregularibus, medio inflatis, sessilibus, octosporis, 36-40 \approx 10-12; sporidiis stipatis, 1-septatis, loculo superiore crassiore, obtuso, 13-14 \approx 4-5, hyalinis. In ramulis Euphorbiæ palustris, Troyes Galliæ (BRIARD).

41. **Sphærella Hieracii** Cooke et Mass. Grev. XV, p. 111. — Peri- 2518

theciis sparsis, epiphyllis, punctiformibus, subconicis, basi immersis; ascis clavatis; sporidiis uniseriatis vel partim biseriatis, ellipsoideo-lanceolatis, apicibus obtusis, 1-septatis, non constrictis, hyalinis, initio 2-nucleatis.

Hab. in foliis subsiccis Hieracii Pilosellæ, Tunbridge Wells Britanniæ (E. G. BAKER).

42. **Sphærella solidaginea** E. et K. Journ. Myc. 1887, p. 127. — Pe- 2519

ritheciis erumpenti-superficialibus, 80-100 μ . diam., subglobosis, laxiuscule parenchymaticis, supra pertusis, sparsis vel gregariis; ascis 35 \approx 9-10; sporidiis distichis, clavato-oblongis, guttulatis, pr. medium subconstrictis, hyalinis, 20 \approx 3,5.

Hab. in foliis emortuis Solidaginis canadensis, Manhattan Amer. bor. (KELLERMAN et SWINGLE).

43. **Sphærella maculicola** Wint. in Hedwigia 1885. — Peritheciis 2520

gregariis in macula determinata, irregulari, pallide fusca, linea obscuriori limitata epiphyllis, immersis, globosis, membranaceis, atris, poro pertusis, 100 μ . latis; ascis elongato-oblongis, fere fu-

siformibus, h. e. utrinque longe attenuatis, substipitatis, 8-sporis, 50-60 \approx 9-12; sporidiis congregatis, elongato-subclavatis, utrinque attenuatis et rotundatis, medio uniseptatis, sed non constrictis, 14-16 \approx 3,5 hyalinis.

Hab. ad folia *Helichrysi* (an petiolati? vel nudifolii?)
Somerset-East, Promont. Bonæ Spei (MAC OWAN).

44. **Sphærella Ritro** Pass. Diagn. F. nuovi I, n. 23. — Peritheciis 2521 sparsis, tectis, subglobosis; ascis ex ovata basi superne attenuatis, 8-sporis; sporidiis tristichis vel congregatis, oblongis, hyalinis, 10-15 \approx 3-5.

Hab. in caule putrescente *Echinopis Ritro*, Vigheffio pr. Parma. — A *Sphærella Compositarum* ascis sporidiisque multo minoribus, et a *Sphærella Jurineæ* ascorum forma distinguenda.

45. **Sphærella aliena** Pass. in Erb. cr. it. ser. II, n. 1278. — Peritheciis 2522 sparsis epidermide velatis, subglobosis, atris, 75-100 μ . diam.; ascis brevibus, clavatis, curvulis subsessilibus, 30-37 \approx 7-8; sporidiis subdistichis vel oblique monostichis, oblongis, hinc paullum attenuatis, medio tenuissime septatis, hyalinis, 15 \approx 3-3½.

Hab. in summo caule arido *Centaureæ nigrescentis*, Collecchio Parma (G. PASSERINI). — A *Sphærella Compositarum* Auersw. plane diversa.

46. **Sphærella majuscula** Cooke Journ. of Bot. 1883, p. 138. — Epiphylla, gregaria sparsave, sæpe maculæformis; peritheciis majusculis, globosis, atris, semiinnatis, prominulis; ostiolo punctiformi; ascis cylindrico-clavatis; sporidiis arcte ellipticis, uniseptatis, loculo inferiore leviter tenuiore, 12 \approx 3.

Hab. in foliis emortuis *Senecionis rotundifolii*, Stewart Island Novæ Zelandiæ (KIRK).

47. **Sphærella hypericina** Ellis Bull. Torr. Bot. Club. IX, p. 74. — 2524 Amphigena; peritheciis exiguis, erumpentibus, hinc illinc 2-6-aggregatis; ascis oblongis; sporidiis inordinatis, clavato-oblongis, uniseptatis, plerumque curvulis, hyalinis, 10 \approx 3.

Hab. in foliis Hyperici prolifici in Amer. bor. (ELLIS).

48. **Sphærella tingens** Niessl in Hedw. 1883, p. 13. — Sparsa vel 2525 subgregaria, matricem rubro vel roseo tingens; peritheciis minutissimis, $1/10$ mm. diam., tectis globosis, membranaceis, fuscis, ostiolo punctiformi; ascis fasciculatis, clavatis, 49-50 \approx 12-15, sessilibus octosporis; sporidiis stipatis, cuneato-cylindraceis, subrectis, inferne leniter angustatis sed utrinque rotundatis, medio 1-septatis, 4-guttulatis, hyalinis, 16-19 \approx 4.

Hab. in foliis vivis Arenariæ ciliatæ, Albulapass Helvetiæ (WINTER). — Macula rubellâ matricali mox inter Sphærellas dignoscenda species.

49. **Sphærella subnivalis** Rehm in Hedwigia 1885, p. 237. — Pe- 2526
ritheciis gregariis in foliis dealbatis sessilibus, fusco-nigris, glo-
bosis, poro minutissimo pertusis, 0,15 mm. diam., in parenchyma
fuscum contextis, glabris; ascis latis, basi sessilibus, sursum at-
tenuatis, crassis, 8-sporis, 40-45 \approx 9; sporidiis clavatis, 2-cellula-
ribus, cellula superiore latiore, plerumque utrinque 2-nucleolatis,
rectis, hyalinis, distichis, 12-15 \approx 3-4,5; paraphysibus nullis.

Hab. ad folia sicca, vix nive liberata, Cerastii latifolii juxta
glaciem æternam montis Ortler prope Sulden in Tirolia (D. REHM).

50. **Sphærella densa** Rostrup Islands Svampe p. 8. — Peritheciis 2527
dense stipatis, amphigenis, rufo-fuscis, sphæroideis; ascis oblongo-
clavatis, 38-42 \approx 7-9, 8-sporis; sporidiis distichis, fusoideo-oblon-
gatis, 1-septatis, hyalinis, 12-14 \approx 4; paraphysibus nullis.

Hab. in foliis Arenariæ norvegicæ in insula Islandia.

51. **Sphærella Alsines** Pass. Diagn. F. nuovi I, n. 10. — Peritheciis 2528
crebre sparsis, punctiformibus, atris, ostiolo acutiusculo, contextu
celluloso, fuligineo; ascis aparaphysatis, obovato-piriformibus vel
gibbis, basi acutis, sessilibus, 8-sporis, 45 \approx 17-20; sporidiis tristi-
chis vel congregatis, oblongis, medio septatis, non constrictis, utrin-
que rotundatis, loculo altero paullo longiore et angustiore, 15-17 \approx 5.

Hab. in foliis inferioribus aridis Alsines laricifoliæ in M.
Prinzera prov. Parma.

52. **Sphærella Caryophylli** Pass. Diagn. F. nuov. I, n. 12. — Peri- 2529
theciis sparsis vel subgregariis, minimis, ostiolo acuto, atris, nitidis,
cellulis ampliusculis fuligineis contextis; ascis oblongo-clava-
tis, 8-sporis, 60-75 \approx 15; sporidiis cuneiformibus, ad septum non
constrictis, 12,5-15 \approx 3,5, hyalinis.

Hab. in vaginis siccis Dianthi Caryophylli et in bracteis
foliisque siccis Dianthi proliferi, Vigheffio pr. Parma.

53. **Sphærella Pulviscula** Cocc. et Mor. Enum. Fung. della provincia 2530
di Bologna Cent. III, p. 27, t. II, f. 36-30. — Peritheciis globo-
sis, perpusillis, 60-72 μ . d., crebre sparsis, granuliformibus, atris,
nitidis; ascis brevibus, obovatis, subsessilibus, 22-26 \approx 9-11, 8-spo-
ris; sporidiis subtristichis, navicularibus, medio septatis, hyalinis,
8-11 \approx 2-2½; paraphysibus nullis.

Hab. in caulinis et foliis Dianthi brachyanthi in Pyre-
næis Galliæ.

54. Sphærella Octopetalæ Oudem. Contr. à la Flore myc. de No. 2531
waja Semlja p. 14, t. I, f. 14. — Peritheciis epiphyllis, subgrega-
riis, minutissimis 50 μ . diam., atris, absque ullo macularum ve-
stigio, parte basilari in foliorum parenchymate absconditis, api-
cali vero supra epidermidem prominente, e parenchymate laxo, fu-
scescenti-nigro contextis; ascis maturis, 70-80 \approx 25, membrana sur-
sum præsertim gelatinoso-incrassatis, 8-sporis, paraphysibus defi-
cientibus; sporidiis hyalinis, di-vel tri-stichis, oblongo-obovatis,
21 \approx 7, septatis, ad septum levissime constrictis, parte dimidia an-
tica latiore, subhemisphærica, circa 9 μ ., postica paululum angu-
stiore, magis oblonga, circa 12 μ . longa.

Hab. in foliis Dryadis octopetalæ, Nowaja Semlja.

55. Sphærella fuegiana Speg. F. Fueg. n. 197. — Maculis nullis; 2532
peritheciis hypophyllis, sparsis, lenticularibus, 90-100, atris gla-
berrimis, membranaceo-coriacellis, contextu dense parenchymatico
olivaceo; ascis cylindraceo-obclavatis, apice rotundato-subtruncatis
crasse tunicatis non vel vix 1-foveolatis, basi abrupte minuteque
coarctato-stipitatis, 30-40 \approx 8-10, octosporis, a paraphysatis; spori-
diis elliptico-obovatis, utrinque obtusiusculis, 7,5-8,5 \approx 3-4, loculo
supero nonnihil crassiore brevioreque, hyalinis.

Hab. ad folia Acænæ adscendentis languida rarissime in
Shammacus Fuegiæ.

56. Sphærella Potentillæ Oudem. Contr. à la Flore myc. de No. 2533
waja Semlja p. 15, t. I, f. 15. — Peritheciis in foliis aridiis ho-
spitantibus.....; ascis inæquilateralibus, oblongo-clavatis, apa-
raphysatis, 70-80 \approx 14, membrana ubique æquali, vulgo 20 in eodem
perithecio; sporidiis distichis, rectis, hyalinis, oblongo-obovatis, 1-
septatis, 18-23 \approx 5-7, parte dimidia anteriore latiore, rotundata,
posteriore multo angustiore, ad altitudinem septi paululum con-
strictis.

Hab. in foliis aridis Potentillæ fragiformis, Nowaja Semlja

57. Sphærella Earliana Winter in Journ. of Myc. pag. 101. — Pe- 2534
ritheciis amphigenis, densissime stipatis, greges parvos, angulato-
rotundatos, 1-2 mm. latos, nigros formantibus, minutissimis, glo-
bosis, poro simplici pertusis, atris, 60-70 μ . diam.; ascis fascicu-
latis e basi subventricosa sursum parum attenuatis, brevissime sti-
pitatis, 8-sporis, 26-30 \approx 7; sporidiis inordinatis clavatis, medio uni-
septatis, non constrictis, hyalinis, 8 \approx 2; paraphysibus nullis.

Hab. in Fragaria culta, Anna Amer. bor. (F. S. EARLE). —
Ascis et sporidiis a *S. Fragariæ* distinguenda.

58. **Sphærella Serpylli** Pass. Diagn. Funghi nuovi, I, n. 26. — Pe- 2535
ritheciis minimis, superficialibus, globosis, solitariis vel subgregariis
atris; ascis oblongis, subsessilibus, basi plus minus inflatis v. gib-
bis, octosporis, $37-45 \approx 10-12$; sporidiis distichis vel inordinatis,
oblongis, medio septatis et leniter constrictis, hyalinis, endoplasma-
te granuloso opaco, $15-20 \approx 5-7$.

Hab. in calycibus siccis Thymi Serpylli, Vigheffio prope
Parma. — A *Sphærella calycicola* cui affinis, characteribus al-
latis videtur distinguenda.

59. **Sphærella Pectinis** Pass. Diagn. F. nuov. III, n. 9. — Peritheciis in 2536
matrice fuscescente crebre sparsis, punctiformibus, atris, ostiolo
acutiusculo; ascis ovatis, basi abrupte breviter stipitatis, 8-sporis,
 $40-42 \approx 12,5-15$; sporidiis stipatis oblongis, utrinque rotundatis,
medio septatis, $12,5 \approx 5$.

Hab. in fructibus dejectis Scandicis Pectinis-Veneris,
Tornovo in prov. Parmensi.

60. **Sphærella Eryngii** (Fr.) Cooke Handb. n. 2761, Sacc. Syll. I,
p. 511, Winter in Hedwigia 1886, p. 19. — Ascii $40-44 \approx 7$; spo-
ridia cylindrico-fusiformia, $19-21 \approx 2,5$.

61. **Sphærella Pachypleuri** Fuck. in Heuglin Reisen Nordpol. vol. III 2537
et Oudemans Contr. Flor. Myc. Now. Seml. pag. 6, t. II, f. 2. —
Peritheciis gregariis sparsisve, minutis, aterrimis, semiliberis, e
basi globosa antice obtuse conicis, ostiolis demum perforatis; ascis
fasciculatis, oblongis, sessilibus, 8-sporis, $52 \approx 10$; sporidiis disti-
chis, oblongo-subclavatis, rectis, inæqualiter didymis, loculis gut-
tulatis, hyalinis, $16 \approx 4$; spermatiis inter ascos in sporophoribus bre-
vioribus ortis, anguste fusiformibus, curvatis, 2-4-guttulatis seu
obscure 2-3-septatis, antice abrupte ciliatis, cilio spermatii dimi-
diā partem superante, $24 \approx 2$.

Hab. in Pachypleuro alpino exsiccatō in insula Nowaja
Semlja.

62. **Sphærella patagonica** Speg. F. Pat. n. 124. — Maculis nullis sed 3538
toto folio ex albo griseo-arescente; peritheciis sparsis v. hinc inde
laxe aggregatis, subepidermicis, minutis, lenticulari-globosis, $70-90$
 μ . diam., tenuissime membranaceis, grosse parenchymatico-cellulo-
sis, ostiolo rotundo perforatis; ascis e globoso ovatis, apice sub-
acutato-rotundatis crassissimeque tunicatis, basi stipite crasso bre-
vissimo (vix evoluto) suffultis, $20 \approx 16$, octosporis, aparaphysatis;
sporidiis inordinate polystichis, elliptico-clavulatis, utrinque ob-
tusiuscule rotundatis, 8-9 $\approx 2-3$, medio 1-septatis, non vel vix

subconstrictis, loculo supero nonnihil crassiore brevioreque, hyalinis.

Hab. ad folia exsiccata emarcida *Bolacis glebariae* in Gregory bay Patagoniæ.

63. **Sphaerella brionnensis** Sacc. et Malbr. Fung. Gall. Ser. V, n. 2159. 2539

— Peritheciis in foliis arescendo expallentibus, subsparsis, saepius epiphyllis, punctiformibus, globulosis, papillulatis, initio epidermide tectis, nigris; ascis tereti-fusoideis, subsessilibus, aparaphysatis, 35-45 \approx 6-7, octosporis; sporidiis oblongo-fusoideis, subrectis, 15-18 \approx 4-5, 4-guttulatis, medio dein septatis, non v. vix constrictis hyalinis.

Hab. in foliis arescentibus *Angelicae silvestris*, Brionne Eure Galliæ.

64. **Sphaerella adusta** Fuck. in Reis. nach Nordpolarmeer et Oud. 2540

Contr. Fl. Myc. Now. Seml. p. 151, tab. II, fig. 3. — Peritheciis in maculis fusco-nigris, rugulosis, adustis, quandoque totum folium caulemque occupantibus, gregariis, minutis, atris, depresso globo-sis, ostiolo prominulo papilliformi, perforato; ascis fasciculatis, sessilibus, 8-sporis, 56 \approx 16; sporidiis distichis, oblongo-clavatis, rectis, antice obtusissimis, basi acuminatis, valde inæqualiter didymis, septo in sporidii angustiore parte, ad septum non constrictis, hyalinis, 16 \approx 6.

Hab. in foliis et caulibus siccis *Epilobii latifolii*, Nowaja Semlja.

65. **Sphaerella Enotheræ** Ell. et Ev. in Journ. of Myc. 1885, p. 151. 2541

— Peritheciis erumpentibus, hemisphæricis, 90-100 μ . diam., sursum late pertusis, dense gregariis et saepe in series latas in valvis capsularum dispositis; ascis oblongis, sursum leniter attenuatis, basi abrupte contractis atque in pedicellum breve desinentibus; sporidiis distichis, forma variis, oblongis vel ovato-oblongis, 10-15 \approx 3, nucleatis vel oblongo-fusiformibus, vix curvulis, obscure 1-septatis, 15-20 \approx 3-3,5.

Hab. in capsulis Enotheræ biennis, Newfield Amer. bor.

66. **Sphaerella depazeæformis** (Auersw.) Sacc. Syll. I, p. 513, Winter in Hedwigia 1886, p. 20. — Huc ducenda sec. Winter (Hedw. l. c.) *Læstadia Oxalidis* (Rabenh.) Sacc. Syll. I, p. 429, n. 1635.

67. **Sphaerella Clematidis** Oud. Contr. Fl. Myc. Pays-Bas XI, p. 21. 2542

— Peritheciis sub epidermide nidulantibus, dense sparsis, maturis atris, poro minuto pertusis, $1/10$ mm. latis, parenchymatico-membranaceis; ascis clavatis, paullum curvatis, 60-70 \approx 14; sporidiis

hyalinis, utrinque obtusissimis, prope medium 1-septatis, 14 \approx 5 tristichis (?).

Hab. in caulis Aristolochiæ Clematitis, prope Harlem Hollandiæ.

68. **Sphærella intermixta** Niessl in Hedw. 1882, p. 112. — Peritheciis 2543 gregariis, minutissimis, circa 150 μ . diam., ostiolo punctiformi, membranaceis, atris, glabris, nitidis; ascis numerosis, cylindraceis, clavatis, stipite brevi, 40-50 \approx 8-9, 8-sporis; sporidiis distichis, cuneatis, superne late rotundatis, inferne attenuatis, rectis vel parum curvatis, medio uniseptatis, hyalinis, 9-12 \approx 3.

Hab. in foliis subsiccis Campanulæ Zoisii in alpinis Carnioliae (Voss).

69. **Sphærella Cyananthi** Pat. in Rev. Myc. 1886, p. 180. — Peritheciis 2544 minutis, numerosis, nigris vel brunneo-rufis; ascis ovoides, 33 \approx 20, sessilibus; sporidiis uniseptatis, hyalinis, 13 \approx 5, demum viridulis, uno apice attenuatis.

Hab. in caulis emortuis Cyananthi, Yunnan Sinarum (Ab. DELAVAY).

70. **Sphærella Euphrasiæ** Pass. Diagn. Funghi nuovi, I, n. 25. — Peritheciis 2545 minutis, sparsis, lenticularibus, fuscis, primo tectis, dein per epidermidem fissam emersis, nucleo albo farctis; ascis brevibus, subcylindricis, basi breviter angustatis, circiter 47-50 \approx 12-13, 8-sporis; sporidiis subdistichis, oblongis, didymis, leniter constrictis, loculis inæqualibus, altero subcuneato angustiore, 12,5-15 \approx 6-7,5, hyalinis.

Hab. in caulis siccis Euphrasiæ luteæ in alveo Bagazza, Vigheffio prope Parma.

71. **Sphærella phlogina** Ell. et Ev. Journ. Myc. 1888, p. 65. — Peritheciis 2546 sparsis, erumpenti-superficialibus, conico-hemisphæricis, $\frac{1}{6}$ mm. diam., epiphyllis; ascis ovato-oblongis, 65-75 \approx 20-22, apophysatis; sporidiis clavato-oblongis, 1-septatis leniter constrictis, utrinque rotundatis, flavidо-hyalinis, plurinucleatis, 20-30 \approx 7-9.

Hab. in foliis emortuis Phlogis longifoliæ, Belt Mts. Montana Amer. bor. (ANDERSON). — Affinis *S. Stellinearum*.

72. **Sphærella Mariæ** Sacc. et Bomm. in Bull. Soc. R. Bot. Belg. 2547 1886, p. 173. — Peritheciis innatis, sparsis, numerosissimis, nigris, epiphyllis, prominentibus, 90-120 μ . diam.; ascis cylindro-claviformibus; sporidiis distichis, hyalinis, elongatis, uniseptatis, loculis inæqualibus, 4-guttulatis, ad septum constrictis 14-18 \approx 4,5-5.

Hab. in foliis emortuis *Digitalis luteæ*, Hastières Belgii.

— Affinis *Sph. silvaticæ*.

73. **Sphærella ciliata** Ell. et Ev. Proc. Phil. 1890, p. 231. — Peritheciis subglobosis, 150 μ . diam., epidermide tectis, sed matrice haud immersis, ostiolo papillato; ascis clavato-cylindraceis, 40 \times 7; sporidiis biseriatis, clavato-oblongis, hyalinis, 1-septatis, constrictisque, 10-13 \times 3.

Hab. in caulis emortuis *Steironematis ciliati*, London Canada Amer. bor. (DEARNESS).

74. **Sphærella magellanica** Speg. F. Pat. n. 125. — Maculis nullis 2549 sed matrice tota sordide griseo-arescente; peritheciis hinc inde sparsis, subepidermicis, globuloso-lenticularibus, 120-130 μ . diam., prominulis, atris, glabris, ostiolo rotundo donatis, membranaceis, contextu parenchymatico, minute celluloso, atro-fuligineo; ascis cylindraceis vel cylindraceo-obclavatis, subpachydermaticis, apice subtruncato-rotundatis, basi abrupte in stipitem brevissimum nodulosum productis, 48-55 \times 10-15, octosporis, a paraphysatis; sporidiis recte di-tri-stichis, elliptico-clavulatis, utrinque plus minusve obtuse rotundatis 15-17 \times 5-6, medio 1-septatis leniterque constrictis, loculo supero nonnihil crassiore, breviore acutioreque, hyalinis.

Hab. in scapis dejectis putrescentibus *Armeriæ andinæ* pr. Gregory bay Patagoniæ.

75. **Sphærella sycicola** Ell. et Ev. Journ. Myc. 1887, p. 45. — Maculis amphigenis, minutis, 1-2 mm. diam., sordide albis, suborbicularibus vel partim venulis foliorum limitatis, centro tenuibus et translucidis; peritheciis paucis, 1-3, saepius uno tantum in centro maculæ, epiphyllis, nigro-nitidulis, 100 μ . diam., sublenticularibus, latiuscule perforatis, grosse cellulosis; ascis tereti-oblongis, 40-50 \times 6-7, sessilibus; sporidiis biseriatis, ovato-oblongis, hyalinis, 1-septatis leniterque constrictis, 8-10 \times 3-3,5, utrinque rotundatis vel subacutis.

Hab. in foliis vivis *Sycii angulatae*, Missouri Amer. bor. (GALLOWAY).

76. **Sphærella Belladonnæ** Br. et Har. Rev. Myc. 1891, p. 15. — Maculis albis, oblongis, irregularibus, ambitu angulosis, usque ad 3-7 mm. diam., brunneo-cinctis; peritheciis epiphyllis, innatis, subepidermicis, sparsis, plus minus adproximatis, punctiformibus, brunneis, 200-250 μ . diam.; ascis cylindraceo-clavatis, breviter pedicellatis, 8-sporis, 52-60 \times 12-14, p. sp.; sporidiis distichis, ellipsoi-

- 628 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Sphærella.
deo-fusoideis, obtusiusculis, 1-septatis, ad septum constrictis, loculis subæqualibus, hyalinis, 20-24 \approx 6-8.
Hab. in foliis vivis Atropæ Belladonnæ, Pont-sur-Seine, Aube Galliæ (P. HARIOT).
77. **Sphærella Gunneræ** Spieg. F. Fueg. n. 198. — Maculis nullis v. 2552 pallescentibus, indeterminatis; peritheciis epiphyllis, sparsis, immerso-prominulis, e conico lenticularibus, 80-90 μ . diam., glabris, atris, coriacellis, contextu atro indistincto, ostiolo minuto perforatis; ascis cylindraceo-clavatis, sessilibus, antice rotundatis, tenui tunicatis, postice attenuato-rotundatis, 45 \approx 9, octosporis, a paraphysatis; sporidiis elliptico-subclavulatis, utrinque obtusiusculis, 8-9 \approx 3, medio septatis, non constrictis, hyalinis.
Hab. ad folia viva vel languida Gunneræ lobatae in pratis uliginosis, Basket island Fuegiæ.
78. **Sphærella Pirolæ** Rost. Fungi Groenl. p. 551. — Peritheciis am. 2553 phigenis in maculis rufescentibus, subzonatis, magnis, prominulis; ascis cylindraceo-clavatis, 50-60 \approx 8-10; sporidiis oblongo-fusoides, 15 \approx 4.
Hab. in foliis vivis Pirolæ grandifloræ in Groenlandia.
79. **Sphærella Deschmannii** Voss Mycol. Cara. III, p. 156. — Maculis 2554 flavis vel rubris, rotundato-ellipticis, difformibus, circ. 5-10 mm. diam. vel 10-20 mm. long., 10 lat., interdum effusis et foliis magnam partem occupantibus; peritheciis dense gregariis, sessilibus, punctiformibus, globosis, parenchymatico-contextis, atris; ascis fasciculatis, cylindraceis vel anguste fusoideis, in stipitem brevem productis, apice rotundatis, 4-8-sporis, 30-35 \approx 6-11; sporidiis inordinate tristichis, fusoideis, utrinque rotundatis, rectis vel curvulis, didymis, medio non constrictis, loculo superiori parum latiore, guttulatis, demum hyalinis, 21-23 \approx 3.
Hab. ad folia arida Gentianæ Pneumonanthis in Carniolia super.
80. **Sphærella Ebuli** Rich. Cat. Champ. Marn. n. 1221. — Peritheciis 2555 sparsis, epidermidem subgriseam reddentibus; ascis clavatis; sporidiis 1-septatis, ad septum subconstrictis.
Hab. in caulinis emortuis Sambuci Ebuli in Aulnay-l'Aître, Lisse etc., Marne Galliæ.
81. **Sphærella Alyxiæ** C. et M. Grev. XVI, p. 5. — Amphigena; peritheciis 2556 gregariis, innatis, subglobosis, epidermide denigrata tectis, demum collabascentibus; ascis clavato-cylindraceis, octosporis; sporidiis lanceolatis, uniseptatis, hyalinis, 25 \approx 7.

Hab. in foliis emortuis Alyxiæ buxifoliæ, Brighton Victoria Australiæ (F. CAMPBELL).

82. **Sphaerella pandurata** Ell. et Ev. in Bull. Torr. bot. Club. 1883, 2557 p. 117. — Hypophylla; peritheciis globosis, 0,17-0,25 mm. diam., parenchymate foliorum immersis, ostiolo papilliformi minuto, epidermide nigriscente tecto, rotundato, prominulis; ascis oblongo-cylindraceis, 50 \times 7; sporidiis distichis, fusiformibus, 4-nucleatis, flaveolis, medio constrictis, loculis ad septum turgidioribus, 10-12 \times 3.

Hab. in foliis delapsis, Plainfield Amer. bor. (G. F. MESCHUTT).

83. **Sphaerella analoga** Sacc. et Paol. Myc. Malac. n. 103, t. II, f. 3. — 2558 Maculis amphigenis sed distinctius epiphyllis, inæqualiter circularris, albis, anguste fusco-marginatis; peritheciis epiphyllis, lenticularibus, punctiformibus, minute pertusis, 100 μ . diam.; ascis oblongis, brevissime apiculato-stipitatis, apice obtuse attenuatis crasseque tunicatis, 50 \times 15, 8-sporis; sporis di-tristichis, oblongo-retibus, apice rotundatis, deorsum leniter attenuatis, uniseptatis, vix constrictis, 15-16 \times 3.

Hab. in foliis languidis ignotis (ovatis, coriaceis, subintegris, myrtoideis), Malaca (SCORTECHINI).

84. **Sphaerella cyclogena** Sacc. et Paol. Myc. Malac. n. 103, t. II, f. 2. — 2559 Maculis amphigenis, circularibus, pallide rufo-ochraceis, 2-3 mm. diam., anguste fusco-cinctis; peritheciis laxe gregariis, punctiformibus, lenticularibus, atris, pertusis, 100-120 μ . diam.; ascis fusoides, sæpe curvulis, 70 \times 11-12, aparaphysatis, octosporis; sporidiis distichis, oblongis, utrinque obtusiusculis, uniseptatis non constrictis, 4-guttulatis, hyalinis.

Hab. in foliis languidis plantæ ignotæ (caule quadrangulari, spinis axillaribus, foliis oppositis, amplis simplicibus), Malaca (SCORTECHINI). — Maculis eumorphis fulvis distincta.

II. Arboricolæ.

85. **Sphaerella Cercidis** Pass. in Erb. Critt. Ital. Ser. II, n. 1463. — 2560 Peritheciis hypophyllis, sparsis vel laxe gregariis, punctiformibus, tectis dein denudatis, atris; ascis forma et longitudine variis, cylindricis, clavatis aut medio ventricosis; sporidiis oblongo-cuneatis, medio tenuissime septatis, vix vel minime constrictis, 17-20 \times 7 $\frac{1}{2}$.

Hab. ad folia humi strata Cercidis japonicæ in Horto Botanico Parmensi Ital. bor. (PASSERINI).

86. **Sphaerella vesicaria** Pass. in Erb. Critt. It. Ser. II, n. 1367. — 2561 Peritheciis superficialibus, sparsis vel subgregariis, subglobosis,

atris, ostiolo papillari, cellulis amplis, fuscis contextis; ascis oblongis, basi saccatis, breviter et abrupte stipitatis, superne attenuatis, $75 \approx 25$; sporidiis subtristichis oblongis, medio septatis, non vel vix constrictis, loculo altero leniter attenuato, hyalinis, $17,5 \approx 7,5$.

Hab. in leguminibus aridis Coluteæ arborescentis, Parma Italiæ borealis.

87. **Sphærella Sophoræ** Winter Contr. Fl. Myc. Lus. V, pag. 13. — 2562

Peritheciis in macula pallida, exarida, fusco-atrocincta, 2-5 mm. lata, rotundata, amphigena immersis, ostiolo punctiformi perforatis, globosis, fuscis, $90-110 \mu$. diam.; ascis elongato-oblongis, sessilibus, sursum vix attenuatis, rotundatis, 8-sporis, $56-68 \approx 15-16$; sporidiis subdistichis, oblongis, didymis, hyalinis, medio profunde constrictis, inæquilateralibus, cellula superiori latiori, guttulatis, $15-17 \approx 7-8$.

Hab. in Sophoræ sp. foliis vivis, Coimbra Lusitaniæ (MOLLER).

88. **Sphærella Macowiana** Winter Ex. Pilze p. 9, in Flora 1884. — 2563

Peritheciis densissime gregariis, greges forma varia, a nervis limitatos, maculiformes, cinereos, angulatos formantibus, minutissimis, valde prominentibus, globosis, atris, ostiolo punctiformi, 70μ . diam.; ascis oblongo-cylindraceis, interdum deorsum parum inflatis, sessilibus, 8-sporis, $35-44 \approx 9-11$; sporidiis distichis, oblongis, utrinque parum attenuatis, uniseptatis, ad septum vix constrictis, sæpe inæquilateralibus et parum curvatis, hyalinis, $12-14 \approx 3,5$.

Hab. in foliis languidis Melianthi majoris in Promontorio Bonæ Spei Afr. austr. (MAC OWAN).

89. **Sphærella Hesperidum** Penz. et Sacc. Contr. Fungh. agrum II, 2564

p. 12. — Peritheciis laxe aggregatis, amphigenis, primum epidermide velatis, dein prorumpentibus, in arida foliorum macula incidentibus, subglobosis, aterrimis, $130-160 \mu$. d.; ascis clavatis, subsessilibus, apice truncatis, crasse tunicatis, $50-70 \approx 14-16$, octosporis; sporidiis distichis, fusoideis, $16-18 \approx 4-5$, uniseptatis, medio haud constrictis, hyalinis.

Hab. in foliis vivis Citri Limonum, Valle di Latte prope Ventimiglia Ital. bor.

90. **Sphærella assimilata** J. Kunze. — Sacc. Syll. I, p. 486, Winter in Hedwigia 1886, p. 21. — Sporidia $8-10 \approx 2,5-3$.

91. **Sphærella curva** Karst. in Hedw. 1883, n. 12, p. 3, Rev. atque 2565 Synop. Ascom. fenn. p. 65. — Peritheciis?; ascis oblongatis, curvulis, infra medium inæqualiter inflatis $46 \approx 14$; sporidiis ovoideo-

oblongatis, uniseptatis, ad septum leviter constrictis, rectis, dilute luteolis, 12, \approx 4-5.

Hab. in foliis emortuis *Ribis alpin i* ad Mustiala Fenniæ.

92. **Sphærella maculiformis** (Pers.) Auersw. Sacc. Syll. I, p. 477. — 2566

Var. **Pseudoplatani** Pass. in litt. Brun. in Rev. Myc. 1886, p. 205.

— Peritheciis subglobosis, minutis, valde approximatis, hypophyllis; ascis numerosissimis, cylindraceis, 50 \approx 5; sporidiis?....

Hab. in foliis *Aceris Pseudoplatani*, Saintes Galliæ.

93. **Sphærella corylina** Karst. in Hedw. 1884, n. 1, pag. 4, Rev. 2567

atque Synop. Ascom. fenn. p. 64. — Peritheciis hypophyllis per totam paginam dispersis, initio epidermide tectis, dein liberis, sub-superficialibus conoideo-sphæroideis, poro pertusis, atris, 60-80 μ . diam.; ascis cylindraceo-elongatis, vulgo curvulis, sæpe prope medium leviter inflatis, 30-44 \approx 8; sporidiis 8-nis, distichis, oblongatis vel ovoideo-oblongatis, uniseptatis, eguttulatis, 8-14 \approx 2-4.

Hab. in foliis *Coryli Avellanæ* ad Mustiala Fenniæ.

94. **Sphærella dendroides** Schw. Syn. Carol. n. 221, Journ. of. Bot. 2568

1883, p. 108. — Epiphylla aggregata, astoma, maculas maximas cinereas dendroides exhibens; peritheciis nigris erumpentibus; ascis saccatis, late clavatis; sporidiis lanceolatis, uniseptatis, hyalinis, loculo inferiore paullo tenuiore, leniter constrictis, 24-25 \approx 4.

Hab. in foliis *Caryæ*, Amer. bor.

95. **Sphærella Firmianæ** Pass. Diagn. Fungi nuovi, I, n. 13. — Pe- 2569

ritheciis laxe gregariis, minutis tectis, ostiolo acuto erumpente, atris; ascis cæspitosis oblongis, infra medium plus minus inflatis et sæpe inæquilateralibus, basi constricta stipitatis, 50-65 \approx 14-15; sporidiis subdistichis, oblongis, medio septatis, loculo altero vix angustiore, 18-20 \approx 7-7,5.

Hab. in petiolis putrescentibus *Firmianæ* (*Sterculiæ*) platanifoliæ in Horto botanico, Parma.

96. **Sphærella succedanea** Pass. Diagn. Fung. nuovi, I, n. 15. — Pe- 2570

ritheciis epiphyllis in maculis irregularibus, exaridis subgregariis, globosis vel lenticulari-depressis, atris; ascis fasciculatis, ovato-oblongis vel gibbis aut clavatis, basi breviter constricto-stipitatis, 8-sporis, 45-62 \approx 15-20; sporidiis bi-tristichis vel congregatis, medio septatis, non constrictis, hyalinis, 15-20 \approx 6-7.

Hab. in foliis languidis *Plasmoparâ* jam invasis *Vitis viniferae*, sociâ *Phoma succedanea*, Parma. — Peritheciis semper epiphyllis, nucleo durissimo parentibus et sporidiis angustioribus non constrictis, a *Sphærella Vitis* videtur distinguenda.

97. **Sphærella japonica** Pass. Diagn. Funghi nuovi, I, n. 16. — Pe. 2571
 ritheciis amphigenis, gregariis vel sparsis, epidermide tectis punctiformibus, non pertusis, atris, contextu obscure celluloso; ascis fasciculatis, non paraphysatis, oblongo-clavatis vel infra medium inflatis, 8-sporis, 62-85 \approx 10-20; sporidiis subdistichis oblongis, prope medium septatis, non constrictis, hyalinis, 20-25 \approx 5-7.

Hab. in foliis putrescentibus *Evonymi japonici* in Horto botanico, Parma.

98. **Sphærella Mayteni** Speg. F. Fueg. n. 199. — Maculis nullis v. 2572
 totum folium ambientibus arescenti-cineraceis, indeterminatis; peritheciis hypophyllis, laxe gregariis, epidermide velatis, prominulis e lenticulari conoideis 100-120 μ . atris, glabris, coriacellis, contextu parenchymatico fuligineo, ostiolo minuto perforatis; ascis obclavatis, antice truncato-rotundatis, crassiuscule tunicatis, postice abrupte coarctatis ac pedicello parvulo noduloso suffultis, 50-55 \approx 10, octosporis, a paraphysatis; sporidiis distichis elliptico-elongatis subclavulatis utrinque obtusis, 12-13 \approx 4, medio 1-septatis non v. vix constrictis, hyalinis.

Hab. ad folia emortua adhuc pendula *Mayteni* prope Shammacus Fuegiæ. — A *Sph. Aiacu* Speg. toto cœlo diversa!

99. **Sphærella Aiacu** Speg. F. Fueg. n. 193. — Maculis marginalibus, 2573
 majusculis determinatis, fuscescentibus, linea incrassatula rufescente cinctis; peritheciis epiphyllis densissime gregariis, minutis, subglobosis, 70-80 μ ., epidermide velatis, prominulis, glabris, atris, coriacellis contextu parenchymatico olivaceo, ostiolo impresso donatis; ascis fusoideo-obclavatis, apice truncato-rotundatis subcrassiuscule tunicatis, late 1-fovelatis, basi in pedicellum brevissimum crassumque cuneato-productis, 30-35 \approx 5, octosporis, a paraphysatis; sporidiis clavulato-acicularibus utrinque obtusiuscule acutatis 10-12 \approx 2, medio 1-septatis non constrictis, hyalinis.

Hab. ad folia languida *Mayteni* prope Picton island Fuegiæ.

100. **Sphærella Mygindæ** Winter Contr. Fl. Myc. Lus. V, p. 13. — 2574
 Peritheciis densis, gregariis, macula cinerea, rotundato-difformi, 1-10 mm. lata, interdum effusa et folii magnam partem occupante, insidentibus, primo immersis, demum superficialibus, punctiformibus, globosis, poro pertusis, atris, 80-90 μ . diam.; ascis elongato-obclavatis, in stipitem brevem productis, 36-48 μ . longis, infra medium 7 μ . latis, sursum usque 5 μ . attenuatis; sporidiis fusiformibus, saepe parum inaequilateralibus, bicellularibus, cellula superiori parum latiori, hyalinis, inordinate distichis, 10-12 \approx 2,5.

Hab. ad Mygindæ pallentis folia arida, Coimbra Lusitanæ (MOLLER).

101. **Sphærella Aloysiæ** Passer. Diagn. nuov. Funghi I, n. 27. — Pe- 2575
ritheciis minutis, sparsis, epidermidi immutata adnata, atris; ascis
oblongo-clavatis, basi abrupte breviter stipitatis, 8-sporis, 50-70
 \approx 15; sporidiis distichis, oblongis, didymis, vix constrictis, vase
pluriguttulatis, hyalinis, 17-22,5 \approx 7,5.

Hab. in ramulis siccis Aloysiæ in Horto botanico, Parma.

102. **Sphærella spinicola** Passer. Diagn. Fung. nuovi I, n. 28. — Pe- 2576
ritheciis crebre sparsis, superficialibus, minimis, punctiformibus,
atris; ascis ventricosis aut saccatis, 25-30 \approx 7,5-12; sporidiis disti-
chis v. conglobatis, oblongis, utrinque rotundatis, medio septatis,
vix v. non constrictis, hyalinis, 10 \approx 3.

Hab. in spinis Hippophaes rhamnoidis in alveo Tari, For-
novo in prov. Parmensi.

103. **Sphærella Embothrii** Speg. F. Fueg. n. 196. — Maculis nullis v. 2577
vix manifestis, totum folium ambientibus, cinerescentibus; perithe-
ciis hypophyllis, dense aggregatis, immersis, epidermide velatis vix
prominulis, e conico lenticularibus, 50-70 μ . diam., coriacello mem-
branaceis, contextu parenchymatico-olivaceo parum distincto, ostio-
lo impresso perforatis; ascis obclavatis, antice truncato-rotundatis,
postice abrupte coarctatis ac pedicello brevissimo, crassiusculo suf-
fultis, 25-30 \approx 10, octosporis, apophysatis; sporidiis elliptico-cla-
vulatis, utrinque obtusiusculis, 8 \approx 3, medio 1-septato-constrictis,
loculis subcoarctatis, hyalinis.

Hab. ad folia languida v. dejecta Embothrii coccinei in
insula Aicina, Beagle channel Fuegiæ.

104. **Sphærella Banksiæ** Cooke et Mass. Grev. XVI, p. 114. — Epi- 2578
phylla; maculis nullis; peritheciis gregariis, subinnatis, atris, poro
pertusis; ascis clavatis, octosporis; sporidiis biseriatis, ellipticis,
didymis, hyalinis, 12 \approx 5.

Hab. in foliis languidis Banksiæ integrifoliæ, Melbourne
(Miss CAMPBELL).

105. **Sphærella erysiphoides** Sacc. Mich. II, p. 604. — Peritheciis 2579
gregariis in pag. inf. dealbata innato-superficialibus, globoso-de-
pressis, 90-100 μ . diam., pertusis; ascis tereti-clavatis, 40-45 \approx 7,
apophysatis, octosporis; sporidiis distichis oblongo-clavulatis, 9-
10 \approx 2,5-3, 1-septatis non constrictis, granulosis, hyalinis.

Hab. in foliis languidis Tecomæ radicantis, Saintes, sociâ

- 634 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Sphærella.
forma spermogonica *Phyllosticta erysiphoides*, Saintes Galliæ (BRU-
NAUD). — Habitus fere *Erysiphes*.
106. **Sphærella Capreolatæ** Passer. Diagn. Fung. nuovi IV, n. 6. — 2580
Peritheciis amphigenis, in maculis exaridis, subdiscoideis, rubrocinctis gregariis vel subsparsis, punctiformibus, atris, nitidulis; ascis tubulosis v. clavatis v. medio inflatis, 40-50 ≈ 8-15; sporidiis elongatis, hyalinis, medio septatis, utrinque rotundatis, 15-17 ≈ 5.
Hab. in foliis languidis Bignoniæ capreolatae, socio *Glæo-*
sporio Bignoniæ in Horto bot. Parma.
107. **Sphærella staphylina** E. et E. Journ. Myc. 1887, pag. 128. — 2581
Maculis amphigenis, irregularibus, saepe angustis et elongatis, plerumque paucis in quoque folio, 2-4 mm. v. etiam 1-1,5 cm. lat., interdum folii dimidium occupantibus; peritheciis minutis, utrinque perspicuis, sublenticularibus, subastomis, membranaceis, nigris, 75-100 µ. diam.; ascis oblongis, sessilibus, 40-60 ≈ 12, a paraphysatis; sporidiis conferto-distichis, oblongo-ellipsoideis, subinæquilateralibus, 1-septatis et constrictis, flavidо-hyalinis, 12-15 ≈ 5.
Hab. in foliis vivis Staphyleæ trifoliæ, socio *Macrosporio* in Manhattan Kansas (W. T. SWINGLE). — Adsunt quoque pycnidia stylosporis ellipsoideis, subfuscis, continuis, circ. 5 ≈ 2,5.
108. **Sphærella collina** Secc. et Sp. Syll. I, p. 493 — var. **caulicola** 2582
Berl. et F. Sacc. Contr. Myc. Lus. p. 4. — Peritheciis subgregariis, erumpentibus; ascis amplis, sursum tenuioribus, 32-36 ≈ 12-14; sporidiis fusoideis, 12-15 ≈ 3-4, ad medium 1-septatis, vix constrictis, loculo superiore vix crassiore.
Hab. in sarmentis Lonicerae etruscae in Lusitania.
109. **Sphærella Implexæ** Passer. Diagn. Funghi nuovi I, n. 22. — 2583
Maculis amphigenis, griseis, submarginalibus, fusco-limitatis; peritheciis amphigenis, sparsis, ostiolo nigro vix emerso; ascis oblongo-clavatis, basi attenuato-stipitatis, 60 ≈ 12; sporidiis distichis, oblongis, medio septatis, non v. vix constrictis; loculis guttulatis, altero majore, hyalinis, 17-20 ≈ 5-7,5.
Hab. in foliis languidis Lonicerae implexæ in Horto botanico, Parma.
110. **Sphærella ramulorum** Passer. Diagn. Fung. nuovi I, n. 21. — 2584
Peritheciis crebre sparsis, minutis, ostiolo conico epidermidem perforante; ascis ovatis vel ovato-oblongis, utrinque attenuatis vel etiam clavatis, 8-sporis, 40-80 ≈ 12; sporidiis bi-tristichis, oblongo-fusiformibus, medio septatis, non constrictis, hyalinis, 12-15 ≈ 3-5.

Hab. in ramulis vivis Lonicerae Caprifolii, Vigheffio pr. Parma.

111. **Sphærella Symphoricarpi** Passer. Diagn. F. N., IV, p. 3, n. 4. — 2585

Peritheciis sparsis, minutis, epidermide nigrefacta velatis, punctiformibus, membranaceis, cellulis amplis fuscidulis contextis; ascis ovato-elongatis, basi abrupte breviter stipitatis, 8-sporis, 45×15 ; sporidiis ad ascorum basim conglobatis, superne subdistichis, oblongo-spathulatis, non vel vix constrictis, hyalinis, $17,5-20 \times 6-7$.

Hab. in ramulis vivis Symphoricarpi racemosi, Parma.

112. **Sphærella Periplocæ** Passer. Diagn. F. N., IV, p. 3, n. 5. — Pe- 2586

ritheciis subgregariis, tectis, per corticem fissum erumpentibus, globosis, atris, ostiolo obtuso; ascis obclavatis, apophysatis, basi breviter stipitatis, rectis vel curvis, $62-87 \times 15-17,5$; sporidiis subdistichis, ovato-elongatis, utrinque rotundatis, prope medium septatis, non vel vix constrictis, hyalinis, $16-20 \times 5-6$, extreto altero longiore et aliquantulum angustiore.

Hab. in ramulis siccis Periplocæ in Horto botanico, Parma.

113. **Sphærella Patouillardii** Sacc. in Add. Syll. p. 407. — Maculis 2587

submarginalibus, candidantibus, nitidis, amphigenis; peritheciis hypophyllis, epidermide tectis, gregariis, punctiformibus, globoso-depressis, minutis, $150-200 \mu$. diam., poro pertusis, contextu indistincto, olivaceo-fusco; ascis dense fasciculatis, tereti-clavatis, breve stipitatis, non vel obsolete paraphysatis, $70-80 \times 12$, octosporis; sporidiis distichis, fusoideis, leniter curvis, utrinque obtusiusculis, 1-septatis (vel saltem plasmate bipartito), $20-25 \times 3-4$, hyalinis.

Hab. in foliis Buxi sempervirentis, Eaux Bonnes in Pyrenæis (PA TOUILLARD). — ***S. Briardi** Sacc. Rev. Myc. 1890 Oct.: Maculis flavo-citrinis, brunneo-marginatis; ascis teretibus, sessilibus, $54-60 \times 10-12$; sporidiis 1-septatis, $26-28 \times 4$. In foliis Buxi sempervirentis, Melun Galliæ (Doct. ROUSSEL).

114. **Sphærella ambiens** Starb. Skand. Pyr. p. 17, t. I, f. 4 (*Myco-* 2588

sphærella). — Peritheciis epiphyllis, interdum hypophyllis, sparsis vel gregariis, maculas irregulares, cinerescentes formantibus, basi semper epidermide innatis, $100-125 \mu$. diam.; ascis cylindraceis, sessilibus, $35-40 \times 6$; sporidiis octonis, ellipsoideis v. ellipsoideo-fusoideis, distichis, hyalinis v. dilutissime lutescentibus, medio inæqualiter septatis, $8-12 \times 2,5-3$.

Hab. in foliis Berberidis vulgaris pr. Rodga in parœcia Simonstorp Ostrogothia Scandinaviæ.

115. **Sphærella australis** Spieg. F. Fueg. n. 195. — Maculis nullis v. 2589 amphigenis, parum manifestis, indeterminatis, rufescentibus v. fuscescenscentibus; peritheciis hypophyllis, hinc inde densiuscule multigregariis, immersis, epidermide velatis e conoideo lenticularibus, 75-80 μ . diam., vix prominulis, atris, glabris, tenui-membranaceis, contextu parenchymatico, olivaceo, ostiolo impresso; ascis cylindraceo-obclavatis, antice truncato-rotundatis, crassiuscule tunicatis, postice abrupte coarctatis ac pedicello brevissimo noduloso suffultis, 20-25 = 5-6, octosporis, apophysatis; sporidiis di-tristichis, e cylindraceo subclavulatis, sursum obtusiusculis, deorsum acutiuscule rotundatis, 9-10 = 2, medio 1-septatis, non v. vix constrictulis, hyalinis.

Hab. ad folia dejecta Berberidis ilicifoliæ in dumetis in Staten island Fuegiæ.

116. **Sphærella Puiggarii** Spieg. F. Puigg. n. 276. — Maculis epiphyllis, minutis, candidis, determinatis, 1-2 mm. diam.; peritheciis lenticularibus minutis, atris, 100-120 μ . diam.; ascis obclavulatis, 50 = 10-15; sporidiis elliptico-ovatis, hyalinis, 16-18 = 4-5, medio 1-septatis vix subconstrictis (an demum 3-septatis?).

Hab. ad folia viva Bouddleiæ speciei cujusdam in dumetis prope Apiah Brasiliæ.

117. **Sphærella conferta** Spieg. Fung. Guar. Pug. I, n. 216. — Maculis amphigenis, parvulis, 1-2 mm. diam., arecentibus, sordide fusco-griseis, obscure determinatis, irregulariter discoideis, angulosis vel repandulis, densiuscule sparsis; peritheciis hypophyllis, rarissime epiphyllis, numerosissimis, densissime constipatis, maculas nigras efficientibus, epidermide primo tectis, dein suberumpentibus, atris, glabris, globoso-lenticularibus, 70-80 μ . diam., minute papillulato-ostiolatis, membranaceis, contextu parenchymatico, olivaceo; ascis cylindraceo-obclavatis vel subovatis, antice subtruncato-rotundatis, postice abrupte, nodulose brevissimeque stipitatis, 30-35 = 8-10, apophysatis, 8-sporis; sporidiis distichis, elliptico-elongatis, rectis vel vix inaequilateralibus, utrinque rotundatis, 10-13 = 2.5-3, medio 1-septatis, non constrictis, loculis in senectute (an semper?) obscure diblastis, 1-2-guttulatis, supero saepè subcrassiore, hyalinis.

Hab. ad folia languida vel dejecta Sapindaceæ cujusdam in silvis prope Guarapí Brasiliæ. — Species habitu mox distinguenda, facillime ad genus *Sphaerulinam* transferenda.

118. **Sphærella exarida** Winter in Rev. Myc. 1885. — Peritheciis in 2592

macula, plerumque magna, irregulari, determinata, fuscescente-grisea, arida, elevata, fusco-nigro-limitata gregariis, immersis, globosis, ostiolo papillato, pertuso, prominulo, membranaceo, atro, 90-115 μ . diam.; ascis oblongis vel e basi latiori sursum parum attenuatis, sessilibus, 8-sporis, 42-52 \approx 12-14; sporidiis conglobatis, anguste clavatis, utrinque rotundatis, interdum parum curvatis, medio septatis, non constrictis, hyalinis, 17-21 \approx 3.5-4.

Hab. ad folia viva Sapindaceæ cujusdam, Guarapi Paraguay Amer. austr.

119. **Sphærella Sapindi** Ell. et Ev. in Journ. of Myc. 1886, p. 101. 2593

— Peritheciis epiphyllis, globosis, prominulis, $\frac{1}{3}$ mm. diam., sparsis, maculis rotundis, definite marginatis, albis, 3-4 mm. diam., insidentibus; ascis oblongo-cylindraceis, 60 \approx 12; sporidiis distichis, subellipsoideis utrinque acutiusculis, 1-septatis, ad septum vix constrictis, 16 \approx 4, hyalinis.

Hab. in foliis vivis Sapindi marginati, Missouri Amer. bor. (B. T. GALLOWAY).

120. **Sphærella? Balansæ** Speg. F. Guar. II, n. 86. — Maculis nul-

lis; peritheciis hypophyllis, hinc inde densissime confertis, plagulas atras subdothideaceas, 1-2 mm. lat., angulosas granulosas fusco-atras efficientibus, globoso-lenticularibus, 80-90 μ . diam., prominulis, saepe confluentibus (dothideaceis?) membranaceo-coriacellis, atris glabris, contextu subpellucido parenchymatico olivaceo donatis, ostiolo latissimo 20 μ . diam., pertusis; ascis cylindraceo-obclavatis vel subfusoideis, apice subtruncato-rotundatis, deorsum cuneatis stipiteque brevissimo crassiusculo suffultis, 40 \approx 12-14, octosporis, aparaphysatis; sporidiis 2-3-stichis, cylindraceo-subclavulatis, superne obtusiuscule rotundatis, 20 \approx 4-5, medio 1-septatis non v. vix constrictis, rectis v. subcurvulis, hyalinis.

Hab. ad folia viva Sapindaceæ cujusdam in silvis pr. Guarapi Brasiliæ. — Species priori peraffinis sed satis ac facile distincta, notis nonnullis ad Dothideaceas vergens.

121. **Sphærella Umbellulariæ** Cooke et Hark. in Grev. XIII, p. 21. — 2595

Hypophylla; maculis suborbicularibus confluentibusve et irregularris, purpureo-brunneis, margine nigro limitatis; peritheciis hinc illinc aggregatis, semiimmersis, atris; ascis clavatis; sporidiis biserialibus, arcte ellipticis, uniseptatis, hyalinis, 15 \approx 4.

Hab. in foliis emortuis Umbellulariæ in California.

122. **Sphærella tahitensis** Sacc. Misc. Myc. I, p. 27. — Peritheciis 2596

hypophyllis, velatis, hinc inde dense gregariis, punctiformibus, len-

ticularibus, pertusis, 80 μ . diam.; ascis tereti-fusoideis, subsessilibus, apice rotundatis, 48 \approx 7-8, octosporis; sporidiis distichis fusoideo-clavulatis, 1-septatis, leniter constrictis, 4-guttulatis, 14 \approx 3, hyalinis.

Hab. in foliis emortuis *Mangiferæ* indicæ (sociâ *Pestalozziæ funereâ* f. *Mangiferæ*: conidiis quam in typo paullo minoribus, 22 \approx 6) in insula oceanica Tahiti (G. BRUNAUD).

123. **Sphærella Sassafras** Ell. et Ev. in Torr. bot. Club 1883, p. 98. 2597

— Peritheciis minutis, semiimmersis, in superficie inferiore foliorum sparsis vel subaggregatis, sed non in quaque macula decolorata adstantibus; ascis 35 \approx 5; sporidiis distichis, oblongo-ellipsoideis, uniseptatis, 4 \approx 1,7, hyalinis.

Hab. in foliis emortuis *Sassafras officinalis*, Fairfield, Ohio Amer. bor. (W. A. KELLERMANN).

124. **Sphærella rubiginosa** Cooke in Grevillea XIII, pag. 90. — Epi-

phylla; peritheciis sparsis, minutis, punctiformibus, sub prominulis, atris; ascis clavatis, sessilibus; sporidiis ellipticis, obtusis, medio leniter constrictis, uniseptatis, chlorinis, 10 \approx 5.

Hab. in foliis emortuis *Pittospori rubiginosi*, Johnston River, Queensland Australiæ. — A *Sph. Pittospori* omnino diversa, tametsi fructificatio similis.

125. **Sphærella Terebinthi** Pass. Diagn. F. nuov. III, n. 8. — Peri-

theciis hypophyllis minutis gregariis, tectis, maculas fuscas venis limitatas formantibus; ascis numerosis, forma variis, basi breviter stipitatis, raro clavatis, rectis, saepius curvis, prope basim vel medio ventricosis, 55-75 \approx 15-20; sporidiis octonis subdistichis v. conglobatis, oblongis, didymis vix constrictis, loculo inferiore attenuato longiore, hyalinis, 20-22,5 \approx 5.

Hab. in foliis dejectis *Pistaciæ Terebinthi*, Parma in Horto botanico. — Ascis numquam clavato-cylindricis et sporidiis non ellipticis nec loculis æqualibus, a *Sph. Pistaciæ* Cooke diversa.

126. **Sphærella Pistaciæ** Cooke Journ. of Bot. 1883, p. 109. — Hy-

pophylla; peritheciis semiimmersis, atris, paucis (6-10) in cæspitulos aggregatis; ascis clavato-cylindricis; sporidiis subellipticis, uniseptatis; loculis æqualibus, hyalinis.

Hab. in foliis *Pistaciæ veræ*, Marseille Galliæ (Roux).

127. **Sphærella Gordoniæ** Cooke Journ. of Bot. 1883, p. 109. — Hy-

pophylla; peritheciis sparsis tectis, vix visilibus; ascis subclavatis; sporidiis inordinatis, ellipticis, uniseptatis, hyalinis, medio vix constrictis, 10 \approx 4.

Hab. in foliis *Gordonia* & *Lasianthæ*, Darien Georgia Amer. bor. (RAVENEL).

128. **Sphærella Panacis** Cooke Journ. of Bot. 1883, p. 138. — Hy-
pophylla, gregaria, maculæformis; peritheciis minimis, subglobosis,
atris, semiinnatis, maculas orbiculares vel irregulares formantibus;
ascis cylindrico-clavatis, sessilibus; sporidiis arcte ellipticis, hya-
linis, uniseptatis, $10 \approx 2.5$.

Hab. in foliis *Panacis crassifolii*, Nova Zelandia, South Island (KIRK).

129. **Sphærella papyrifera** Pass. Diagn. Fung. nuov. I, n. 20. — Pe-
ritheciis gregariis in maculis fuscis oblongis, subglobosis, atris;
ascis magnitudine varia, clavatis vel cylindricis; sporidiis oblongis,
subspathulatis, obscure tenuiter septatis, hyalinis, endoplasmate gra-
nuloso, interdum uni-biguttulatis, $20 \approx 7-7.5$.

Hab. in petiolis *Araliæ papyriferae* gelu tactis, in Horto
botanico, Parma. — Differt a *Sphærella Firmiana* Pass. peri-
theциis in maculas aggregatis et ascorum forma; et a *Sphærella*
Araliæ C. et Harkn. saltem sporidiorum forma et magnitudine.

130. **Sphærella Drymidis** (Berk.) *Sphæria Drymidis* Curr. Linn. 2604
Trans. XXII, p. 333, Journ. of Bot. 1883, p. 109. — Epiphylla;
maculis determinatis, albis, orbicularibus, depresso-papillatis, depresso-
punctatis, atris, semiinnatis; ascis cylindraceis; sporidiis ellipticis, uni-
septatis, loculo inferiore tenuiore, hyalinis, $12 \approx 4$.

Hab. in foliis *Drymidis*, Juan Fernandez.

131. **Sphærella Nigredo** (Schw.) Cooke Journ. of Bot. 1883 pag. 109, 2605
Sphæria Schw. Am. n. 1799. — Hypophylla, vix innata, aggrega-
ta; peritheciis accumulatis majusculis pro ratione, atris, rugosis, pa-
pillatis, passim solitariis, interdum aggregatis et tunc crusta pul-
veraceo-cinerescente, orta ex parenchymate, aspersis; ostioliis pa-
pillæformibus subapertis; ascis clavatis; sporidiis sublanceolatis,
uniseptatis, loculis subconicis, hyalinis, $8-10 \approx 3$.

Hab. in foliis *Rhois glabræ* Amer. bor. — Cfr. Syll. n. 4474.

132. **Sphærella Ilicis** Ellis in Amer. Nat. XVII, 1883, pag. 317. — 2606
Amphigena; maculis rotundatis, superne albis, inferne brunneis,
margin'e elevato purpureo circumdati; peritheciis punctiformibus,
subglobosis, semiinnatis, prominulis, ostiolo ampio; ascis oblongo-
cylindricis, $40-55 \approx 7.5-11$; sporidiis biserialibus, clavato-oblongis,
subhyalinis, uniseptatis, leviter constrictis, $13-15 \approx 3$.

Hab. in foliis subvivis *Ilicis glabræ*, New Jersey Amer. bor.
(ELLIS).

133. **Sphærella Prini** Cooke in Rav. Amer. Fungi n. 753, Journ. of Bot. 1883, p. 106. — Epiphylla, sparsa vel totam paginam occupans; peritheciis semiinnatis, prominulis, atris; ascis clavatis, sessilibus; sporidiis minutissimis, arcte ellipticis, utrinque obtusis, uni-septatis, hyalinis, $5 \approx 1,8$.

Hab. in foliis Prini glabri, S. Carolina Amer. bor. (RAVENEL).
— Perithecia 0,1-0,14 mm. diam.

134. **Sphærella cornifolia** (Schw.) Cooke in Rav. Amer. Fungi n. 681, 2608
Journ. Bot. 1883, p. 108, *Sphæria Corni* Schw. n. 1792. — Hypophylla, maculas efformans, maximas orbiculatas ambitu indeterminatas, *Sph. effiguratæ* instar; cæspitibus trium aut quatuor peritheciorum punctiformium absque ulla crista, inter se distantibus sed frequentibus in eadem macula; peritheciis ipsis astomis, innatis, evacuatis, extus punctatis; ascis clavatis, abbreviatis; sporidiis ellipticis, biseriatis, hyalinis, uniseptatis, $5 \approx 2$.

Hab. in foliis Corni floridæ in Carolina inferiore Americæ borealis.

135. **Sphærella carpogena** Pass. Diagn. F. nuovi I, n. 9. — Peritheciis sparsis, subcuneatis, minutis, ostiolo atro, acuto vix perspicuo; ascis oblique ovoideo-oblongis, $50-75 \approx 15-20$; sporidiis tristichis vel conglobatis, oblongo-ovatis, medio septatis, non constrictis, hyalinis, $15-17,5 \approx 5$.

Hab. in fructibus Liriodendri Tulipiferæ in Horto bot., Parma. — *Sphærellæ Tassianæ* fructificatione similis sed peritheciorum forma et situ distinguenda.

136. **Sphærella Liriodendri** Cooke Journ. of Bot. 1883, pag. 108. — 2610
Epiphylla; maculis orbicularibus, brunneis 1 cm. diam.; peritheciis subinnatis punctiformibus, atris; ascis clavato-cylindricis; sporidiis ellipticis, uniseptatis, hyalinis, $16 \approx 5$. — *Spermogonium, Phyllosticta Liriodendri* Cooke.

Hab. in foliis Liriodendri Tulipiferæ, Darien Georgia.

137. **Sphærella Pernettiæ** Speg. F. Fueg. n. 201. — Maculis nullis; peritheciis hypophyllis, sparsis, e globoso lenticularibus 100-120 μ . diam., epidermide velata, prominulis atris, glabris, coriaceo-membranaceis, contextu indistincto fuligineo, opaco, ostiolo pusillo perforatis; ascis obclavatis, antice truncato-rotundatis, non vel vix 1-foveolatis postice attenuato-coaretatis, stipite brevissimo crasso noduloso suffultis, $40-45 \approx 10-12$, octosporis, apophysatis; sporidiis di-polystichis e cylindraceo fusoides, utrinque obtusiusculis, $15-17 \approx 3$, saepius curvulis, medio 1-septatis non constrictis, hyalinis.

Hab. ad folia emortua arida dejecta vel adhuc pendula Pernettiae mucronatae in Staten island Fuegiæ.

138. **Sphærella Polifoliæ** Ell. et Ev. Proc. Phil. 1890, p. 231. — Pe-
ritheciis epiphyllis, maculis griseo-nigris, indefinitis, 2 mm. vel
ultra diam., insidentibus, erumpentibus, asperis, minutis, supra late
pertusis; ascis oblongis, 35-40 \approx 6-8, aparaphysatis; sporidiis di-
stichis, clavato-oblongis, 1-septatis, 10-12 \approx 2,5-3.

Hab. in foliis vivis v. partim emortuis Andromedæ polifo-
liæ, London Canada Amer. bor. (J. DEARNNESS).

139. **Sphærella apiahyna** Speg. F. Puigg. n. 275. — Maculis arescen-
ti-impressis, determinatis, rufescentibus, 1-3 mm. diam., hypophyllis;
peritheciis sparsis globulosis immersis, 90-120 μ . diam.; ascis ob-
clavatis 40 \approx 10; sporidiis cylindraceo-clavulatis hyalinis, 9-10 \approx 3,
uniseptatis.

Hab. ad folia languida coriacea Ericaceæ cujusdam in dumetis prope Apiahyn Brasiliæ.

140. **Sphærella hæmatodes** B. et C. in Herb. Berk. Cooke in Journ. of Bot. 1883, p. 70. — Epiphylla; maculis orbicularibus, sparsis vel confluentibus, rubro-fuscis, late marginatis; peritheciis mini-
mis, nigris, semiimmersis, punctiformibus; ascis cylindrico-clava-
tis; sporidiis arcte ellipticis, continuis, hyalinis, 8-10 \approx 2,5.

Hab. in Kalmia glauca in Amer. bor. — *Sph.* coloratæ
valde affinis.

141. **Sphærella arbucola** Peck in Torr. bot. Club. 1883, p. 75. — 2615
Maculicola; maculis suborbicularibus, inæqualibus, nigrescentibus vel cinereis, brunneis, paginam inferiorem foliorum occupantibus; peritheciis minutis, nigris, epiphyllis, sparsis vel in glomerulos minutos congestis, epidermidem perforantibus; ascis oblongis, sæpe sursum acutis, 40-60 μ . longis; sporidiis stipatis, acutis, unisepta-
tis, 11-15 \approx 4, hyalinis.

Hab. in foliis emortuis Arbuti Menziesii, Santa Cruz in California (C. G. PRINGLE).

142. **Sphærella Gallæ** Ell. et Ev. in Torr. bot. Club. 1884, p. 75. — 2616
Peritheciis minutis, sparsis vel 2-3 gregariis, epidermidem rum-
pentibus; ascis 40-45 \approx 10, sessilibus; sporidiis stipatis, bi-tristi-
chis, subtilibus, clavatis, 1-septatis, 12-15 \approx 3.

Hab. in gallis Vaccinii corymbosi, Newfield Amer. bor.

143. **Sphærella Rhododendri** Cooke Journ. of Bot. 1883, p. 108. — 2617
Epiphylla; peritheciis semiinnatis, nitidis, atris, demum asperato-
prominulis, in maculas irregulares congestis; ascis cylindricis; spo-

- 642 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Sphærella.
ridiis arcte ellipticis, uniseptatis, rectis, utrinque obtusis, hyalinis,
 $10-12 \approx 2,5$.
Hab. in foliis languidis et emortuis Rhododendri, Forden
Britanniæ.
144. **Sphærella Gardeniæ** Cooke Journ of Bot. 1883, p. 108. — Hy- 2618
pophylla; peritheciis sparsis, punctiformibus, semiinnatis, atris;
ascis clavatis; sporidiis inordinatis, elongato-ellipticis unisepta-
tis, hyalinis, $12 \approx 3,5$. — Spermogonium: *Phyllosticta Gardeniæ*
Cooke.
- Hab.* in foliis Gardeniæ floridæ, Carolina infer. Amer. bor.
145. **Sphærella Weinmanniæ** Cooke in Grevillea XIV, pag. 130. — 2619
Epiphylla; peritheciis gregariis, subglobosis, atris, semiimmersis,
punctiformibus, pertusis; ascis clavatis; sporidiis lanceolatis, uni-
septatis, $15-18 \approx 4$, hyalino-luteolis.
Hab. in foliis emortuis Weinmanniæ racemosæ, New Zea-
land Austral. (COLENSO).
146. **Sphærella cinerescens** Fuck. — Sacc. Syll. I, p. 493, Winter
in Hedwigia 1886, p. 21. — Perithecia maculas cinerescentes, plus
minusve effuse formantia.
147. **Sphærella cerasina** Cooke. — Sacc. Syll. I, p. 493 — var. **padina** 2620
Karst. in Hedw. 1884, n. 1, pag. 2, Revisio atque Synop. Ascom.
fenn. p. 64. — Peritheciis hypophyllis, innato-prominulis, puncti-
formibus, sphæroideis, in maculas inæquales, subinde orbiculares,
plus minus dense aggregatis, astomis; ascis cylindraceo-clavulatis,
 $28 \approx 6$; sporidiis ovoideo-oblongatis, uniseptatis, $8 \approx 3$.
Hab. in foliis Pruni Padi in regione Aboensi.
148. **Sphærella Lenticula** Cooke Rav. Amer. Fungi n. 800. — Hy- 2621
pophylla; peritheciis globosis, atris in pustulas elevatas, lenticula-
res dense stipatis, circiter 1 mm. diam.; ascis breviter clavatis; spo-
ridiis inordinatis, ellipticis, uniseptatis $8-9 \approx 3-3,5$.
Hab. in foliis Cerasi carolinianæ, Carolina infer. Ameri-
cæ borealis.
149. **Sphærella crenata** Brun. Miscell. mycol. p. 11. — Maculis in 2622
folii pagina superiore nigrescentibus, in inferiore brunneo-griseis,
magnitudine et forma variis, irregularibus; peritheciis hypophyl-
lis, sparsis, punctiformibus, erumpentibus, brunneis; ascis cylindra-
ceis v. subclavatis, breve pedicellatis, $85-90 \approx 8$; sporidiis octonis,
monostichis v. oblique monostichis, ovoideo-subfusoideis, 1-septa-
tis, ad septum non v. vix constrictis, $10-14 \approx 7,5-8$, hyalinis.
Hab. in foliis Spirææ crenatæ, Saintes Galliæ.

150. **Sphærella fruticum** Starb. Skand. Pyr. p. 17, f. 5. — Peritheciis punctiformibus, laxe gregariis, interdum 2-3-connatis, sphæroideis, 80-100 μ . diam., epidermide vix tectis, hyphis mycelii plus minus crassis insidentibus; ascis ovato-oblongis, sessilibus v. paulatim pedicellatis, 25-30 \approx 10-12; sporidiis octonis, ovoideo-fusoides v. ovoideis, primo guttulis minusculis præditis, deinde prope medium 1-septatis paullumque constrictis, superne distichis, inferne tristichis, 9-10 \approx 3.

Hab. ad caules Rubi idæi pr. Rogda in parœcia Simonstorp Ostrogothia (E. HAGLUND) et ad Are Frösnäset in Jemtlandia Scandinaviæ (A. J. GREVILLIUS).

151. **Sphærella rosigena** Ell. et Ev. Journ. Myc. 1887, pag. 45. — 2624 Maculicola; maculis amphigenis, rufo-brunneis, purpureo-marginalis, definitis, 3-4 mm. diam.; peritheciis epiphyllis, dense sparsis, minutis, 60-75 μ . diam., partim erumpentibus, subastomis, nigris; ascis subclavato-oblongis, 25-30 \approx 8-10; sporidiis distichis, clavato-oblongis, hyalinis, 1-septatis, 10-12 \approx 2, subacutis.

Hab. in foliis vivis Rosarum cultarum, Louisiana Amer. bor. (LANGLOIS). — A *Læstadia Rosæ* distinguenda.

152. **Sphærella cerasicola** Passer. Diagn. Funghi nuovi I, n. 17. — 2625 Peritheciis minutissimis, sparsis, epidermide tectis eamque punctiformiter sublevantibus, globulosis, atris, ostiolo simplici, pertuso, membranaceis, eximie parenchymaticis; ascis oblongo-piriformibus vel cuneatis, 8-sporis, 30-36 \approx 12-15; sporidiis congregatis v. subtristichis, obovato-oblongis, prope medium septatis, loculo altero breviore et angustiore, hyalinis, 10-12 \approx 5.

Hab. in ramulis Pruni cerasi, gummosi infectis, socio *Cyniothyrio Cerasi*, Vigheffio pr. Parma.

153. **Sphærella rhodophila** Passer. Diagn. F. nuovi, I, n. 18. — Hy- 2626 pophylla subgregaria; peritheciis minutis prominulis, maculas griseas, venis limitatas, formantibus; ascis brevibus, saepius basi vel medio ventricosis, interdum globoso-ovatis, 6-8-sporis; sporidiis oblongis, didymis, hyalinis, medio leniter constrictis, 20-23 \approx 8.

Hab. in foliis dejectis Rosæ in Horto botanico, sociâ *Disco-sia Artocreate*, Parma.

154. **Sphærella Aristoteliæ** Cooke in Grevillea XV, pag. 16. — Ma- 2627 culis epiphyllis, pallidis, orbicularibus, late fusco-cinctis; peritheciis minutis, immersis, membranaceis, globoso-depressis, perforatis; ascis clavatis, octosporis; sporidiis cylindraceo-ellipsoideis, utrinque rotundatis, uniseptatis, vix constrictis, 18-20 \approx 4.

Hab. in foliis vivis Aristoteliæ racemosæ, New Zealand Australasiæ (COLENZO).

155. **Sphærella fraxinea** Peck in 35 Rep. St. Mus. 1884, pag. 145. 2628

— Peritheciis numerosis, minutis 75 μ . diam., nigris, epiphyllis v. hypophyllis, saepe in maculis suborbicularibus coacervatis; ascis oblongis, saepe sursum leniter attenuatis, 35-40 μ . long.; sporidiis stipatis, oblongo-ovoideis, uniseptatis, 10-12 \approx 4.5, inæqualibus, parte majore tripla vel quadrupla.

Hab. in foliis Fraxini americanæ in montibus Helderberg Amer. bor.

156. **Sphærella fraxinicola** (Schw.) Cooke Journ. of Bot. 1883, p. 107, 2629

Sph. fraxinicola Schw. Am. n. 1787. — Hypophylla; peritheciis subinnatis, nigris, demum fisis orificio longitudinaliter, difformibus, paucis tantum conjunctis maculam atram efficientibus minorem, maculis quasi confluentibus inter se; ascis clavatis, abbreviatis, sporidiis inordinatis, subellipticis, uniseptatis, hyalinis, loculo inferiore tenuiore, 7.5 \approx 3.

Hab. in foliis Fraxini americanæ, Darien Georgia, Amer. bor. Cfr. Syll. II, n. 4478.

157. **Sphærella effigurata** (Schw.) Cooke Journ. of Bot. 1883, p. 107, 2630

Sphæria Schw. Amer. n. 1790. — Maculis longe lateque effusis, nigro-cinerescentibus ambitu determinatim effigurato, et ob frequentiam peritheciorum in margine quasi nigrocincto. saepe totum folium in pagina aversa occupans; peritheciis innumeris, minutis, accumulatis in his maculis, astomis, subinnatis, convexulis, nigris valde invicem approximatis et crusta cinerescenti quasi inter se connexis; ascis clavatis; sporidiis ellipticis, uniseptatis, vix constrictis, hyalinis, 15 \approx 4.

Hab. in foliis Fraxini acuminatæ, Amer. bor. Cfr. Syll. n. 4479.

158. **Sphærella pterophila** Passer. Diagn. nuovi Funghi I, n. 24. — 2631

Peritheciis minutis, aggregatis, atris, matricem plus minus infuscantibus; ascis oblongo-clavatis v. etiam brevibus, ovatis aut gibbis, 45-75 \approx 15, 8-sporis; sporidiis distichis v. congregatis, cuneato-oblongis, extremo altero acutiusculo, altero rotundato, medio septatis, non constrictis, hyalinis, 17-5 \approx 5.

Hab. in samaris Fraxini Orni in Horto botanico, Parma.

159. **Sphærella oleina** Cooke et Rav. Amer. Fungi n. 754, Journ. of 2632

Bot. 1833, p. 107. — Epiphylla; maculis albidis, suborbicularibus, demum confluentibus, rubro-marginatis; peritheciis mininis, puncti-

formibus, vix prominulis, atris, hinc illic subcircinatis; ascis clavato-cylindricis; sporidiis ellipticis, uniseptatis, hyalinis, leniter constrictis, $12 \approx 4$. — Spermogonium: *Phyllosticta oleina* Cooke.

Hab. in foliis Oleæ americanæ, Aiken Carolina Amer. bor.

160. **Sphærella Chionanthi** (B. et Br.) Cooke Bot. Journ. 1883, p. 107,

Didymella Chionanthi Syllog. I, n. 2143. — Adde: ascis cylindrico-clavatis; sporidiis fusiformibus, 1-septatis, hyalinis, $18 \approx 5$.

— In foliis *Chionanthi zeylanicæ*, Ceylon (nec in ramis).

161. **Sphærella ulmifolia** Pass. in Erb. critt. ital. Ser. II, n. 1281. — 2633

Maculis candidis, sparsis, parvulis, discoideis; peritheciis epiphyllis, paucis, punctiformibus, atris; ascis e basi ventricosa attenuatis, breviter abrupte stipitatis, 8-sporis, $50-70 \approx 18-20$; sporidiis subdistichis, oblongo-cuneatis, ad septum non vel vix constrictis, $20-22 \approx 7-8$.

Hab. in foliis vivis Ulmi campestris, Parma (G. PASSERINI).

— Facie *Hendersoniam ulmifoliam* Pass. in Thum. Mycoth. univers. n. 1978, refert, quæ formam ejusdem stylosporam sistit.

162. **Sphærella platanifolia** Cooke Journ. of Bot. 1883, pag. 106. — 2634

Hypophylla, sparsa; peritheciis exiguis, atris, semiimmersis, punctiformibus; ascis clavatis, sessilibus; sporidiis biseriatis, subellipticis, uniseptatis, hyalinis, loculo inferiore tenuiore, $8 \approx 4$.

Hab. in foliis Platani occidentalis, Georgia Amer. bor. (RAVENEL).

163. **Sphærella incanescens** (Schw.) Cooke Journ. of Bot. 1883, p. 107, 2635

Sphæria Schw. Amer. n. 1796. — Maculis incanescensibus, late effusis, indeterminatis, quasi pruinatis, insident perithecia punctiformia subglobosa, minutissima, nigra, demum evacuata, sæpe quasi truncata aut collapsa; ascis cylindricis; sporidiis ellipticis, uniseptatis, hyalinis, $8 \approx 3$.

Hab. in foliis Tiliæ in Amer. borealis (SCHWEINITZ). — Cfr. Syll. n. 4477.

164. **Sphærella platyphæca** Karst. in Hedw. 1883, n. 12, p. 179 et Rev. 2636

atque Synop. Ascom. fenn. p. 66. — Peritheciis.....; ascis sessilibus v. brevissime stipitatis, inferne inæqualiter saccato-inflatis (ut in *Sph. Tassiana*), $65-80 \approx 20-28$; sporidiis octonis, ovoideo-oblongatis, medio v. paullo extra medium uniseptatis, ad septum vix v. leviter constrictis, vulgo guttulatis, $22-35 \approx 7-10$.

Hab. in foliis emortuis Tiliæ platyphylla in Mustiala. —

Forte cum *Sphærella Tassiana* conjungenda.

165. **Sphærella bracteophila** Passer. Diagn. Fung. nuovi, I, n. 14. — 2637

Peritheciis epiphyllis, sparsis, subglobosis, atris; ascis oblongis, clavatis v. medio turgidulis, 8-sporis; sporidiis oblongis, apicibus rotundatis, altero angustiore, medio septatis, subconstrictis, 18-23,5 \approx 5-6,5.

Hab. in bracteis dejectis *Tiliæ europææ*, Parma.

166. **Sphærella maculans** Passer. Diagn. F. nuov. III, n. 10. — Pe. 2638
ritheciis in maculis epiphyllis albicantibus, fusco-marginatis, paucis,
crassiusculis, atris; ascis sæpius ovoideo-elongatis v. subclavatis,
8-sporis, 50-75 \approx 12-15; sporidiis oblongis, bi-tristichis, medio se-
ptatis, hyalinis, 15 \approx 5.

Hab. in foliis vivis *Populi albæ*, Vigheffio, Parma. — Asco-
rum et sporidiorum magnitudine a *S. maculosa* Sacc. et a *S. ma-
culari* (Fr.) pariter differt. Maculis exceptis, ad *S. crassam* Auersw.
propius accedit.

167. **Sphærella populifolia** Cooke Rav. Amer. Fungi n. 689, Journ. 2639
of Bot. 1883, p. 197. — Hypophylla; peritheciis innato-prominu-
lis, punctiformibus, globosis, nigris, in maculas minutas crebras
dense aggregatis v. sparsis; ascis cylindricis; sporidiis sublanceo-
latis; uniseptatis hyalinis, 16-18 \approx 3,5-4.

Hab. in foliis *Populi angulatæ*, Aiken Carolina austr.
Amer. bor.

168. **Sphærella salicicola** (Fr.) Fuck. — Sacc. Syll. I, p. 487. — Var.
borealis Karst. in Hedw. 1884, p. 3. — Ascis 50-65 \approx 10-12; spo-
ridiis 9-12 \approx 2-4. — Ad folia emortua *Salicis phyllicifoliæ*, *S. ca-
preæ* et *S. auritæ* in Fennia, vulgatissima.

169. **Sphærella Æsculi** Cocc. et Mor. Funghi Bol. IV, pag. 9, t. II, 2640
f. 2-5. — Peritheciis numerosis, minutis, erumpentibus, epiphyllis,
in areolis magnis subdecoloratis, oblongis, irregularibus gregariis,
globulosis, subdepressis, 90-98 μ . diam.; ascis cylindraceis, 42-46
 \approx 8-10; sporidiis hyalinis, distichis, medio subconstrictis, 1-septa-
tis, ovoideo-oblongis, utrinque obtusiusculis, 12-14 \approx 4-5.

Hab. in foliis *Quercus Æsculi*, Montese in agro Bononiensi
Italiæ.

170. **Sphærella aquatica** Cooke Rav. Amer. Fungi n. 690, Journ. of 2641
Bot. 1883, p. 106. — Hypophylla; peritheciis globosis, atro-brun-
neis, in maculas orbiculares dense stipatis, primo cuticula tectis,
demum emergentibus; ascis clavato-cylindricis; sporidiis elongato-
ellipticis, uniseptatis, hyalinis, 20 \approx 4.

Hab. in foliis *Quercus aquaticaæ*, Darien Georgia (RAVENEL).

171. **Sphærella Phellos** (Schw.) Cooke Journ. of Bot. 1883, p. 106, 2642

Sph. Phellos Schw. n. 2805. — Hypophylla; peritheciis paucis sparsim in macula griseo-fusca aggregatis, pagina aversa innatis, prominulis, subglobosis, minutis, nigris, pertusis; ascis clavatis; sporidiis arcte ellipticis, uniseptatis, hyalinis, $8-10 \approx 2,5$.

Hab. in foliis *Quercus Phelli*, Carolina infer. Amer. bor.

172. **Sphærella antarctica** Speg. F. Fueg. n. 194. — Maculis nullis; 2643 peritheciis hypophyllis, sparsis, parenchymate immersis, epidermide velatis, vix prominulis e globoso conicis, $80-100 \mu$. diam., atris, glabris, coriaceo-membranaceis, contextu majuscule parenchymatico, olivaceo; ostiolo papillulato; ascis cylindraceis, non vel vix obclavulatis, apice rotundatis, tunica vix incrassata, basi subcoarctatis ac stipite brevissimo, noduloso suffultis, $25-30 \approx 3,5-4$, octosporis, a paraphysatis; sporidiis elongato-subclavulatis, utrinque acutiusculis, $7,5-8,5 \approx 1,5$, medio septatis non constrictis, leniter curvulis, hyalinis.

Hab. ad folia dejecta *Fagi antarcticæ*, ubique vulgata in Fuegia.

173. **Sphærella Castagnei** Har. et Br. Rev. Myc. 1891, p. 18, *Sph. 2644 Celtidis* Har. et Br. Rev. Myc. 1890, p. 177, nec Pass. — Peritheciis hypophyllis, congregatis, confertis, nigris, minutissimis, $70-80 \mu$. diam., maculas atras, 1-1,5 mm. diam. formantibus; ascis cylindraceo-fusoideis, sessilibus, 8-sporis, $32-40 \approx 8$; sporidiis distichis vel conglobatis, fusoideis, obtusulis, 1-septatis, subconstrictis, $12-16 \approx 3-4$, hyalinis.

Hab. in foliis emortuis *Celtidis australis*, Marseille Galliae (CASTAGNE).

174. **Sphærella Celtidis** Pass. Fung. nuovi, IV, p. 4, n. 7 et in Roum. 2645 Fungi selecti exs. n. 5043. — Peritheciis hypophyllis, sparsis vel subgregariis, erumpentibus, minutis, globoso-conicis, ostiolo acuto, atris; ascis subclavatis, a paraphysatis?; sporidiis elongatis, medio septatis vix vel minime constrictis, hyalinis, $22,5 \approx 6,5$, loculo altero angustiore.

Hab. in foliis dejectis *Celtidis australis* in Horto botanico, Parma.

175. **Sphærella morifolia** Pass. in Erb. Critt. Ital. ser. II, n. 1464. 2646 — Peritheciis hypophyllis, sparsis vel subgregariis, superficialibus, globosis, atris; ascis brevibus, ut plurimum curvulis, basi inflatis, inaequilateralibus, $50-55 \approx 10-15$; sporidiis distichis vel acervato-subtristichis, oblongo-cuneatis, ad septum non vel leniter constrictis, $17-25 \approx 5-5\frac{1}{2}$, hyalinis.

Hab. in foliis dejectis Mori albæ, Parma Ital. bor.

176. **Sphærella alnicola** Peck 40 Rep. p. 68. — Peritheciis parvis, 2647
85-113 μ . lat., hypophyllis, botryoideis vel sparsis, nudis, nigris; ascis oblongis vel subclavatis, 50-62,5 \approx 12,5-15; sporidiis confertis, lanceolatis, 1-septatis, saepe subincurvis, 22,5-27,5 \approx 4.

Hab. in foliis emortuis Alni viridis, Mt. Marcy Amer. bor.

— *Sphærellæ Alni-viridis* affinis sed peritheciis majoribus, stromate dematiaceo carens et sporidiorum forma diversa.

177. **Sphærella minutissima** Peck 40 Rep. p. 68. — Peritheciis nu- 2648
merosissimis, foliorum paginam inferam occupantibus, minutissi-
mis, 50-63 μ . lat., epidermide velatis, nigris; ascis oblongis vel
sursum subangustatis, 40-50 \approx 7,5-10; sporidiis confertis, oblongis,
rectis, obscure septatis, 15-17,5 \approx 4.

Hab. in foliis emortuis Alni incanæ, Adirondack mountains
Amer. bor.

178. **Sphærella Opuntiæ** E. et E. Journ. Myc. 1888, p. 97. — Peri- 2649
theiciis gregariis, 100-112 μ . diam., in cæspitulos 2-10 mm. diam.
confertis; ostiolis erumpentibus, globosis, imperfecte 4-sulcato-fis-
sis; ascis oblongo-cylindraceis, 60 \approx 8-9, sessilibus, aparaphysatis;
sporidiis distichis, clavato-fusoideis, 1-septatis, ad septum subin-
curvulis, guttatis, flavidis, 20-22 \approx 3,5.

Hab. in foliis emortuis Opuntiæ, Louisiana Amer. bor.

179. **Sphærella Macluræ** Ell. et Ev. Proc. Phil. 1890, p. 231. — Ma- 2650
culis rubro-brunneis margine saturatiore cinctis, 3-10 cm. diam.
vel confluendo latioribus, fragilibus, centro pallidioribus et mox
laceratis; peritheciis innatis, minutis 75 μ . diam., vertice erumpen-
te; ascis oblongo-cylindraceis, 50 \approx 8-10, aparaphysatis; sporidiis
biseriatis, oblongo-piriformibus, constrictis, leniter curvis, 12-14 \approx
5; apicibus subacutis.

Hab. in foliis Macluræ aurantiacæ, Emma, Saline Co. Mo.
Amer. bor. (C. H. DEMETRIO).

180. **Sphærella sycophila** Winter in Hedw. 1886, pag. 101 (*sycophila*). 2651
— Peritheciis in macula exarida, permagna, usque 35 mm. lata, ir-
regulari, fusca, late viridi cincta, indeterminata, hypophylla grega-
riis, immersis, globosis, poro punctiformi pertusis, membranaceis,
atris demum hemisphærice prominulis, 60-70 μ . latis; ascis obcla-
vatis, e basi crassiori sursum parum attenuatis, sessilibus, octospo-
ris, 24-26 \approx 8; sporidiis congregatis, fusoideis, utrinque attenuatis
sed rotundatis, inæquilateralibus et interdum parum curvatis, me-
dio 1-septatis, haud constrictis, hyalinis, 12 \approx 2.

Hab. in foliis Fici cuiusdam in insula S. Thomé Afr. occ.

181. **Sphærella Podocarpi** Cooke Journ. of Bot. 1883, p. 106. — Amphi- 2652
gena; peritheciis numerosis, atris epidermide tectis; ascis cylindri-
co-clavatis; sporidiis elongato-ellipticis, uniseptatis, hyalinis, $12 \approx 3$.

Hab. in foliis Podocarpi, Java (KUNZE). — Perithecia sub-
inde totam folii paginam occupant.

182. **Sphærella Taxodii** Cooke in Rav. Fungi Amer. n. 686. — 2653
Amphigena; peritheciis sparsis, subprominulis, atris 130μ . diam.,
poro pertusis; ascis cylindricis; sporidiis arcte ellipticis, unise-
ptatis hyalinis, $8 \approx 2,5$.

Hab. in foliis Taxodii distichi, Carolina inf. Amer. bor.
(RAVENEL).

183. **Sphærella conigena** Peck in 33 Rep. St. Mus. p. 34. — Peri- 2654
theciis minutis, erumpentibus, nigris; ascis latis, obovatis vel sub-
clavatis, sæpe apice acutis; sporidiis oblongis vel subcylindraceis,
denuo uniseptatis, $25-40 \mu$. longis.

Hab. in conis Thujæ in montibus Helderberg Amer. bor.

184. **Sphærella Retinosporæ** Berl. et Bres. Micr. Trid. p. 27, t. II, 2655
f. 7. — Peritheciis sparsis, minutis, $90-100 \mu$. diam., epidermide
tectis, pertusis, contextu parenchymatico, atro; ascis roseolatis v.
fasciculatis, sessilibus, clavatis, curvis, $30-35 \approx 9-10$; sporidiis octo-
nis, oblongis vel leniter cuneatis, ad medium 1-septatis, non con-
strictis, $9-11 \approx 3-3,5$, hyalinis.

Hab. in foliis emortuis Retinosporæ squarrosæ prope
Trento Tiroliæ cisalpinæ.

185. **Sphærella acicola** Cooke et Hark. in Grevillea XIII, p. 20. — 2656
Sparsa, minuta, subinnata; peritheciis globoso-depressis, membra-
naceis, fuscis; ascis breviter clavatis, octosporis; sporidiis clavato-
ellipticis, uniseptatis, hyalinis, flavidis, loculis altero subgloboso,
altero angustiore subconico, $7 \approx 3$.

Hab. in acubus Pini in California (HARKNESS).

186. **Sphærella Peckii** Sacc., *S. conigena* Peck 38 Rep. Stat. Mus. 2657
p. 104, nec ejusd. 33 Rep. — Peritheciis minutis, sparsis v. gregariis,
vix prominulis, erumpentibus, nigris; ascis subcylindraceis, $60-85 \approx$
 12 ; sporidiis stipatis oblongo-clavatis, uniseptato-constrictis, $10-$
 $12 \approx 4-5$, loculis inæqualibus, inferiore acutiore, superiore crassio-
re rotundatoque.

Hab. in conis dejectis Abietis canadensis in montibus Hel-
derberg Amer. bor. — Differt a *S. Pinsapo* longitudine ascorum
et fabrica sporidiorum nec non matrice.

187. **Sphærella Andersonii** Sacc. *Sphærella conigena* Ell. et Ev. Proc. 2658

Phil. 1890, p. 230, nec Peck. — Peritheciis gregariis, minutis, apice nigro, levi, conico-papilliformi excepto, immersis; ascis anguste clavato-cylindraceis, subitus sensim attenuatis, 75-80 \approx 5, aparaphysatis; sporidiis monostichis, ovoides, 1-septatis, ad septum constrictis, hyalinis, 6-7 \approx 3-3,5.

Hab. in apice squamarum strobili *Abietis Douglasii*, Belt Mts. Montana Amer. bor. (F. W. ANDERSON).

III. In Monocotyledoneis.

188. **Sphærella Gastonis** Sacc. in Rev. Myc. 1885, p. 158. — Peritheciis velatis, punctiformibus laxe gregariis; ascis fusoideis, sessilibus, 40 \approx 10; sporidiis oblongis, 1-septatis 7-8 \approx 2-3, hyalinis.

Hab. in foliis *Cocos nuciferae* in insula Tahiti (GAST. BRUNAUD).

189. **Sphærella sabaligena** Ell. et Ev. in Journ. of Myc. 1886, p. 101. 2660

— Peritheciis gregariis, 100-125 μ . diam., cuticula cinerescente, vix rupta tectis; ascis subovatis, 22-25 \approx 12-15; sporidiis stipatis, oblongo-clavatis, 1-septatis, ad septum constrictis, 10-12 \approx 3,5-4, subhyalinis.

Hab. in foliis emortuis *Sabalae Palmetto* in America boreali (LANGLOIS).

190. **Sphærella incisa** Ell. et M. in Journ. of Myc. 1885, pag. 99. — 2661

Peritheciis membranaceis, gregariis, globosis vel depresso globosis, $\frac{1}{4}$ mm. diam., aparaphysatis, epidermide nigrefacta tectis; sporidiis fusiformibus, utrinque appendicula acuta præditis, endochromate distinete bipartito, pallide luteis, 40-50 \approx 3-4.

Hab. in petiolis emortuis *Sabalae serrulatae*, New Florida Amer. bor.

191. **Sphærella bacillifera** Karst. in Hedw. 1883, n. 12, p. 31 et Rev. 2662

atque Synop. Ascom. fenn. p. 68. — Peritheciis sparsis vel subgregariis, subcutaneis, subsphaeroideis, glabris, ostiolo brevi erumpente, atris, exiguis; ascis elongato-clavatis, curvatis, apice rotundatis, vix stipitatis, 80-105 \approx 20-24; sporidiis octonis, tristichis, bacillari-fusoideis, curvulis, uniseptatis, chlorino-hyalinis, 45-60 \approx 6.

Hab. in foliis *Scheuchzeriae palustris* emortuis in parœcia Tammela Fenniae.

192. **Sphærella caricicola** Fuck., Winter in Hedwigia 1886, p. 19, *Læstadia caricicola* (Fuck.) Sacc. Syll. I, n. 430. — Ascii 40-44 \approx 10-

11; sporidia bicellularia, utriusque rotundata, ad septum constricta, hyalina, $14 \approx 4$.

193. **Sphærella Pontederiæ** Peck 40 Rep. p. 69. — Maculis majusculis, 12-20 μ . long., oblongis vel ellipsoideis, interdum confluentibus, supra brunneis, subtus nigrescenti-brunneis vel griseolis; peritheciis minutis, 75 μ . lat., hypophyllis, nigris; ascis oblongis v. subfusoideis, 50-63 \approx 12,5; sporidiis confertis vel biseriatis, oblongo-clavatis, uniseptatis, interdum 4-guttulatis, 15-20 \approx 5-6,5.

Hab. in foliis languidis Pontederiæ cordatae, Whitehall, Washington County Amer. bor. — *S. Caladii* subaffinis.

194. **Sphærella microscopica** Pass. in Brun. Nouv. Fragm. p. 27. — Peritheciis epidermide innatis, punctiformibus, minutissimis, sparsis, brunneis, membranaceis; ascis minutissimis, obovatis vel piriformibus, saepe gibbosis, 25 \approx 12; sporidiis inordinatis, lanceolatis, medio 1-septatis, hyalinis, 10 \approx 2,5-3.

Hab. in caulis emortuis Typhæ angustifoliæ, Rochefort, Saintes Galliæ.

195. **Sphærella tædiosa** Pass. in Brun. Champ. Saint. V, p. 2. — Peritheciis sparsis, tectis, contextu celluloso, fuligineo, parvis, ostiolo acuto, nigro, vix erumpente; ascis forma variabili, obovatis vel basi ventricosis apiceque attenuatis, rectis vel curvis, 8-sporis, 20-32 \approx 10-12,5, aparaphysatis; sporidiis 2-3-stichis, vel inordinate congestis, oblongis, utrinque rotundatis, hyalinis, 7,5-10 \approx 2,5-3, septo ope chlorurojoduri zinci tantum perspicuo.

Hab. in caulis emortuis Sparganii ramosi, Saintes, Saint-Sever, Rochefort Galliæ.

196. **Sphærella subcongregata** Ell. et Ev. in Journ. of Myc. 1886, p. 101. — Peritheciis sparsis, depresso globosis, $1/6$ mm. diam., cuticula nigrescente, ostiolo minuto perforata tectis; ascis oblongis, sessilibus, circiter 35 \approx 7; sporidiis distichis vel stipatis, hyalinis fusiformi-oblongis vel clavato-oblongis, 1-septatis, 9-11 \approx 2,5.

Hab. in caulis emortuis Smilacis, Newfield Amer. bor., soia *Diplodina Smilacis*.

197. **Sphærella Tassiana** De Not. — Sacc. Syll. I, p. 530. — Var. **rupe fortensis** Pass. in litt. Brun. in Rev. Myc. 1886. p. 205. — Ascis oblongis, 37 \approx 12; sporidiis inordinatis, oblongis, hyalinis, 1-septatis.

Hab. in foliis emortuis Typhæ latifoliae in Horto botanico Rochefort Galliæ.

198. **Sphærella Evansiæ** Pat. in Rev. Myc. 1886, pag. 83. — Peritheciis brunneis, minutis, 120-150 μ . diam., sparsis, primum epi-

dermide tectis dein emersis; ascis subsessilibus, aparaphysatis, 50-90 \approx 18-22, 8-sporis; sporidiis distichis, bilocularibus, altero rotundato, altero vero attenuato, septo parum constrictis, hyalino-viridulis, 15 \approx 4.

Hab. in foliis emortuis Evansiæ fimbriatae, Tibet Asiae centralis (Ab. DAVID).

199. **Sphærella Morææ** Pass. Diagn. Fung. nuov. III, n. 11. — Pe- 2669
ritheciis sparsis vel subgregariis, tectis, dein liberis, subglobosis; ascis aparaphysatis oblongo-clavatis, substipitatis, 65-87 \approx 15-20, 8-sporis; sporidiis oblongo-spathulatis, subtristichis, ad septum non vel vix constrictis, hyalinis, 20-25 \approx 7,5-10.

Hab. in caulinis, pedunculis foliisque aridis Morææ sinensis in Horto botanico, Parma. — *Sphærellæ Iridis* Auersw. affinis, sed ascis sporidiisque grandioribus diversa.

200. **Sphærella Hemerocallidis** Pass. Diagn. Fung. nuovi I, n. 33. — 2670
Peritheciis gregariis vel sparsis, epidermide fusca tectis, subglobosis, ostiolo acuto, matricem plus minus infuscantibus; ascis ovato-oblongis, vix stipitatis, 8-sporis, 45-57 \approx 20; sporidiis tristichis, oblongo-ovatis, utrinque rotundatis, tenuiter medio septatis, non constrictis, 17,5 \approx 7,5, hyalinis.

Hab. in caulinis siccis Hemerocallidis fulvæ, Vigheffio prope Parma. — Ascorum forma et sporidiis non constrictis a *Sph. Pale* Sacc. videtur distinguenda.

201. **Sphærella Dioscoreæ** Pass. Diagn. Fung. nuovi, I, n. 32. — Pe- 2671
ritheciis minutissimis, sparsis; ascis aparaphysatis!, obovatis vel subventricosis, 8-sporis, 50-75 \approx 10-15; sporidiis di- tristichis, oblongis, uniseptatis, loculis subæqualibus, hyalinis, 20 \approx 5-6,5.

Hab. in caulinis Dioscoreæ Batatae in Horto botanico, Parma. — An *Didymella Dioscoreæ* (B. et C.) Sacc. Syll. I, p. 560 huc spectet, e diagnosi brevi et manca haud facile judicandum.

202. **Sphærella fusca** Pass. Diagn. Fung. nuovi, I, n. 31. — Peritheciis minutis, pustulæformibus, laxe gregariis epidermide fuscata tectis, ostiolo papillæformi atro emergente, maculas oblongas, nigricantes, in matrice dealbata efficientibus vel matricem omnino nigrificantibus; ascis ovatis, basi brevissime constricto-stipitatis, superne attenuatis, 8-sporis, 45-50 \approx 17; sporidiis subtristichis, oblongis, utrinque rotundatis, medio septatis, non constrictis, hyalinis, 17,5 \approx 5.

Hab. in caulinis foliisque aridis Gladioli segetum, Vigheffio prope Parma.

203. **Sphærella minimipuncta** Cooke Rav. Amer. Fungi n. 681, Journ. 2673
Bot. 1883, p. 136. — Sparsa vel aggregata; peritheciis punctiformibus, emergentibus, atris; ascis clavatis, breviter stipitatis; sporidiis ellipticis, continuis (sempre?), hyalinis, 8-3.
- Hab.* in caulis Gladioli, S. Carolina Amer. bor.
204. **Sphærella Orchidearum** Karst. Rev. atque Synop. Ascom. fenn. 2674
p. 68. — Peritheciis sparsis vel gregariis, hinc inde aggregatis, epidermide tectis, lentiformibus, atris, nitentibus, saepe basi hyphis fuligineis, radiantibus obsitis, demum poro pertusis 0,2 mm. circ. lat.; ascis cylindraceo-clavatis; sporidiis ignotis; paraphysibus nullis.
- Hab.* in caulis aridis Gymnadeniæ conopseæ in Mustiala.
205. **Sphærella peregrina** Karst. — Sacc. Syll. I, p. 528 — var. **minima** (Johans.) Berl. et Vogl. Add. Syll. p. 83, *Mycosphærella peregrina* var. *minima* Johans. Svamp. Island p. 166. — Omnibus partibus minor; peritheciis subgregariis, strias formantibus, 270-370 μ . diam.; ascis 20-27 \approx 10-14; sporidiis congregatis, rectis vel curvulis, ad septum non constrictis, 11-16 \approx 2-3,5 (raro 20 μ . longis).
- Hab.* in foliis et calamis Scirpi cæspitosi, Hofios Islandiæ borealis.
206. **Sphærella Philodendri** Pat. Bull. Soc. Myc. 1888, pag. 112. — 2676
Peritheciis minutis amphigenis, nigris, glabris, centro ostiolatis, maculis albis late brunneo- v. fulvo-cinctis insidentibus; ascis apice obtusis, basi incrassatis, subsessilibus, 33-40 \approx 10, a paraphysatis; sporidiis octonis, rectis, cylindraceis, utrinque rotundatis, hyalinis, 1-septatis, loculis inæqualibus, 11-13 \approx 3.
- Hab.* in foliis vivi Philodendri cujusdam in Puerto-Zamuro Venezuela.
207. **Sphærella junciginea** Cooke Grev. XIX, p. 3. — Peritheciis in 2677 maculis orbicularibus gregariis, atris, parenchymate immersis, in acervulos elongatos phyllachoriformes, epidermide nigrificata velatos, plerumque subparallele aggregatis, sphæroideis, demum rima longitudinaliter exposita; ascis cylindraceis, octosporis; sporidiis subfusiformibus, didymis, medio vix constrictis, rectis curvulisve, luteo-hyalinis, 12-14 \approx 3-3,5.
- Hab.* in culmis Junci vaginati, Stewart Island Novæ Zealandiæ (KIRK). — Affinis *Sph. phyllachoroidi*.
208. **Sphærella præparva** Pass. et Beltr. F. Sicul. n. 7. — Peritheciis minutissimis, sparsis epidermide innatis, atris, 40-45 μ . diam.; ascis obovatis, basi abrupte constrictis, 30 μ . long.; sporidiis tri-

stichis, fusiformibus, curvulis, medio septatis, non constrictis, hyalinis, 22-25 \approx 4½.

Hab. in calamis et bracteis *Junci acuti*, Licata Siciliæ.

209. **Sphærella depressa** (Berk.) Cooke. Huc trasferenda *Physalospora depressa* (Berk.) Sacc. Syll. I, n. 1709, nam sporidia lanceolata, 1-septata, hyalina.

210. **Sphærella Rostkoviæ** Speg. F. Fueg. n. 202. — Maculis nullis 2679 vel late ambientibus, indeterminatis e pallescenti fuscescentibus; peritheiis sparsis immersis, vix prominulo-lenticularibus, 90-95 μ . d. atris glabris coriacellis, contextu densiuscule parenchymatico fusco-atro; ascis ovatis, antice truncato-rotundatis, crassissime tunicatis, latissime 1-foveolatis, postice rotundato-coarctatis, 26-28 \approx 8, stipite brevissimo noduloso suffultis; sporidiis inordinatis elliptico-clavulatis, utrinque obtuse rotundatis, 8 \approx 1,5-2, primo grosse 4-guttulatis, dein 1-septatis non constrictis, hyalinis.

Hab. in culmis dejectis putrescentibus *Rostkoviæ* (*Junci*) grandifloræ in Staten island Fuegiæ.

211. **Sphærella parallelogramma** Rehm in Hedwigia 1885, p. 239. 2680 — Periteciis minutissimis, globosis, atris, gregarie confertis, in maculis epidermidis parallelogrammiformibus, griseis vel griseo-atris, 2-5 mm. long., 1-2 mm. lat., secundum longitudinem culmorum conspicuis, sessilibus, primitus immersis, glabris, parenchymate fuscō contextis; ascis pseudo-piriformibus, latis, basi sessilibus, crassis, 8-sporis, 25-27 \approx 9; sporidiis polystichis, fusiformibus, acutis, rectis, 2-cellularibus, cellulis æqualibus, sœpe binucleatis, hyalinis, 10-12 \approx 2; paraphysibus nullis.

Hab. ad culmos sicclos gramineos prope glaciem æternam montis Ortler prope Sulden in Tirolia (DR. REHM).

212. **Sphærella Bambusæ** Pat. Bull. Soc. Myc. 1888, p. 112. — Periteciis globosis, minutissimis, 50-70 μ ., gregariis, numerosis, nigris, cellularibus, apice poro pertusis; ascis clavatis, apice obtusis, subsessilibus, 8-sporis, a paraphysatis; sporidiis hyalinis, ovoido-elongatis, rectis, 1-septatis, loculis æqualibus, 11-13 \approx 2-2,5.

Hab. in culmis *Bambusæ* putridis in Caracas Venezuelæ.

213. **Sphærella Cruris-galli** E. et K. Journ. Myc. 1887, p. 105. — 2682 Periteciis immersis, utrinque perspicuis sed supra magis prominulis, æqualiter sparsis vel minute gregariis, globosis, 100-115 μ . diam., ostiolo lato, rotundo; ascis oblongis, 50-55 \approx 10-12, a paraphysatis; sporidiis confertis, oblongo-fusoideis, subinæquilateralibus, 1-septatis, plerumque ad septum constrictis, hyalinis, 14-16 \approx 4.

Hab. in foliis aridis Panici Cruris-galli, Manhattan Americae borealis (SWINGLE).

214. **Sphærella exitialis** Mor. Enum. F. Bol. IV, p. 10. — Peritheciis non maculicolis, adproximatis v. saepius sparsis, amphigenis, globosis, epapillatis, ostiolo majusculo præditis, erumpentibus, saturate brunneis, 75-90 μ . diam., contextu e cellulis polyhædricis, 6-9 μ . diam. formato; ascis cylindraceis, subincurvulis, subtus saepius ventricosis, breve pedicellati, 45-24 \approx 16-20; sporidiis subdistichis vel confertis, hyalinis, cylindraceo-ovoideis, 1-septatis, loculo uno breviore et crassiore, altero longiore et tenuiore, 14-16 \approx 5-6.

Hab. in culmis languidis Tritici vulgaris prope Mezzolara in agro Bononiensi Italiæ. *Triticico exitialis*.

215. **Sphærella Spartinæ** E. et E. Journ. Mycol. 1888, p. 97. — Peritheciis laxe parenchymaticis, subastomis, ellipsoideis, 170-190 \approx 100-112, immersis, æqualiter denseque sparsis; sporidiis oblongo-ellipsoideis, flavidis, medio constrictis et septo parum conspicuo præditis, 4-guttulatis, 12-16 \approx 4,5-5,5.

Hab. in foliis emortuis et vaginis Spartinæ cynosuroidis prope Lincoln, Nebraska Amer. bor. — Ascii immaturi sed evidenter præsentes.

216. **Sphærella Hordei** Karst. Hedw. 1890, pag. 271. — Peritheciis sparsis, subcutaneis, epiphyllis, sphæroideis vel sphæroideo-depressionis, atris vel fusco-atris, osculo hiantibus, 0,1 mm. vel paullo ultra; ascis fusoideo-elongatis, subinde basi plus minus ventricosis obliquisque, 55-62 \approx 13-16; sporidiis octonis, di- vel subtristichis, oblongatis vel fusoideo-oblongatis, uniseptatis, ad septum constrictis, loculis 2-guttulatis rectis, hyalinis, 18-24 \approx 6-8.

Hab. in foliis Hordei vulgaris languescentibus in parœcia Libelits Fenniæ (ERNST SAVANDER). — *Hordeo noxia*.

217. **Sphærella graminicola** Fuck. — Syll. f. I, p. 528 — *f. alpina* 2686 Rehm Ascom. n. 794. — Peritheciis grosse cellulosis, brunneis, calvis; ascis ovoideis, vertice incrassatis, sessilibus, 8-sporis, 36 \approx 15; sporidiis oblongis subclavatis, rectis, 1-septatis, pluriseriatis, 15 \approx 4.

Hab. in foliis emortuis siccis Graminum in alpinis am Ortler Tiroliæ, ad 2000 m. circ.

218. **Sphærella Dactylidis** Pass. Diagn. F. nuovi, I, n. 37. — Peritheciis sparsis, erumpentibus, tectis, ostiolo acuto; ascis ex ovata basi attenuatis, rectis vel gibbis, breviter abrupte stipitatis, 45-50 \approx 20-

23, 8-sporis; sporidiis conglobatis vel subtristichis, obovato-oblongis, utrinque rotundatis, uniseptatis, non vel vix constrictis, 18-19 \approx 7-8, hyalinis, endoplasmate opaco, granuloso.

Hab. in culmo et in rachide *Dactylidis* glomeratae, Vigheffio prope Parma.

219. **Sphærella Eulaliæ** Pass. Diagn. Fung. nuovi, I, n. 36. — Maculis nullis; peritheciis punctiformibus, sparsis vel in series linea-
res breves digestis, ostiolo vix perspicuo, primo tectis, dein plus minus nudatis, erumpentibus; ascis ovato-oblongis, basi gibbo-ven-
tricosis, breviter abrupte stipitatis, 50 \approx 15, 8-sporis; sporidiis sub-
tristichis, ovoideo-oblongis, rectis, ad septum leviter constrictis,
hyalinis, loculis varie guttulatis, altero crassiore, 22,5 \approx 7,5.

Hab. in foliis siccis *Eulaliæ japonicæ* in Horto botanico,
Parma.

220. **Sphærella maydina** Pass. Diagn. Fung. nuovi, I, n. 35. — Peritheciis minutissimis, superficialibus, sparsis, cellulis fuligineis gran-
diusculis contextis; ascis parvulis, ovato-oblongis, subsessilibus, 8-
sporis, 25 \approx 10; sporidiis conglobatis, oblongis, prope medium se-
ptatis et breviter constrictis, utrinque rotundatis, hyalinis, 10 \approx 3,5.

Hab. in culmis putridis *Zeæ Maydis*, Vigheffio pr. Parma.
— A *Sph. zeina* peritheciis multo minoribus e matrice griseola
oriundis præcipue distinguenda.

221. **Sphærella zeina** Pass. Diagn. Fung. nuovi, I, n. 34. — Peritheciis 2690
in matrice dealbata erumpentibus, sparsis, minutis, globosis, atris;
ascis ovatis vel oblique ovatis, 25 \approx 14-15, 8-sporis; sporidiis ob-
longis, non constrictis, didymis, hyalinis, 10 \approx 2,5.

Hab. in culmis fracidis *Zeæ Maydis*, Vigheffio pr. Parma.

222. **Sphærella Triseti** Speg. F. Fueg. n. 203. — Maculis nullis; pe- 2691
ritheciis hypophyllis sparsis vel laxe gregariis ac subseriatibus, im-
mersis, non v. vix prominulis, lenticularibus, 70-80 μ . diam., tenui-
membranaceis, contextu parenchymatico fuligineo, ostiolo latissimo,
20 μ . diam., impresso perforatis, globosis, levibus; ascis cylindra-
ceo-ovatis, antice truncato-rotundatis, crasse tunicatis non foveo-
latis, postice brevissime cuneatis, 24-26 \approx 8, stipite brevissimo cras-
so nodulo suffultis, octosporis, aparaphysatis; sporidiis di- poly-
stichis, subclavulato-ellipticis, utrinque truncato-rotundatis, 10-12
 \approx 2,5, medio 1-septatis constrictulis, loculo supero nonnihil cras-
siore, hyalinis.

Hab. ad folia languida *Triseti* in summis montibus in Sta-
ten island Fuegiæ.

223. **Sphaerella californica** Cooke et Harkn. Journ. of Bot. 1883, p. 136. — 2692
 — Peritheciis exiguis, sparsis, subsphæroideis, innato-prominulis, nigro-fuscis, poro pertusis; ascis clavato-cylindricis; sporidiis arcte ellipticis, uniseptatis, nec constrictis, hyalinis, 8 \approx 2.
Hab. in Graminaceis, California Amer. bor. (HARKNESS).
224. **Sphaerella philochorta** Cooke Journ. of Bot. 1883, p. 137. — 2693
 Epiphylla, sparsa; peritheciis minutis, globosis, prominulis, atris, epidermide ostiolo papillato pertusis, velatis; ascis clavatis; sporidiis arcte cylindrico-ellipticis, utrinque obtusis, uniseptatis, vix constrictis, hyalinis, 13-17 \approx 3.
Hab. in foliis graminum, Maine Amer. bor.
225. **Sphaerella epistroma** Cooke Journ. of Bot. 1883, p. 137. — 2694
 Culmicola, aggregata, innato-prominula, minutissima, nigra, striae breves, 2 mm. long., densas efficiens; ascis cylindricis; sporidiis fusiformibus, rectis curvulisve, uniseptatis, nec constrictis, hyalinis, 16-18 \approx 3.5.
Hab. in culmis graminum in Britannia.
226. **Sphaerella disseminata** De Not. et Car. Erb. critt. it. ser. II, n. 391, S. M. I, 2036, nec *Leptosphaeria disseminata* De Not. Recl. — Adde: ascis brevibus obclavatis, subsessilibus, 45-48 \approx 18, 8-sporis; sporidiis ovoido-oblongis, di-tristichis, uniseptatis vix constrictis, rectis curvulisve, 17-20 \approx 4, hyalinis.
227. **Sphaerella Mühlenbergiæ** Ellis in Amer. Nat. XVII, 1883, p. 317. — 2695
 — Peritheciis erumpentibus, minutis, plerumque elongato-serialibus; ascis oblongis, 35 \approx 9,5; sporidiis ellipticis, uniseptatis, 11-15 \approx 3-3,5; stylosporis in peritheciis majoribus, oblongo-fusiformibus, 13-19 \approx 4, binucleatis.
Hab. in foliis Mühlenbergiæ, Newfield New Jersey in America boreali (ELLIS).
228. **Sphaerella Zizaniæ** (Schw.?) Ell. Journ. Mycol. 1887, *Sphaeria* 2696 Schw. Syn. N. Amer. n. 175 (?). — Peritheciis immersis, sparsis, non seriatis; ascis tereti-oblongis, turgidis, 45-55 \approx 10-12; sporidiis distichis clavato-oblongis, 1-septatis constrictisque, hyalinis, rectis curvulisve, 15-20 \approx 5-6.
Hab. in foliis languidis Zizaniæ aquatice, Delaware Amer. bor. (COMMONS). — Videtur species Schweinitzii, cuius vero exempl. originalia sunt sterilia.
229. **Sphaerella loliacea** Pass. Diagn. Fung. nuovi, I, n. 38. — Peritheciis minimis, punctiformibus, sparsis vel laxe gregariis; ascis

parvulis obpiriformibus vel clavatis, aparaphysatis, 8-sporis; sporidiis oblongo-cuneatis, ad septum vix constrictis, hyalinis, 16 ± 5 .

Hab. in spicis aridis *Lolii perennis*, Vigheffio pr. Parma.
— Ascorum basi angusta ab affinibus *Sphærella Tassiana* et *S. Maydis* præcipue distinguenda.

IV. In Acotyledoneis.

230. **Sphærella altera** Pass. Diagn. Fung. nuovi, I, n. 39. — Peritheciis minutis, tectis, in seriem simplicem dispositis, maculas lineares formantibus, per epidermidem fissam vix perspicuis, contextu minuto celluloso, fuligineo; ascis aparaphysatis, oblongis, basi plus minus inflatis, ad apicem attenuatis, 45 ± 12 , 8-sporis; sporidiis subtristichis, oblongis, utrinque obtusis, prope medium septatis, vix vel non constrictis, endoplasmate granuloso, non guttulatis, hyalinis, 15 ± 5 .

Hab. in caulis et ramis putrescentibus *Equiseti ramosi*, Gajone prope Parma.

231. **Sphærella Trichomanis** Cooke in Grevillea XIV, pag. 12. — Sparsa vel subcæspitosa; peritheciis semiimmersis, punctiformibus atris, cuticulâ tenui primo tectis, punctato-pertusis; ascis clavatis, 30 ± 5 , octosporis; sporidiis cylindricis, utrinque rotundatis, uniseptatis, hyalinis, 8 ± 2 .

Hab. in frondibus vivis *Trichomanis* prope Samoa in Australia.

232. **Sphærella aquilina** Fr. — Sacc. Syll. I, p. 532 — var. **infera** 2700 Karst. in Hedw. 1884, n. 1, p. 4, Rev. atque Synop. Ascom. fenn. pag. 68. — A typo recedit peritheciis hypophyllis; ascis 42-58 \pm 8-9 sporidiisque ad septum vix vel leviter constrictis, $8-12 \pm 2-4$.

Hab. in frondibus *Pteridis aquilinæ* emortuis ad Mustiala Fenniæ.

233. **Sphærella Lycopodii** Peck 39 Rep. p. 51, t. I, f. 12-15. — Peritheciis minutis, 100μ . lat., nigrescentibus; ascis oblongis v. subcylindraceis, apicem versus sæpe subattenuatis, $30-40 \pm 10$; sporidiis oblongis $11,5-12,5 \pm 4-5$.

Hab. in squamis spicarum *Lycopodii clavati*, Adirondack mountains Amer. bor. — A *Sph. lycopodina* differt ascis sporidiisque minoribus.

234. **Sphærella parasitica** Wint. in Hedw. 1886, p. 19, *Epicymatia Winteri* Kunze Fungi selecti exsicc. n. 65. — Peritheciis gregariis, minutissimis, globoso-piriformibus, vertice pertusis, tenuis-

sime membranaceis, glabris, fuscis, $70 \approx 50\cdot60$; ascis elongato-ovoideis vel oblongis, utrinque attenuatis, sessilibus, 8-sporis, $24\cdot32 \approx 10\cdot12$; sporidiis inordinatis, conglobatis, bacilliformibus, anguste clavatis utrinque rotundatis, hyalinis, bicellularibus, haud constrictis, $14\cdot18 \approx 3,5$.

Hab. in *Cenangio fuliginoso* parasitans, in Germania.

MYCOSPHÆRELLA Johans. Svamp. Island. p. 163 (1884) pro minori parte (Etym. *myces* fungus et *Sphærella*). — Perithecia *Sphærellæ*. Asci 16-spori, aparaphysati. Sporidia ovata vel oblonga, 1-septata, hyalina *).

1. **Mycosphærella polyspora** Johans. Svamp. Island. pag. 164, 2703 tab. XXIX, fig. 7. — Peritheciis sparsis vel gregariis, globoso-conoideis, erumpentibus, atris, $80\cdot120 \mu$. diam.; ascis aparaphysatis, ovoideis saepe inæquilateralibus, sessilibus, 16-sporis, $36\cdot44 \approx 13\cdot18$; sporidiis conglobatis, ovato-clavulatis, utrinque rotundatis, medio vel prope medium septatis, guttulatis, rectis, hyalinis, $11\cdot14 \approx 4\cdot6$, septo vix constrictis.

Hab. in pedunculis et fructibus aridis *Azaleæ* procumbentis, Eskifjördur Islandæ orient.

APIOSPORA Sacc. Syll. I, p. 539.

1. **Apiospora Striola** (Pass.) Sacc. Syll. I, pag. 539. — Var. **minor** 2704 Berl. et F. Sacc. Contr. Myc. Lus. p. 3, t. 83, f. 3. — Peritheciis seriatim dispositis, immersis culmine superficiem rimose findentibus et elevantibus; ascis clavatis, $70\cdot80 \approx 12\cdot15$, octosporis; sporidiis subdistichis, oblongis, $20\cdot22 \approx 6$, basi 1-septatis, hyalinis.

Hab. in culmis emortuis *Arundinis Donacis*, Coimbra Lusitanæ.

2. **Apiospora Urticæ** Rehm Beitr. Ascom. in Hedw. 1882, p. 133. — 2705 Peritheciis gregariis, sub peridermio caulinum nidulantibus, dein liberis, subglobosis, sessilibus, in collum breviusculum, peridermium perforans protractis, nigris, circ. 0,4 mm. diam.; ascis cylindro-clavatis, teneris, 8-sporis $90 \approx 12$; sporidiis dacryoideis, nempe cellula superiore subrotunda, $\frac{3}{4}$ long. totius sporidii, inferiore sub-

*) Beat. Johanson pro nomine vastissimi generis *Sphærellæ* (quod nomen olim pro alga quadam usurpatum fuit) illud *Mycosphærellæ* proposuit; sed innovationem hanc inconsultam demonstraverunt cl. A. N. Berlese et G. B. De Toni (*Intorno al genere Sphærella*, Venezia 1887).

rotunda $\frac{1}{4}$; utraque cellula nucleis 2 parvulis prædita, hyalinis subdistichis, 18 \approx 10; paraphysibus articulatis, ramosis.

Hab. ad caules sicclos Urticæ, Hausstattalpe in monte Bene-dictenwand alpium Bavariæ (DR. ARNOLD).

STIGMATEA Fr. — Sacc. Syll. I, p. 541.

I. Eu-Stigmatae.

1. **Stigmata vexans** Wint. in Hedw. 1885, I. — Peritheciis in macula non determinata, ochracea vel fuscidula, epiphylla gregariis, depresso-hemisphæricis, dimidiatis, crasse membranaceis, atris ostio-lo punctiformi, sed distincto pertusis, 180-230 \approx 96-105; ascis oblongis vel e basi ventricosa sursum parum attenuatis, membrana crassa præditis, sessilibus, 8-sporis, 58-78 \approx 24-32; sporidiis conglobatis, didymis, olivaceis, cellula superiori fere globosa, 12-14 \approx 12, cellula inferiori obconica, 9-9,5 \approx 7-9; paraphysibus non distinctis, mucilagineis.

Hab. ad folia viva Brayeræ anthelminthicæ, in Abyssinia (KECK).

2. **Stigmata silvatica** Sacc. Mich. II, pag. 604. — Peritheciis gregariis vel hinc inde aggregatis, hypophyllis, subsuperficialibus, globulosis, 110-120 μ . diam., pertusis, solidiusculis; ascis tereti-clavulatis, 40-44 \approx 6-7, parce (spurie?) paraphysatis, octosporis; sporidiis distichis oblongis, deorsum acutiusculis, 10 \approx 3, uniseptatis, non vel vix constrictis, hyalinis.

Hab. in foliis languidis vel emortuis Juniperi communis, Fontainebleau Galliæ (THERRY).

3. **Stigmata Nicholsoni** Cooke Grev. XI, pag. 16. — Amphigena; peritheciis circinantibus, erumpentibus, nigris, nitidulis, convexis, denique collapsis, glabris; ascis clavatis; sporidiis ellipticis, utrinque rotundatis, 1-septatis, constrictis, 18 \approx 8, pallide fuscis.

Hab. in foliis Pruni lusitanicæ, Newcastle Britan. (G. NICHOLSON).

4. **Stigmata nitidula** (Lév.) Sacc. Sic nuncupanda *Sphærella nitidula* Sacc. Sylloge I, n. 1886.

II. *Stigmatula* Sacc.

5. **Stigmata nitens** Pat. Bull. Soc. Myc. 1888, p. 112. — Epiphylla, greges subcirculares, 3-5 mm. diam. formans; peritheciis hemisphæricis, numerosis, distinctis, superficialibus, atro-nitidis, ostiolo vix

Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Didymella. 661

prominulo; ascis brevibus, crassis, obtusis, $80 \approx 26-30$; sporidiis ovoideis, obtusis, continuis sed medio subconstrictis, hyalinis, $20 \approx 10$.

Hab. in foliis coriaceis vivis, Maipures Venezuelæ.

6. **Stigmatea Melastomatum** (Lév.) Sacc. Sic nuncupanda *Læstadia Melastomatum* (Lév.) Sacc. Syll. I, n. 1631.
7. **Stigmatea Francevilliana** Pat. Bull. Soc. Myc. 1887, p. 125, t. X, 2710 fig. 5. — Peritheciis convexis, nigris, nitidis, $\frac{1}{2}-1$ mm. diam., basi tantum epidermide insculptis, ostiolo prominulo; ascis piriformibus, breve stipitatis; paraphysibus copiosis, gracilibus ramosis; sporidiis ellipsoideis, hyalinis, continuis, $15-17 \approx 6$.

Hab. gregatim in caulis vetustis Rubiaceæ, Franceville Congo.

8. **Stigmatea Armandi** Pat. in Rev. Myc. 1886, p. 83. — Peritheciis punctiformibus, nigris, globulosis; ascis subsessilibus, $42-45 \approx 12-15$; paraphysibus linearibus; sporidiis uno apice attenuatis, $12-15 \approx 3-4$, hyalinis vel pallide viridulis.

Hab. in petiolis et foliis Clematidis Armandi, Tibet Orient. Asiæ (AB. DAVID).

9. **Stigmatea mucosa** Pat. in Rev. Myc. 1886, p. 82. — Peritheciis minutis, primum epidermide tectis, demum, ea rupta, prominulis, nigris, rotundis, vertice poro albo pertusis; ascis sessilibus, octosporis, $35-45 \approx 12-15$; sporidiis ovoideis, hyalinis, continuis, $12-15 \approx 4-5$; pseudo-paraphysibus gelatinosis, ramosis, massa mucosa ascos obvallantibus.

Hab. in foliis emortuis Agrostidis albæ, Tibet Oriental. Asiæ (AB. DAVID).

DIDYMELLA Sacc. Syll. I, pag. 545.

I. Herbicolæ.

1. **Didymella Fuchsiae** Cooke et Hark. in Grevillea XIV, pag. 9. — 2713 Sparsa vel subgregaria; peritheciis globoso-depressis, atris, epidermide tectis, ostiolo conico, prominulo emergente; ascis cylindrico-clavatis, octosporis; sporidiis biserialibus, rectis curvulisve, uniseptatis, nec constrictis, binucleatis, hyalinis, $15 \approx 5$.

Hab. in caule Fuchsiae in California (HARKNESS).

2. **Didymella Trifolii** (Fuck.) Sacc. Syll. I, pag. 554, *Didymosphaeria Trifolii* Winter in Hedwigia 1886, pag. 24. — Sporidia 17-20 $\approx 5-6$.
3. **Didymella picea** (Sollm.) Sacc. *Sphæria picea* Sollm. Hedw. 1863, 2714 n. 19, p. 192, t. XIII, f. 12-16, nec Pers. — Peritheciis inæqualiter seriatis, innato-erumpentibus, subglobosis, nigricantibus, minu-

tis, glabris, obtuse papillatis, hyphis mycelii fuligineis ramosis repentibus cinctis; ascis tereti-clavatis, apice rotundatis subsessilibus, paraphysatis; sporidiis distichis, hyalinis, obovatis, 1-septato-constrictis, 17-18 \approx 7.

Hab. in caulis siccis Trifolii pratensis prope Coburg Germaniae.

4. **Didymella Lupini** (Cooke et Hark.) Berl. et Vogl. Add. Syll. p. 88, 2715
Sphaeria Lupini Cooke et Hark. in Grev. XIII, p. 18. — Sparsa, punctiformis; peritheciis tectis, globoso-depressis, leniter papillatis; ascis clavatis; sporidiis ellipticis, uniseptatis, hyalino-flaveolis (immaturis?), plasmate granuloso, 15-17 \approx 6-7.

Hab. in caule Lupini in California (HARKNESS).

5. **Didymella Melonis** Pass. in Erb. Critt. Ital. Ser. II, n. 1465, 2716
 — Peritheciis sparsis, mycelio haud perspicuo, subglobosis ostiolo acuto; ascis cylindrico-clavatis, 8-sporis, 62-80 \approx 10-12; sporidiis distichis, constrictis, hyalinis, loculis biguttulatis, 17,5 \approx 1,5; paraphysibus brevioribus, paucis.

Hab. ad caules, ramos petiolosque aridos Cucumeris Melonis prope Parma Italiæ bor.

6. **Didymella Cookeana** Berl. et Vogl. Add. Syll. p. 87, *Sphaeria Me-garrhizæ* Cooke et Harkness in Grev. XIII, pag. 18. — Sparsa, tecta; peritheciis subglobosis, mox depressis, atris, ostiolo papillato; ascis subclavatis; sporidiis biserialibus, ellipticis, uniseptatis, vix constrictis, hyalinis, pallide luteolis, 12-14 \approx 6.

Hab. in Megarrhiza californica in California (HARKNESS). — «An a sequente vere diversa?»

7. **Didymella Megarrhizæ** Cooke et Hark. in Grevillea XIV, p. 9, 2718
 — Gregaria; peritheciis subconicis, atris, parvulis, nitidis, initio velatis, demum ostiolo acuto erumpentibus; ascis cylindrico-clavatis, octosporis; sporidiis ellipticis, uniseptatis, medio constrictis, rectis, hyalinis, 16-18 \approx 7.

Hab. in caule Megarrhizæ in California (HARKNESS).

8. **Didymella Rehmii** (J. Kunze) Sacc., *Didymosphaeria Rehmii* Kunze 2719 Fung. Sel. exsic. n. 1875 (absque diagnosi). — Peritheciis dense gregariis epidermide velatis, tumidulis, globoso-depressis, nigris, vix papillatis; ascis clavulatis, 80-90 \approx 10; paraphysibus filiformibus obvallatis, octosporis; sporidiis monostichis, ellipsoideis, obtusiusculis, uniseptatis, 4-guttulatis, leniter constrictis, 15-17 \approx 5, hyalinis.

Hab. in caulis subvivis Verbenæ officinalis pr. Eisleben Germaniæ.

9. **Didymella? nigrella** (Fr.) Sacc. *Sphæria nigrella* Fr. S. M. II, 2720 p. 512, Cooke Handb. n. 2728. — Peritheciis subglobosis, levibus, umbilicato-perforatis, nigris, maculæ elongatae determinate nigræ insidentibus; sporidiis distichis fusiformibus, hyalinis, plasmate bipartito 20 μ . longis.

Hab. ad caules Galeopsidum et Angelicæ in Suecia, Britannia, Belgio. — Macula constanter præsens, levigata. Perithecia superficiali-innata, umbilicata, intus albo-farcta.

10. **Didymella Angelicæ** (E. et E.) Sacc. *Sphærella Angelicæ* Ell. et 2721 Ev. Proc. Phil. 1890, pag. 231. — Peritheciis sparsis, subcutaneis cumque epidermide secedentibus, globosis, $\frac{1}{4}$ mm. diam., infra collabascentibus, ostiolo papillato, epidermidem perforante et vix exstante; ascis clavato-cylindraceis, copiose paraphysatis, 65-70 \approx 7-9; sporidiis biseriatis, oblongo-ellipsoideis, 2-nucleatis, 12-15 \approx 3,5, demum ovato-oblongis et 1-septatis.

Hab. in caulis emortuis Angelicæ atropurpureæ, London Canada Amer. bor. (DEARNESS).

11. **Didymella Heribaudii** Har. et Br. Rev. Myc. 1890, n. 47. — Pe. 2722 ritheciis minutis erumpentibus, nigris, numerosis, subinde confluentibus, irregularibus, oblongis vel punctiformibus, depressis; ascis oblongis, cylindraceis vel clavatis, sessilibus vel breve stipitatis, 8-sporis, 32-45 \approx 8; paraphysibus cohærentibus; sporidiis distichis, oblongis vel tereti-fusoideis, 1-septatis, obtusiusculis, hyalinis, 14-16 \approx 2,5-3.

Hab. in caulis siccis Thalictri minoris, an Puy de Crozel prope Clermont Ferrand Galliæ (HÉRIBAUD).

12. **Didymella chætostoma** Pass. Diagn. Fung. nuovi, III, n. 12. — 2723 Peritheciis sparsis subglobosis, tectis, ostiolo acuto setulis rigidis convergentibus 30-40 μ . long., obsito, erumpente; ascis cylindricis, breviter stipitatis, 8-sporis, 60-80 \approx 9-10; paraphysibus filiformibus stipitatis; sporidiis monostichis ovatis, prope medium septatis et constrictis, hyalinis, 20 \approx 7,5, loculo altero minore.

Hab. in caulis siccis Artemisiæ camphoratæ, Vigheffio prope Parma Italiæ bor..

13. **Didymella prominens** Ell. et Ev. in Journ. of Myc. 1886, p. 101. 2724 — Peritheciis erumpenti-superficialibus, sparsis, ovato-hemisphæricis, $\frac{1}{4}$ mm. diam., atris, rugosis, ostiolo prominulo, tuberculiflori-conoideo vel breviter cylindraceo, levi donatis; ascis clavato-

cylindraceis, subsessilibus, 40-45 \approx 6-7; paraphysibus filiformibus, cinctis, octosporis; sporidiis subfusoideis v. cymbiformibus, 1-septatis, hyalinis, 10-12 \approx 3-4, ad septum vix constrictis.

Hab. in caulis herbaceis emortuis Ambrosiæ trifidae in America boreali (LANGLOIS).

14. **Didymella superflua** (Fuck.) Sacc. Syll. I, p. 555 — var. **Humuli** 2725

Karst. Symb. ad Myc. Fenn. XVI (1884), pag. 150 et Rev. atque Synop. Ascom. fenn. pag. 61. — Peritheciis gregariis, primum epidermide tectis, dein denudatis, ut plurimum cupulato-collapsis; ascis cylindraceis, subsessilibus, 55-60 \approx 8-10; sporidiis 9-15 \approx 4-5.

Hab. ad caules exsiccatos Humuli Lupuli in Mustiala Fenniae. — Status pycnidicus est *Phoma sarmamentella*.

15. **Didymella inconspicua** Johans. Svamp. fran Island. pag. 167, 2726

t. XXIX, fig. 8. — Peritheciis sparsis, parenchymate innatis, epidermide leviter nigrefacta tectis, ostiolo rotundo pertusis, atris, glabris, minutis, 95-135 μ . diam.; ascis octosporis, paraphysatis, ovoideo-elongatis, saepe inaequilateralibus, 55-65 \approx 18-20; sporidiis distichis vel congregatis, ovoideo-elongatis vel cuneato-elongatis, utrinque rotundatis, prope basim 1-septatis, non constrictis, guttulatis, rectis, hyalinis vel chlorino-hyalinis, 20-23 \approx 7-9, loculo majore 15-18 μ . long., minore 4,5-6 μ . long.

Hab. in foliis emortuis Saxifragæ oppositifoliæ, Eski-fjördur Islandiæ bor.

16. **Didymella Chamæjasmes** (Fuck.) Berl. et Vogl. Add. Syll. p. 88, 2727

Sphaeria Chamæjasmes Fuck. in Reis. nach Nordpolarf. III, et Oudem. Contr. Fl. Myc. Nov. Seml. p. 152, tab. II, fig. 5. — Peritheciis gregariis, raro sparsis, sub epidermide nidulantibus, demum subliberis, minutis, atris, e basi globosa antice conicis, acutis; ascis oblongis, sessilibus, 8-sporis, 104 \approx 12; sporidiis oblique submonostichis, oblongis, rectis, utrinque obtusis, æqualiter didymis et ad septum facile diffractis, 16 \approx 6, hyalinis.

Hab. in caulis exsiccatis Androsaces Chamæjasmes, Nowaja Semlja.

17. **Didymella nivalis** (Fuck.) Berl. et Vogl. Add. Syll. p. 88, *Sphaeria* 2728

nivalis Fuck. in Die zweite deutsch. Nordpolarf. p. 93, t. I, f. 5. — Peritheciis sub foliorum caulinque aridorum epidermide albescente nidulantibus, gregariis, minutis, globosis, aterrimis, ostiolo atro, acuto, prominulo; ascis oblongo-ovatis, utrinque attenuatis, tunica crassa præditis, 8-sporis, 94 \approx 20; sporidiis congregatis subdi-

stichisve, oblongis, inæqualiter didymis, utrinque obtusis, ad septum parum constrictis, hyalinis, 20 ± 8 .

Hab. in caulis foliisque exsiccatis *Epilobii latifolii*, Kaiser-Franz-Josephs-Fjord in Groenlandia.

18. **Didymella Rehmiana** Bäuml. Mycol. III, p. 1. — Peritheciis gre-
gariis sub epidermide dealbata nidulantibus, demum prominulis, glo-
bulosis, fuscis, ostiolo rotundato pertusis, $100-120 \mu$. diam., con-
textu grosse parenchymatico, fuligineo; hyphis septatis 4μ . cr.,
basi cinctis; ascis tereti-clavatis, breve pedicellatis, 8-sporis, $45-$
 $50 \pm 8-10$; sporidiis distichis, $10-18 \pm 4-5$, utrinque rotundatis, re-
ctis curvulisve, primo continuis demum 1-septatis, medio non con-
strictis, hyalinis; paraphysibus vix visilibus.

Hab. in caulis aridis *Euphorbiæ palustris* pr. St. Georgium, Pressburg Austriæ.

19. **Didymella cornuta** Ell. et Ev. Proc. Phil. 1890, p. 232. — Pe. 2730
ritheciis sparsis, minutis, $175-200 \mu$. diam., sub epidermide late
atrata ostiolisque prominentibus perforata nidulantibus; ascis cy-
lindraceis, subsessilibus, $65-70 \pm 7$; paraphysibus immixtis; spori-
diis biseriatis, fusoides, flavidis, leniter curvulis, 1-septatis con-
strictisque, demum 3-septatis, $20-25 \pm 3-3,5$, utrinque subobtusis.

Hab. in caulis emortuis *Asclepiadis Cornuti*, London Canada (DEARNESS). — «Si sporidia constanter 3-septata, species
ad *Leptosphaeriam* ducenda videtur».

II. Arboricolæ.

20. **Didymella Cocciferæ** H. Fab. Sphér. Vaucl. II, p. 43, f. 12. — 2731
Peritheciis parce gregariis, nunc corticolis et semiinfossis, nunc
lignicolis et superficialibus globosis v. conoideis, vertice porulo per-
tusis, $1\frac{1}{2}-2\frac{1}{3}$ mm. latis; ascis paraphysibus filiformibus dense ob-
vallatis, cylindraceis, stipitatis, 8-sporis, parte sporifera, 100 ± 12 ;
sporidiis oblique monostichis, bi-conoideis, hyalinis, medio septatis
constrictisque, $18-20 \pm 8-9$.

Hab. ad ramos exaridos *Quercus Ilicis* humifusos in apri-
cis Vaucluse Galliæ.

21. **Didymella vagans** H. Fab. Sphér. Vaucl. II, p. 45, f. 16. — Pe. 2732
ritheciis laxe sparsis, solitariis epidermide nigrificata tectis, dein
nudis, globosis, $1\frac{1}{3}-1\frac{1}{2}$ mm. latis, vertice hemisphæricis et porulo
hiante pertusis; ascis dense paraphysatis, clavatis, stipitatis, 8-
sporis, $120-140 \pm 14-18$; sporidiis distichis, bi-conoideis medio con-

strictis septatisque, hyalinis, 4-guttatis, guttis binis intermediis multo majoribus, 27-30 \approx 10.

Hab. ad ramulos dejectos *Quercus cocciferæ* et *Ericæ arboreæ*, Vaucluse Galliæ.

22. **Didymella clavulata** Ellis Amer. Nat. 1883, p. 318. — Peritheciis membranaceis globosis, $1\frac{1}{9}$ - $1\frac{1}{6}$ mm. diam. rugosis, folio innatis, utraque pagina prominulis; ostiolo diametrum perithecii æquante, cylindraceo apice subclavato pertuso; ascis oblongo-cylindraceis, p. sporif. 35-40 \approx 5-6, paraphysatis; sporidiis distichis, oblongo-fusoideis, acutis, 4-nucleatis, denique inæqualiter 1-septatis et obtusiusculis, 7,5-9 \approx 2, flavidulis.

Hab. in pagina super. foliorum *Quercus* (nigræ?), Newfield N. J. Amer. bor.

23. **Didymella australis** H. Fab. Sphér. Vaucl. II, p. 44, fig. 14. — 2734 Peritheciis nunc sparsis, nunc congestis, cortice infixis, epidermide tectis, dein erumpentibus, seminudis, globosis, vertice poro pertusis, $1\frac{1}{2}$ - $2\frac{1}{3}$ mm. latis; ascis paraphysibus densis, filiformibus et longioribus circumdatis clavatis, longe et tenue stipitatis, 8-sporis, 90-150 μ . longis, parte sporifera, 90 \approx 14-18; sporidiis farctis, hyalinis, bi-conoideis, medio septatis constrictisque; loculis binis plerumque 2-nucleatis, 25 \approx 10.

Hab. ad ramos emortuos *Celtidis australis*, Vaucluse Galliæ.

24. **Didymella lophospora** Sacc. et Sp. Syll. I, p. 561 — * **echinophila** 2735 Sacc. Mich. II, p. 595. — A typo differt ostiolo paullo longiore; ascis angustioribus (spor. 14-18 \approx 4-5, appendiculatis; ascis 60-70 \approx 6-7).

Hab. in spinis involucrorum *Castaneæ*, Lyon Galliæ (THERRY).

25. **Didymella acerina** H. Fab. Sphér. Vaucl. II, p. 44, f. 13. — Pe- 2736 ritheciis sparsis superficialibus vel semiimmersis, lignum denudatum et fuligineo-inquinatum incolentibus, globulosis, $\frac{1}{2}$ mm. latis, vertice poro pertusis; ascis dense paraphysatis, cylindraceis, breve stipitatis, 8-sporis, 100-130 \approx 10-12; sporidiis oblique monostichis, hyalinis, medio septatis, constrictisque, altera dimidia parte conoidea, altera semiovata et crassiori; loculis binis multi-guttulatis, 18 \approx 7.

Hab. ad plagas ramorum resectorum *Aceris monspessulanæ*, in silvis Vaucluse Galliæ.

26. **Didymella Eurillæ** Pass. Fungi nuovi, IV, pag. 4, n. 8. — Peritheciis tectis, cortici interiori insculptis, globoso-depressis, ostiolo brevi acuto, contextu celluloso fuligineo; ascis clavatis, attenuato-

stipitatis, paraphysatis, 8-sporis, 80×15 ; sporidiis distichis, fusiformibus, utrinque acutis, medio septatis, plus minusve constrictis, hyalinis, loculis biguttulatis $18-20 \times 5-6$.

Hab. in ramulo sicco Juglandis, Vigheffio prope Parma (EURILLA PASSERINI).

27. **Didymella Salicis** Grove in litt. — Peritheciis peridermio infossis, sparsis vel binis ternisque adnatis vel gregariis, globulosis, dein subdepressis, atris, primo epidermide pustulatim tumida tectis, dein ea lacerata emergentibus et supra evanidâ nudis, papillatis, $\frac{1}{3}$ mm. diam.; ascis clavatis, interdum medio inflatis, vix infra attenuatis, subsessilibus, apice rotundatis, octosporis, $60-70 \times 11-13$; paraphysibus non visis; sporidiis distichis, oblongis, curvulis utrinque obtusis, granulosis ideoque levissime subluteolis, continuis, subinde 2-3-guttulatis, dein senio uniseptatis et medio subconstrictis, $16-20 \times 5$.

Hab. in ramis Salicis, Sutton Coldfield prope Birmingham Britanniæ.

28. **Didymella canadensis** Ell. et Ev. Proc. Phil. 1890, pag. 232. — Peritheciis inæqualiter sed crebre sparsis, cortice nidalibus eumque leviter elevantibus ostiolisque minutis papillatis nigris perforantibus, intus albis, globosis, $\frac{1}{3}$ mm. diam.; ascis tereti-clavatis, $75-90 \times 12-15$, copiose paraphysatis; sporidiis stipato-distichis, cylindraceis, obtusis, hyalinis, 4-nucleatis, medio distinctius leviterque prope utrumque apicem constrictis, $25-34 \times 6-7$.

Hab. in trunco emortuo Salicis, London Canada (J. DEARNES). — «An *Metaspheeria*?»

29. **Didymella Mali** Ell. et Ev. Proc. Phil. 1890, p. 232. — Peritheciis sparsis, $\frac{1}{4}$ mm. diam., cortice infossis, ostiolo acute conico tantum emergente; ascis clavato-cylindraceis, 70×7 , copiose paraphysatis; sporidiis biseriatis, fusoideis, leniter curvis, 4-nucleatis, hand constrictis, $20-22 \times 3$, utrinque acutis.

Hab. in cortice interiore relaxato Piri Mali viventis, Newfield N. J. Amer. bor. — «An *Zignoella* vel *Metaspheeria*?»

30. **Didymella hypophloea** Pass. Diagn. Fung. nuovi, I, n. 40. — Peritheciis sparsis, punctiformibus ovatis, fuscis, opacis, ostiolo acuto; ascis oblongo-clavatis, 75×10 , paraphysibus filiformibus æquantibus intermixtis; sporidiis oblique monostichis, fusiformibus, medio septatis, non constrictis, apicibus acutis, $15 \times 2,5$, hyalinis.

Hab. in cortice interiore Piri Mali, Vigheffio pr. Parma.

31. **Didymella Celastri** Brun. Champ. Saint. V, p. 2. — Peritheciis

approximatis, membranaceis, in cortice nidulantibus lignoque basi leviter insculptis, tectis, denique erumpentibus, globoso-depressis, perforatis, nigris, $\frac{1}{5}$ mm. diam.; ascis cylindraceo-clavatis v. sub-fusoideis, $60 \pm 8-10$, octosporis; sporidiis distichis vel oblique monostichis, oblongo-elongatis, utrinque rotundatis, hyalinis, 1-septatis, vix constrictis, 12 ± 4 .

Hab. in ramulis emortuis *Celastris scandentis*, Rochefort Gallæ.

32. **Didymella Opuntiæ** Pat. Bull. Soc. Myc. 1888, p. 112. — Peritheciis sparsis, majusculis, 1-1,5 mm. diam., nigris, globosis, poro pertusis, subcutaneis, dein superficialibus; ascis clavatis, sessilibus, $100-150 \pm 30-40$, crasse tunicatis, 8-sporis; sporidiis distichis utrinque obtusis, medio septatis et profunde constrictis, hyalinis, granulosis, loculis subconoideis.

Hab. in foliis putridis *Opuntiæ*, Caracas Venezuelæ.

33. **Didymella nigrificans** Karst. in Hedw. 1884, n. I, p. 3. — Rev. 2744 atque Synop. Ascom. fenn. pag. 60. — Peritheciis densissime congestis, cuticula nigrificata, demum fissa tectis, subsphæroideis, basi cupulato-collapsis, pertusis, atris, glabris, circ. 2 mm. diam.; ascis cylindraceis, stipitatis, pseudo-paraphysatis, $70-100 \pm 6$; sporidiis octonis, monostichis, oblongatis vel ellipsoideis, 1-rarissime 3-septatis, eguttulatis, chlorino-hyalinis, nitido strato mucoso obvolutis, $11-15 \pm 4-5$.

Hab. in ramis emortuis *Rosarum* in Mustiala Fenniæ.

34. **Didymella Rauii** (Ell. et Ev) Berl. et Vogl. Add. Syll. p. 86, *Sphæria* Ell. et Ev. in Torr. Bot. Club. 1883, p. 90. — Peritheciis sparsis v. 2-3 in cortice interiore coalitis, epidermide laxe tectis, demum eam ostiolo papillæformi perforantibus; ascis $30-35 \pm 6-7$; paraphysibus gracilibus; sporidiis distichis, oblongo-fusoideis, 1-septato-constrictis, leniter curvulis, utrinque appendicula minuta præditis, luteolis, binucleatis, $7-8 \pm 5$.

Hab. in ramis emortuis *Rosæ* cultæ, Bethlehem Amer bor. (E. A. RAU).

35. **Didymella eriostoma** Sacc. in Mout. Ascom. Lièg. II, p. 4, t. I, 2746 f. 17. — Peritheciis sparsis, atris, globoso-depressis, $\frac{1}{3}$ mm. diam., tectis, ostiolo papillato, fasciculo setularum brevium coronato; ascis tereti-clavatis, copiose filiformi-paraphysatis, 8-sporis, $90-100 \pm 12$; sporidiis mono- vel subdistichis, oblongo-fusoideis, vulgo rectis, medio septatis, leniter constrictis, 4-guttatis, $20-22 \pm 6-7$, hyalinis, utrinque appendicula brevi 2μ . long. auctis.

Hab. in ramulis Sarothamni scoparii, Gomzé, Tilff etc.
Belgii. — *D. Genistæ* affinis.

36. **Didymella buxicola** H. Fab. Sphér. Vaucl. II, pag. 44, fig. 15 — 2747

Peritheciis dense gregariis, cortici infossis, nonnullis autem hinc illinc fere superficialibus, globosis, $\frac{1}{3}$ mm. latis; ostiolis crassis, breve cylindraceis aut conicis, inæqualibus, corticem immutatum v. cinereo-infuscatum exasperantibus; ascis paraphysibus densis circumdatis, 8-sporis, plerisque clavatis, breve aut longissime stipitatis, parte sporifera 90 \approx 12, connallis autem cylindraceis 120 \approx 9; sporidiis in ascis clavatis distichis, in cylindraceis submonostichis oblongis, hyalinis, utrinque obtusis, medio constrictis et subtile septatis, 4-magni-guttatis; altera diuidia parte perpaullo crassiori 20-25 \approx 6.

Hab. ad caudices et ramos emortuos Buxi sempervirentis, Vaucluse Galliæ.

37. **Didymella Olearum** H. Fab. Sphér. Vaucl. II, p. 43, fig. 11. — 2748

Peritheciis rugis matricis prominentibus et atro-inquinatis infossis, nunc solitariis nunc seriatim gregariis, globosis, $\frac{1}{2}$ mm. latis, matricem in pustulas cinereo-atras, ad latera compressas, 1 mm. et ultra longas inflantibus; ostiolo brevi, papilliformi, eximie nitido; ascis dense paraphysatis, brevissime stipitatis, cylindraceis, 8-sporis, 120-150 \approx 6-7.

Hab. ad truncos emortuos Oleæ europææ, in ligno denudato, Avignon Galliæ (Th. DELACOUR.)

38. **Didymella maculiformis** Winter in Hedw. 1884, pag. 169. — Pe. 2749

ritheciis amphigenis, densissime stipatis, greges maculiformes, irregulares, magnos, saepe confluentes, usque 6 cm. latos formantibus, immersis, punctiformibus, e basi globosa, brevissime conicis, poro simplici pertusis, epidermide circa verticem vix prominentem nigrefacta tectis; sporidiis distichis, cylindraceis, inæqualiter didymis, ad septum parum constrictis hyalinis, 26-30 \approx 5-6; ascis oblongis, brevissime stipitatis, sursum parum attenuatis, 8-sporis, 78-82 \approx 16-17, paraphysibus conglutinatis.

Hab. in foliis emortuis Proteæ grandifloræ prope Cape-town in Promontorio Bonæ Spei.

39. **Didymella truncata** Karst. Rev. Myc. 1889. — Peritheciis di-

spersis, erumpenti-superficialibus, conoideo-truncatis, poro latiusculo, atris, 0,2-0,3 mm. diam.; ascis cylindraceo-clavatis, 130-150 \approx 12-14, 8-sporis; sporidiis distichis, ovoideo-elongatis, inæquilate-

670 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Didymella.
ralibus, 1-septatis, ad septum leviter constrictis, subhyalinis, 26-
33 \approx 5-7; paraphysibus numerosis, ramosis, 0,5 μ . cr.
Hab. ad virgulta in Minas Geraes Brasiliæ (EDW. WAINIO).

III. Monocotyledonicolæ.

40. **Didymella? Carphæ** Speg. F. Fueg. n. 205. — Maculis nullis; 2751
peritheciis sparsis v. laxissime gregariis globoso-lenticularibus, 130-
150 μ . d., epidermide velatis, atris, glabris, membranaceo-coriacellis,
contextu densiuscule parenchymatico fuligineo, ostiolo vix papillu-
lato obtuso; ascis cylindraceis, apice truncato-rotundatis crassiu-
scule tunicatis, basi breviter attenuato-pedicellatis, 55-60 \approx 6-7,
paraphysibus crassiusculis guttulatis immixtis, octosporis; sporidiis
elliptico-ovoideis, v. subbiconicis, utrinque obtusiusculis, 10 \approx 4,5-5,
primo continuis hyalinis, dein diblastis pallide fuscescentibus.

Hab. ad scapos vaginasque Carphæ schœnoidis in Staten
island Fuegiæ.

41. **Didymella glacialis** Rehm Beitr. Ascom. Hedw. 1882, p. 121. — 2752
Peritheciis minutis, globosis, atris parenchymate foliorum denigra-
to immersis ejusque superficiem papilla conoidea atra vix superau-
tibus, gregariis; ascis clavatis, 8-sporis, 50 \approx 9; sporidiis ellipticis,
subacutatis, inæquilateralibus, 2-cellularibus, utraque cellula ma-
gno nucleo instructa, hyalinis, distichis, 12 \approx 3; paraphysibus filiformibus.

Hab. ad folia graminea sicca prope Taschach-Gleßcher in
valle Piz alpium Tirolensis, alt. c. 2000 metr. (DR. REHM).

42. **Didymella arctica** (Fuck.) Berl. et Vogl. Add. Syll. p. 88, *Sphæria* 2753
Fuck. Die 2^e Deutsche Nordpolarf. II, 94, et Oudem. Contr. Fl. Myc.
Now. Seml. p. 152, tab. III, fig. 4. — Peritheciis in macula cine-
rescente gregariis, seu linear-dispositis, erumpentibus, globoso-co-
nicis, minutis, aterrimis, demum perforatis; ascis fasciculatis, ob-
longis, contortis, sessilibus, tunica crassa præditis, 72 \approx 17, octo-
sporis; sporidiis imbricato-distichis, oblongis, utrinque parum at-
tenuatis, rectis, uniseptatis, ad septum constrictis, loculis inæqua-
libus, biguttulatis, 24 \approx 8, hyalinis.

Hab. in foliis exsiccatis Poæ alpinæ, Nowaja Semlja.

43. **Didymella pachyspora** Bomm. Rouss. Sacc. sp. nov. — Perithe- 2754
ciis sparsis, globulosis, glabris, initio velatis, dein erumpentibus,
intus griseolis, poro pertusis; ascis oblongis, sessilibus, 60-75 \approx 15,
octosporis, obsolete paraphysatis; sporidiis late ellipsoideis, 14-
18 \approx 9, mono- distichis, 1-septatis, non constrictis.

Hab. in foliis Festucæ glaucæ, Comblain au pont Belgii.

IV. Acotyledonicolæ.

44. **Didymella Chamæcyparissi** Rehm in Voss Mat. Pilzk. IV, p. 10, 2755
 fig. 5. — Peritheciis globosis, immersis, gregariis, minutis, apice
 haud perspicue pertusis, contextu parenchymatico; ascis clavatis,
 6-8-sporis, 60 \times 12; sporidiis distichis, hyalinis, 1-septatis, subcon-
 strictis, 12-15 \times 5, cellula superiore latiore, 4-nucleatis; paraphy-
 sibus filiformibus, 3 μ . cr.

Hab. in foliis emortuis Lycopodii Chamæcyparissi pr.
 Laibach Carnioliae.

45. **Didymella Ulothii** (Körber) Berl. et Vogl. Add. Syll. p. 89, *Didymo-* 2756
sphaeria Winter in Hedw. 1886, p. 26, *Cercidospora Ulothii* Kör-
 ber, Parerga p. 466. — Peritheciis minutissimis, sparsis v. subgre-
 gariis, thallo omnino immersis, globosis, vertice tantum pertuso, de-
 dum depresso, parum prominulis, atris; ascis cylindricis, deorsum
 parum attenuatis, 4-8-sporis, 70-80 \times 10; sporidiis oblique mono-
 stichis, fusoideis, utrinque acutis, bicellularibus, hyalinis, 16-20 \times 5;
 paraphysibus numerosis, filiformibus.

Hab. in thallo apotheciisque Placodii saxicoli in Ger-
 mania.

46. **Didymella epipolytropa** (Mudd.) Berl. et Vogl. Add. Syll. p. 89, 2757
Didymosphaeria Winter in Hedw. 1886, p. 26, *Thelidium epipo-*
lytropum Mudd. Manual of Brit. Lichen. pag. 298, *Cercidospora*
epipolytropa Arnold in Flora 1874, pag. 154. — Peritheciis spar-
 sis, punctiformibus, minutissimis, immersis vel semiemersis, globo-
 sis, atris, sub microscopio viridi-atris, vertice subdepresso et perfo-
 rato; ascis breve cylindraceis v. oblongo-clavatis, 4-8-sporis; pa-
 raphysibus distinctis, tenuibus, filiformibus, subramosis; sporidiis
 elongato-oblongis, fere fusiformibus, bicellularibus, hyalinis, 15-
 18 \times 4-5.

Hab. in thallo discoque apotheciorum Lecanoræ polytro-
 pæ in Germania.

47. **Didymella sphinctrinoides** (Zwackh) Berl. et Vogl. Add. Syll. p. 89, 2758
Didymosphaeria Winter in Hedwigia 1886, pag. 26, *Endococ-*
cus sphinctrinoides Zwackh in Flora 1864, pag. 88. — Perithe-
 ciis.... indescriptis; ascis cylindraceis, angustis, 6-8-sporis, 74-78
 \times 10-12; sporidiis bicellularibus, hyalinis, 18-22 \times 6-7; paraphysi-
 bus filiformibus.

Hab. parasitice in Lecanora subfusca in Germania. — Var.

Immersæ (Arnold) Berl. et Vogl. l. c., *Endococcus sphinctrinoides* var. *Immersæ* Arn. Lichen. Fragm. XVI, p. 25, t. II, f. 14 in Flora 1874, *Didymosphaeria sphinctrinoides* var. *Immersæ* Winter in Hedw. 1886, p. 26: Peritheciis immersis, semiemersis, minutis, globosis, atris; sporidiis oblongis, bicellularibus, hyalinis, 18 ± 6 ; ascis cylindraceis, octosporis; paraphysibus distinctis, capillaribus. In thallo Lecidellæ immersæ in Germania.

HARIOTIA Karst. Journ. Bot. 1889, p. 206 (Etym. a cl. botanico P. HARIOT). — Perithecia carbonacea, erumpentia, glabra, rimâ subdehiscentia, atra. Ascii cylindraceo-clavati, polyspori. Sporidia (24) ovoidea, 1-septata, hyalina. Paraphyses nullæ.

1. **Hariotia strobiligena** (Desm.) Karst. l. c., *Sphaeria* Desm. XIII, 2759 Not. p. 75, *Didymella* Sacc. Syll. I, p. 552. — Ascii 105-120 \approx 12-15, jodo non cœrulecentes; sporidia disticha 12-18 \approx 4-6.

Hab. in conis dejectis Pinî maritimæ in Gallia occident. — *Dichænam strobilinam* in memoriam revocat.

GNOMONIA Ces. et De Not. — Sacc. Syll. I, p. 501.

1. **Gnomonia euphorbiacea** Sacc. et Briard in Rev. Myc. 1885. — 2760 Peritheciis sparsis, epidermide tectis, sphæroideis, basi cupuliformi vel excavata, $1/5-1/4$ mm. diam.; ostiolis punctiformibus, epidermidem perforantibus eique adhærentibus; ascis oblongis, subfusiformibus, breviter stipitatis, octosporis, 72-88 \approx 12; sporidiis monostichis vel centro distichis, ovato-oblongis, obtusis, 1-septato-constrictis, biguttulatis, rectis vel leniter curvulis, 14-16 \approx 6, hyalinis.

Hab. in ramlulis emortuis Euphorbiæ palustris, Troyes Galliae (BRIARD).

2. **Gnomonia tithymalina** Sacc. et Briard in Rev. Myc. 1885. — 2761 Peritheciis numerosis, plus minusve gregariis, sub epidermide nidulantibus, sphæroideis, basi cupulata v. excavata, $1/3-1/2$ mm. diam.; ostiolis obtusis, epidermidem leniter nigrefactam perforantibus; ascis cylindraceo-fusiformibus, sessilibus, 40-44 \approx 8-10, octosporis; sporidiis oblique distichis, oblongis, obtusis, 1-septato-constrictis, 4-guttulatis, rectis v. vix curvulis, 13-15 \approx 4-5, hyalinis.

Hab. in caulis emortuis Euphorbiæ palustris, Troyes (BRIARD).

3. **Gnomonia Amygdalinæ** Fuck., Sacc. Myc. Sibir. p. 12, f. 8, *Gnomoniella* Sacc. Syll. Fung. I, p. 418. — Perithecia lageniformia

obtuse rostellata, $\frac{1}{4}$ mm. lat.; ascis 36-40 \approx 6-7; spor. subfusoidea, 16-18 \approx 3-3,5, matura distincte 1-septata; hinc vera *Gnomonia* nec *Gnomoniella*. — Viget quoque in caulis emortuis Euphorbiæ lutescentis in campis pr. Meticowo Sibiriæ.

4. ***Gnomonia Gei*** Pat. et Doass. Tab. An. Fung. Fasc. IV, pag. 214, 2762 fig. 495. — Peritheciis sparsis, epidermide tectis, ostiolo filiformi, valde elongato præditis; ascis brevibus, sessilibus, 26 \approx 5, sursum truncatis; paraphysibus nullis; sporidiis bacillaribus, acicularibus, 1-septatis, septo vix constrictis, hyalinis, 12-14 \approx 1.

Hab. in caulis Gei pyrenaici, Les Eaux Bonnes Galliæ.

5. ***Gnomonia borealis*** Schroet. Nord. Pilze p. 11. — Peritheciis late 2763 laxe gregariis, epidermide pustulata tectis, 0,6 mm. lat., globulosis, collapsis, nigris, ostiolo subulato, 0,2 mm. long., erumpente; ascis copiosis, fusoideis, sessilibus, 44-50 \approx 10-12, 8-sporis; sporidiis distichis, elliptico-fusoideis, 15-17 \approx 4-5, utrinque rotundatis, medio 1-septatis leniterque constrictis, 2-4-guttulatis, hyalinis.

Hab. in caulis vetustis Geranii silvatici, Magerö, Nordcap Norvegiæ.

6. ***Gnomonia carpophila*** Bomm. Rouss. Sacc. sp. nov. — Peritheciis 2764 sparsis, subseriatis vel 4-5 consociatis, subepidermicis, dein erumpentibus, glabris, conoideis, 300-360 μ . diam., ostiolo perithecium superante, tereti-papillato, sæpe curvulo; ascis clavatis; sporidiis distichis, oblongo-fusoideis, 1-septatis, loculis subinæqualibus, pluriguttulatis, hyalinis, 18 \approx 3,5.

Hab. in capsulis *Oenotheræ bienniæ*, Groenendael Belgii.

7. ***Gnomonia Rubi*** Rehm Ascom. n. 596, Wint. in Hedw. 1887, p. 62. 2765 — *G. tetraspora* **Rubi* Rehm — Syll. I, p. 562. — Peritheciis sparsis, immersis, globosis, rostro crasso cylindrico, longo ornatis, 180-200 μ . diam.; ascis subfusoideis, sessilibus, 45-52 \approx 8-9, tetrasporis; sporidiis subdistichis, inæqualiter fusoideis, utrinque breve appendiculatis, 1-septatis non vel vix constrictis, 4-guttulatis, 14-16 \approx 3,5.

Hab. in foliis dejectis Rubi fruticosi, Zürich Helvetiæ (WINTER).

8. ***Gnomonia spermogoniooides*** Rehm Ascom. Ill. n. 944, Sydow. Myc. 2766 march. 1355 (sub *Ceratosphaeria*). — Peritheciis gregariis, in cortice interiore nidulantibus, globosis, viridifuscis vel fuscis, 0,2-0,3 mm. lat., collo filiformi, 0,3 mm. longo, peridermum perforante et prominente, demum peridermio lacerato denudatis; ascis teneris, ellipticis, 36 \approx 9, 4-8-sporis; sporidiis bacillaribus, obtusis, rectis

- 674 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Gnomonia.
v. subcurvatis, 2-cellularibus, utraque cellula binucleata, hyalinis, 12-15 \approx 2,5, distichis; paraphysibus filiformibus, nucleatis, longis, c. 3 μ . lat.
Hab. ad sarmenta putrescentia Rubi agrestis prope Berolini (SYDOW).
9. **Gnomonia Magnoliæ** Ell. Am. Nat. 1883, pag. 318. — Peritheciis 2767 majusculis foliorum parenchymate immersis, rufis, subulato-conicis, ostiolo tantum visibili; ascis oblongo-ellipsoideis, 40 \approx 7-8; sporidiis fusoideis, acutis, pallide stramineis, obscure nucleatis, 11-19 \approx 2.
Hab. in foliis dejectis Magnoliæ glaucæ, Newfield N. J. Amer. bor. — «An potius *Gnomoniella*?».
10. **Gnomonia acerina** Starb. Skand. Pyr. p. 17, t. I, f. 6. — Peritheciis 2768 hypophyllis, sparsis, minutis, 250 μ . diam., siccis hemisphaerico-prominulis, ostiolo gracili, brevi; ascis ellipsoideo-cylindraceis, 65-75 \approx 12-18, sessilibus; sporidiis tereti-fusoideis, prope basin 1-septatis, loculo sup. 2-3-guttato, infero 1-guttato, curvulis, distichis, 18 \approx 6.
Hab. in foliis siccis Aceris platanoidis in horto botanico Upsaliensi Sueciæ. — *G. errabundæ* subaffinis.
11. **Gnomonia petiolophila** (Peck) Berl. et Vogl. Add. Syll. pag. 90, 2769 *Sphæria petiolophila* Peck in 35 Rep. St. Mus. Bot. 1884, p. 144. — Peritheciis minutis, sparsis primum epidermide tectis, dein, ea perfossa, prominulis, ostiolo papillato v. breviter rostrato instructis, compresso-globosis, nigris; ascis subcylindraceis, angustis 40-45 μ . long.; sporidiis anguste fusiformibus, utrinque acutis, distichis, 12-15 \approx 2, quandoque 3-4-nucleolatis.
Hab. in petiolis emortuis Aceris spicati in montibus Helderberg Amer. bor.
12. **Gnomonia australis** Winter Contr. Fl. Myc. Lus. V, p. 14. — 2770 Peritheciis dense sparsis, hypophyllis, immersis, demum erumpentibus, globosis, membranaceis, fusco-atris, 120-200 μ . diam., rostro crasso, cylindrico, apice sæpe parum compresso et incrassato, usque 180 \approx 20, instructis; ascis fusoideo-elongatis, 8-sporis, 56-60 \approx 8-9, sporidiis oblongis, utrinque parum attenuatis et rotundatis, inæquilateralibus, bicellularibus, 4-guttulatis, hyalinis, ad septum constrictis, 14-16 \approx 4-4,3.
Hab. ad Apolloniæ canariensis folia arida, Coimbra Lusitaniæ (MOLLER). — Var. **Lauri** Wint. l. c.: Peritheciis gregariis in macula indeterminata, obscuriore hypophylla; ascis deorsum in pedicellum brevem attenuatis, usque 78 μ . longis; sporidiis inter-

dum usque 18 μ . longis. Ad folia arida Lauri nobilis prope Coimbra Lusitaniæ.

13. **Gnomonia Sassafras** Ell. et Ev. in Torr. bot. Club. 1883, p. 98. 2771

— Peritheciis erumpentibus, hemisphæricis, circiter 200 μ . diam., in superficie inferiore foliorum sparsis; ostiolis filiformibus 200-250 μ . longis, contextu fibroso, sursum subhyalino, sæpe parum curvatis; ascis lanceolatis, curvis, apophysatis; sporidiis filiformibus, dilutissime flavidis, indistincte multinucleatis, 35-40 \approx 1,7.

Hab. in foliis emortuis Sassafras officinalis, Fairfield, Ohio Amer. bor. (W. A. KELLERMANN). «An *Linospora*?».

14. **Gnomonia perversa** Rehm Ascom. Ill. n. 795. — Peritheciis mi- 2772

nutissimis, sparsis, parenchymati foliorum immersis, nudo oculo vix punctiformiter paginam superiorem foliorum inquinantibus conspicuis eamque non elevantibus, haud rostratis, vertice punctiformi, atris, contextu parenchymatico fusco; ascis fusiformibus, in stipitem tenerum elongatis, 8-sporis, 60 \approx 7; sporidiis 2-3-stichis, ellipticis, rectis vel interdum subcurvatis, haud acutatis, 2-cellularibus, utraque cellula 2-nucleata, hyalina, 12 \approx 3,5; paraphysibus nullis.

Hab. ad folia putrida Alni glutinosæ prope Königstein a. Elbe Saxoniæ (W. KRIEGER). — An potius *Hypospilæ* species? *H. Rehmii* Sacc. proxima.

15. **Gnomonia coriacea** Cooke et Mass. Grev. XVIII, p. 35. — Pe- 2773

ritheciis minutis, in maculas orbiculares congestis, foliorum parenchymati innatis, ostiolo elongato, sursum leniter attenuato; ascis clavato-stipitatis, octosporis; sporidiis uniseptatis, obtusis, hyalinis, 10 \approx 2-3.

Hab. in foliis coriaceis in Brasilia (GLAZIUS).

16. **Gnomonia? magellanica** Speg. F. Pat. n. 127. — Maculis nullis; 2774

peritheciis parenchymate innatis, tectis, amphigenis, sparsis, minutis, globosis, 100-250 μ . diam., glabris, ostiolo setiformi longiusculo exerto, elongatulo, levi, concolori; ascis et sporidiis ignotis.

Hab. ad folia dejecta putrescentia Fagi antarcticæ in silvis Voces bay Patagoniæ.

17. **Gnomonia Rhois** Rich. Cat. Champ. Marn. n. 1231. — Perithe- 2775

ciis subcutaneis, ostiolis spiniformibus erumpentibus; ascis basi attenuatis, apophysatis; sporidiis anguste fusiformibus.

Hab. in petiolis foliorum Rhois typhini, Saint-Amand Galliæ.

Doct. H. REHM medico). — Perithecia rostrata, membranacea, immersa. Asci polyspori, aparaphysati. Sporidia didyma, hyalina. — A genere *Gnomonia* ascis polysporis in primis differt.

1. **Rehmiella alpina** Winter l. c. pag. 2. — Peritheciis sparsis immersis, globosis, ca. $\frac{1}{2}$ mm. diam., glabris, atris, rostro elongato, cylindraceo, saepe curvato usque $\frac{1}{2}$ mm. longo, erumpentibus; ascis oblongatis, sessilibus, polysporis, 50-70 \approx 10-11; sporidiis inordinatis, oblongo-cylindraceis, infra medium uniseptatis et parum constrictis, 4-guttulatis, hyalinis, 10-12,5 \approx 2.

Hab. ad petiolos foliaque putrida Alchemillæ alpinæ, Rigi prope Luzern Helvetiæ (G. WINTER).

PHARCIDIA Körber Parerga Lichenol. p. 469 (Etym. *pharcis* ruga). — Perithecia sparsa vel laxe gregaria, semiimmersa vel demum emergentia, atra, punctiformia, minuta, globosa vel globoso-conoidea, poro pertusa. Asci 4-8-spori, oblongi v. fusoideo-clavati. Sporidia cuneata, ovoidea vel fusiformia, 1-3-septata, hyalina.

1. **Pharcidia Schœreri** (Mass.) Winter in Hedwigia 1886, pag. 11. *Epicymatia Schœreri* (Mass.) Sacc. Sylloge fungorum vol. I, pag. 571. — Perithecia sparsa, primo plus minusve immersa, demum emergentia, globosa, verticem versus parum attenuata, poro pertusa, membranacea, fusca, minutissima, punctiformia; ascis oblongi, utrinque subattenuati, 8-spori, 30-40 \approx 12; sporidia conglobata, oblonga, subclavata v. bacillaria, 1-2-septata, hyalina, 12-14 \approx 3-3,5; paraphyses mucosæ, indistinctæ. — Var. **Croceæ** Arnold (in *Solorina crocea*); sporidiis 12 \approx 2-3.

2. **Pharcidia parvipuncta** (Stein) Wint. in Hedw. 1886, p. 14, *Sagedia parvipuncta* Stein Flechten in Krypt. Flora von Schlesien II, 2, pag. 339. — Peritheciis minutissimis, sessilibus, hemisphaericis v. conoideis, atris, nitidis, molliusculis, 30 μ . diam.; ascis subfusiformi-clavatis, membrana crassa præditis, 8-sporis; sporidiis fusiformibus, abrupte acutatis, 2-4-cellularibus, 12-18 \approx 4-5; paraphysibus mucosis, parum distinctis.

Hab. in thallo Thelidii diaboli in Silesia Germania.

3. **Pharcidia constrictella** Müller in Flora 1874, p. 350 et Wint. in Hedw. 1886, pag. 12. — Peritheciis fere totis immersis, globosis, sursum parum attenuatis, thalli superficie a vertice tantum pertusis, $\frac{1}{5}$ mm. latis, tenuissime membranaceis, fusco-atris; ascis cylindricis, obtusis 4-sporis, circ. 75 \approx 12; sporidiis monostichis, el-

liptico-oblongis, utrinque rotundatis, medio distincte constrictis, 18-22 \approx 8-9.

Hab. in thallo *Psoromatis fulgentis* var. *alpinæ* in Germania.

4. **Pharcidia Arthoniæ** (Arnold) Wint. in Hedw. 1886, p. 12, *Sphaeria Arthoniæ* Arnold Lichenol. Ausfl. VIII, p. 304. — Peritheciis sparsis, fere omnino emersis, minutissimis, globosis, membranaceis fuscis; ascis e basi latiori sursum oblongis, sessilibus, 8-sporis, 40-45 \approx 19; sporidiis congregatis, cylindricis, utrinque rotundatis, 1-3-septatis, hyalinis, rectis, 18-23 \approx 5; paraphysibus nullis.

Hab. parasitica in thallo *Arthoniæ astroideæ* in Germania.

5. **Pharcidia dispersa** (Lahm) Wint. in Hedw. 1886, pag. 13, *Arthopyrenia dispersa* Lahm in Körber Parerga pag. 388. — Peritheciis sparsis, minutissimis, globosis vel hemisphæricis, atris opacis, ostiolo parum distincto, subverruciformi; sporidiis octonis in ascis cylindrico-clavatis, breviter stipitatis, subsoleæformibus, bicellularibus, plerumque cellula superiori latiori, hyalinis, 13-15 \approx 4-5.

Hab. parasitica in thallo *Biatoræ rupestris*, *Callopismatis ferruginei* et *Lecanoræ subfuscæ* in Germania.

6. **Pharcidia Lichenum** Arnold Lichenol. Ausfl. in Tirol. VIII, p. 302 2781 et Wint. in Hedw. 1886, p. 11. — Peritheciis sparsis vel gregariis, plus minusve immersis vel demum emergentibus, immo interdum fere superficialibus, globosis, poro simplici pertusis, minutissimis, membranaceis, fuscis; ascis oblongis, ventricosis utrinque parum attenuatis, substipitatis, 8-sporis, 33-42 \approx 12-14; sporidiis congregatis, clavatis vel cuneatis, vertice rotundatis, bicellularibus, non constrictis, hyalinis, 12-15 \approx 3,5-4; paraphysibus mucosis.

Hab. parasitica in thallo, rarius in fructibus *Lichenum variorum* in Germania.

7. **Pharcidia hygrophila** (Arnold) Wint. in Hedw. 1886, p. 14, *Endococcus hygrophilus* Arnold in Flora 1871, p. 147. — Peritheciis atris, subglobosis, supra thalli granulos dispersis, minimis; ascis elongato-ellipticis, sursum parum, deorsum magis stipitiformi-attenuatis; sporidiis elliptico-oblongis, utrinque attenuatis vel in uno apice late rotundatis, saepe inæquilateralibus, 1-septatis, 15-18 \approx 5-6, octonis, hyalinis, non constrictis.

Hab. in thallo *Buelliae rivularis* in Germania — Var. **Berengeriana** Arnold Lichenol. Ausfl. VI, pag. 1146: Peritheciis sat minutis, atris, globoso-emersis, dispersis; paraphysibus capil-

laribus; sporidiis 1-septatis, cum 2-4 guttulis oleosis, utroque apice plus minus cuspidatis, 22-28 \approx 5-6, 8 in ascis oblongis. In thallo Biatoræ Berengerianæ in Germania.

8. **Pharcidia Porocyphi** (Stein) Wint. in Hedw. 1886, p. 12, *Artho-* 2783
pyrenia Porocyphi Stein Flechten in Krypt. Flora von Schlesien II, 2, pag. 343. — Peritheciis minutissimis, sessilibus, conoideis vel globosis, sine ostiolo distincto, atris, nitidis, 30 μ . diam.; ascis subfusiformibus, 8-sporis; sporidiis bicellularibus, ovoideo-ellipsoideis, cellula superiori plerumque parum majori, 6-8 \approx 2-3.

Hab. in thallo Porocyphi Cataractarum in Silesia Germaniæ.

9. **Pharcidia microspila** (Körber) Wint. in Hedw. 1886, p. 13, *Artho-* 2784
pyrenia microspila Körber Parerga pag. 392. — Peritheciis minutissimis, primo immersis, demum emergentibus, tandem superficialibus, subglobosis, ostiolo papilliformi indistincto, atris, opacis, in maculis rotundatis irregularibusque, circ. 2-3 mm. latis, cinereo-atris, e mycelio fusco, breve articulato formatis, gregariis; ascis oblongis vel elongato-oblongis, medio subventricosis, fere sessilibus, 8-sporis, 42-50 \approx 14; sporidiis inordinatis, cylindrico-cuneatis, 2 (rarissime 4)-cellularibus, haud constrictis, hyalinis, 15-17 \approx 4-5.

Hab. parasitica in thallo Graphidis scriptæ in Germania.

10. **Pharcidia Aspiciliæ** (Lahm) Wint. in Hedw. 1886, p. 13, *Artho-* 2785
pyrenia Aspiciliæ Lahm in Körber Parerga p. 388. — Peritheciis confertis, minutissimis, hemisphaerico-conoideis, atris, subnitidis, ostiolo indistincto, immersis, demum erumpentibus; sporidiis octonis in ascis fusiformi-clavatis, oblongis, bicellularibus, demum guttulis 4 præditis (4-cellularibus?), diam. 3-4 plo longioribus, hyalinis.

Hab. in thallo Aspiciliæ calcareæ in Germania.

11. **Pharcidia Punctillum** (Arnold) Wint. in Hedw. 1886, pag. 12, *Ar-* 2786
thopyrenia Punctilum Arnold Lichenol. Ausfl. XI, pag. 22. — Peritheciis sparsis, minutissimis, punctiformibus, hemisphaerico-emersis, membranaceis, atro-fuscis; ascis e basi latiori, ventricosa sursum attenuatis, 8-sporis, 30-40 \approx 14-16; sporidiis conglobatis, oblongo-clavatis, bicellularibus, hyalinis, medio haud vel perparum constrictis, utrinque rotundatis, 14-18 \approx 3,5-5.

Hab. in thallo Biatorinæ cyrtellæ in Germania.

12. **Pharcidia conspurcans** (Fries) Wint. in Hedw. 1886, p. 13, *Ar-* 2787
thopyrenia conspurcans Fries Lichen. Spitsb. p. 51. — Perithe-

ciis punctiformibus, semiemersis, atris; hymenio absque paraphysibus; sporidiis bicellularibus, incoloribus, utroque apice plus minus obtusis, sæpe cum 2 guttulis oleosis, 10-14 \approx 5, 8 in ascis ventricosis.

Hab. parasitica in thallo Dimelænæ nimbosæ in Germania.

13. **Pharcidia Badiæ** (Arnold) Winter in Hedw. 1886, p. 12, *Arthopyrenia Badiæ* Arnold Lichenol. Ausf. in Tirol. X, p. 13. — Peritheciis majusculis, hemisphæricis, emersis, non pertusis, atris; sporidiis elongatis, bicellularibus, hinc inde cum 4 guttulis oleosis, hyalinis, 15-16 \approx 4-5, 8 in asco; paraphysibus nullis.

Hab. in thallo Lecanoræ badiæ in Germania.

14. **Pharcidia Tabacinæ** Arnold in Flora 1881, p. 176 et Wint. in Hedw. 1886, pag. 12. — Peritheciis punctiformibus, semiemersis, atris, hymenio absque paraphysibus; sporidiis incoloribus, elongato-oblongis, 1-septatis, regulariter 4-guttulatis, 10-12 \approx 3, 8 in asco, medio inflatis.

Hab. in thallo Psoræ tabacinæ in Germania.

15. **Pharcidia Ahlesiana** (Hepp) Wint. in Hedw. 1886, pag. 14. *Sagedia Ahlesiana* Hepp in Zwackh Exsicc. 314, *Endococcus Ahlesianus* Zwackh in Flora 1862, pag. 572. — Peritheciis et ascis ignotis; sporidiis octonis, bicellularibus, interdum subcurvatis, semper incoloribus, utrinque subobtusis, 9-14 \approx 3-4.

Hab. in Parmelia demissa parasitica in Germania.

16. **Pharcidia cerinaria** (Mudd) Sacc. *Sphæria cerinaria* Mudd British Lichens p. 136. — Peritheciis perexiguis, punctiformibus, semiimmersis, nigris, solitariis, 10-12 in quoque disco apotheciorum insidentibus; ascis...?; sporidiis numerosis, oblongis, subconstrictis, bilocularibus, 3,7 \approx 1,8, pallide flavis; paraphysibus nullis.

Hab. in apotheciis Lecanoræ cerinæ pr. Ayton York Brit.

LIZONIA Ces. et De Not. — Sacc. Syll. I, p. 574.

1. **Lizonia bertiodes** Sacc. et Berl. in Rev. Myc. 1885, tab. LIV, fig. 5. — Peritheciis 2-5 acervulatis, stromate plano, rufescente insidentibus, sphæroideis, rugosiusculis, atris; ascis amplis, ovatis, crasse tunicatis, stipitatis, paraphysibus longioribus cinctis, 50-60 \approx 20-25, octosporis; sporidiis inordinate tristichis, oblongo-ovoideis, guttuligeris, demum didymis, 20 \approx 8, pallide luteis, dein fuscellis.

Hab. in pagina superiore foliorum Compositæ corymbiferæ

cujusdam, Santos Brasiliæ. — Sociis *Hypocreacea* quadam scle-cospora (asci 70 ± 14; sp. 50-55 ± 3), *Polystigmina* (sp. 40-50 ± 2 utrinque hamatæ), *Entomosporio*, quorum conceptacula omnino la-tent, et *Helotio* amœne aureo, immaturo.

2. **Lizonia abscondita** Johans. Svamp. fran Island. p. 167, t. XXIX, 2793 f. 9. — Peritheciis superficialibus, gregariis vel subsparsis in ner-vis immersis foliorum insidentibus, subcoriaceis, atris, glabris, filis fuligineis a basi prodeuntibus, plerumque ovoideis, minutis, 130-140 ± 88-100; ascis cylindraceo-clavatis, vix vel brevissime pedi-cellatis, octosporis, paraphysibus nullis, 54-70 ± 16-20; sporidiis distichis, pallide olivaceis, ovoideis, medio vel prope medio septa-tis et constrictis, membrana tenuissima, 12-17 ± 7-10.

Hab. in pag. superiore foliorum exsiccatorum Dryadis octo-petalæ, Eskifjördur Islandiæ orient.

3. **Lizonia halophila** Bomm. Rouss. Sacc. sp. nov. — Peritheciis am-phigenis, dense congestis, superficialibus, ovoideis, glabris, 60-115 μ. diam., 90-110 μ. alt., poro pertusis; ascis saccatis vel obovatis, sessilibus, aparaphysatis, 48-54 ± 18-21, 8-sporis; sporidiis 2-sti-chis vel congregatis, oblongis, uno apice tenuatis, 1-septatis, non constrictis, 21-24 ± 6-6,5, hyalinis.

Hab. in foliis emortuis Honckeneyæ (Arenariæ) peploidis, Nieuport Belgii.

4. **Lizonia? inæqualis** Winter in Hedwigia 1885, p. 261. — Perithe-ciis amphigenis, soros rotundatos vel irregulares, sæpe compactos, $\frac{1}{2}$ -10 mm. latos formantibus, plerumque densissime stipatis, super-ficialibus, globosis, depresso, demum vertice impresso, umbilicato, verruculosis, atris, opacis, ostiolo minutissimo, pertuso, interdum brevissime papilliformi, 210-240 μ. diam.; ascis oblongis, sursum parum, deorsum magis attenuatis, stipite sublongo, gracili prædi-tis, membrana apice valde incrassata, 8-sporis, 70-75 ± 19-20; spo-ridiis subdistichis, ellipticis, 2-cellularibus, cellula superiore multo majori, elliptica, basi subtruncata, cellula inferiore apiculiformi, minutissima, primo hyalinis, demum pallide fuscidulis.

Hab. in foliis vivis Compositæ adhuc ignotæ, scandentis (Mikaniæ spec.?) prope S. Francisco Brasiliæ (E. ULE).

5. **Lizonia Thalictri** Rost. Fungi Groen. p. 556. — Peritheciis ovo-deis, coriaceo-membranaceis, papilla conoidea præditis, siccis col-lapsis; ascis crasse clavatis, apice crassissime tunicatis, 110 ± 45, brevissime pedicellatis; sporidiis octonis, 2-3-stichis, fusoideo-elon-gatis, 1-septatis, 4-guttulatis, mucō hyalino obvolutis, 45-50 ± 10-13.

Hab. in caulis siccis Thalictri alpini, Umanak-fjord
Groenlandiæ.

6. **Lizonia? guaranitica** Speg. Fung. Guar. Pug. I, n. 218. — Macu- 2797
lis amphigenis, subarescentibus non v. obscure determinatis ex or-
biculari irregulariter repando-angulosis, parvis, 2-5 mm. diam., sor-
dide fuscis vel olivascentibus; peritheciis hypophyllis, laxiuscule
gregariis, subsuperficialibus, basi matrici insculptis, primo lepidi-
bus tectis, dein inter illas prominulo-subexsertis, globosis vel glo-
boso-depressis, atris, glabris, levibus, 100-110 μ . diam., vertice
minute perforato ostiolatis, membranaceo-subcoriacellis, contextu
subcrasse parenchymatico, fusco-atro; ascis cylindraceo-obclavatis,
antice truncato-rotundatis crassiusculeque tunicatis, postice abrupte
breviterque attenuato-stipitatis, 45-50 \approx 7-10, aparaphysatis, 8-
sporis; sporidiis hyalinis, 2-3-stichis, fusoideis vel elongato-ellipti-
cis, 18 \approx 2,5, utrinque subtruncato-rotundatis, rectis, inæquilate-
ralibus, medio 1-septatis, non constrictis, primo loculis 2-3-blastis
dein eguttulatis, supero sæpe obtusiore ac crassiore.

Hab. ad folia languida Styracis speciei cujusdam in dumetis prope Borjas Brasiliæ. — An novum genus?

7. **Lizonia? paraguayensis** Speg. Fungi Guar. Pug. I, n. 219. — Ma- 2798
culis nullis; peritheciis hypophyllis in glomerulos sparsis vel laxe
gregariis, rarius confluentibus, parvulis, 0,5-1 mm. diam., dense
constipatis, basi sæpius confluenti-cæspitosis, superficialibus, glo-
bosis vel elliptico-globosis, 100-110 \approx 70-80, umbonato-ostiolatis,
levibus, tomento vel pruinula alba vel fusca laxe adspersis, in sic-
co sæpe collapsis, membranaceo-subcoriaceis, contextu parenchyma-
tico, atro-fuligineo, subopaco; ascis ovatis vel obclavatis, apice ob-
tuse rotundatis vel subtruncatis, crasse tunicatis, basi abrupte sub-
coarctato-stipitellatis, 50-80 \approx 15-20, aparaphysatis, 8-sporis; spo-
ridiis 2-3-stichis vel subinordinatis, elliptico-elongatis vel subfu-
soideis, utrinque obtusiuscule rotundatis, 27-29 \approx 6,5 medio 1-se-
ptatis, non constrictis, æquilateralibus, rectis vel curvulis, eguttu-
latis, hyalinis.

Hab. ad folia viva Styracis speciei cujusdam in silvis apud
pagum Rica Brasiliæ. — An novum genus?

8. **Lizonia Jacquiniae** Br. et Har. Rev. Myc. 1891, pag. 16. — Pe- 2799
ritheciis superficialibus, globulosis, perpusillis, 90-100 μ . diam.,
nigris, conglomeratis, greges minutos, epiphyllos oculo nudo peri-
thecium singulum $\frac{1}{3}$ mm. diam. simulantes, formantibus; ascis
ovoideis, sessilibus, 8-sporis, 20-24 \approx 12-14; sporidiis congregatis,

682 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Melanopsamma.
tereti-oblongis, 1-septatis, obtusis, hyalinis, ad septum vix constrictis, 12-13 \approx 3.

Hab. in foliis Jacquiniae armillaris in insula Jamaica (herb. BERKELEY).

9. **Lizonia Sphagni** Cooke Grev. XVIII, p. 86. — Peritheciis sparsis, 2800 subglobosis, nigris, prominulis, ostiolo mamillato, mox superficialibus; ascis clavatis; sporidiis cylindraceis polo uno vel utroque leniter curvis, 1-septatis, dein 3-septatis?, hyalinis, 40-50 \approx 8.

Hab. in foliis dejectis Sphagni, Maine in Amer. bor.

PSEUDOLIZONIA Pir. N. G. Bot. it. 1889, p. 315 (Etym. *pseudes* falsus et *Lizonia*). — Perithecia carbonacea, discreta vel aggregata, subsphærica, breviter papillata, levia, apice mycelii rhizomorphici, incidentia. Ascii 16-spori, paraphysati. Sporidia didyma, fuscescens. — A gen. *Lizonia* ascis polysporis diversa.

1. **Pseudolizonia Baldinii** Pir. loc. cit. — Peritheciis subsphæricis, 2801 cellulis minutis, polygonis contextis, ostiolo brevi papilliformi; ascis crebris, tereti-clavatis, uno latere convexulis, 122-136 \approx 14-18, brevissime pedicellatis; paraphysibus clavulatis, subseptatis, simplicibus, ascum subæquantibus, evanidis; sporidiis subdistichis, didymis, subfusoideis, inæquilateris, dilute fuligineis, 22-29 \approx 8-9, loculo superiori majori.

Hab. in foliis periandricis Polytrichi communis, Allumiere prope Roma Italia (T. A. BALDINI).

MELANOPSAMMA Niessl — Sacc. Syll. I, p. 675.

1. **Melanopsamma Saccardiana** Bomm. et Rouss. in litt., Sacc. Misc. 2802 Myc. I, pag. 18. — Peritheciis gregariis, minutis, $\frac{1}{6}$ mm. diam., subglobosis, obtuse papillatis, atris, glabrescentibus, sed hyphis dematiaceis (propriis?) rigidulis, fuligineis, plerumque cinctis; ascis subcylindricis, a paraphysatis vel spurie paraphysatis, 60-70 \approx 5-7, apice rotundatis, deorsum attenuatis breviterque stipitatis, octosporis; sporidiis oblique monostichis v. distichis, breve oblongis, utrinque obtusiusculis, diu continuis, tandem distincte constricto 1-septatis, 7-8 \approx 4, 2-4-guttulatis, hyalinis.

Hab. in ligno fagineo putrescente, Groenendael Belgii. — Socia adsunt *Helminthosporium apiculatum* et *Lasiosphæria hispida*.

2. **Melanopsamma Grevillii** Rehm in Starb. Bid. Sver. Ascom. p. 5, 2803 t. I, f. 1. — Peritheciis dense gregariis, superficialibus, primo obconicis, superne applanato-convexis, deinde patellæformiter concavis.

viusculis, ascomata *Tympnidis* in memoriam revocantibus, astomis, valde rugulosis vel fere verrucosis, 0,4-0,5 mm. diam., pilis fusco-atris, in juventute peritheciæ, conidiophoris?, obsessis; ascis e stipite longo filiformi late clavatis, 8-sporis, 40-50 \times 8-10; sporidiis ovoideis vel ovoido-ellipsoideis, uniseptatis, non constrictis, loculis uni-magni- vel pluri-guttulatis, hyalinis, 7-10 \times 3,5-5; paraphysibus guttulatis, gracilibus, non satis perspicuis.

Hab. ad ramos corticatos Pruni Padi prope Johanneslund Ostrogothiæ (GREVILLIUS). — *M. pomiformi* subaffinis.

3. **Melanopsamma? lophiostomoides** Spieg. F. Pat. n. 126. — Perithecii sparsis vel laxe gregariis subseriatis, matrice totaliter immersis, globoso-ovoideis, 300-350 μ . diam., atris, carbonaceis, ostio lo papillulato tantum matricis superficiem attingente non vel vix exerto, rotundo vel compressiusculo, rimose dehiscente (an semper?); ascis cylindraceis, apice obtuse rotundatis crasseque tunicatis, basi in stipitem breviusculum nodulosum attenuato-productis, 130-140 \times 13-14, octosporis, paraphysibus densissimis filiformibus obvallatis; sporidiis oblique monostichis, fusoideo-ellipticis, 24-26 \times 7-8, medio 1-constricto-septatis, primo loculis grosse 2-3-nucleatis ac bis coarctatis (an tetrablastis?), dein simplicibus, levibus, utrinque obtusiusculis, minute subimperspicueque apiculatis, strato mucoso tenuissimo obovolutis, hyalinis vel fumosis.

Hab. in trunco decorticato fluctato secus Rio Galegos in loco Kilik-aiken vocato, Patagoniæ. — Ad *Lophiostomata* nonnihil vergens, notis tamen fructificationis *Melanopsammæ europæe* Sp. peraffinis.

4. **Melanopsamma obtusa** Karst. Hedw. 1888. — Perithecii sparsis, emergendo elevatis, semiimmersis, conoideo-rotundatis vel oblongatis, saepe vertice depresso, atris, subnitentibus, dein poro pertusis, 0,3 mm. diam.; ascis cylindraceo-clavatis, 95-135 \times 11-12 8-sporis; sporidiis monostichis, ovoido-ellipsoideis, utrinque obtusis, medio constricto-septatis, eguttulatis, chlorino-hyalinis, 16-18 \times 6-9; paraphysibus numerosis, coalescentibus, vix 1 μ . cr.

Hab. in ligno Betulæ? ad Vasam Fenniæ.

5. **Melanopsamma ampulligera** Karst. et Starb. Ascom. nov. fenn. p. 2. — Perithecii levibus, gregariis, basi immersis, sphæroideo-conoides, ostiolo, solito papillato, perforatis, glabris, atris, 0,2-0,3 mm. latis; ascis cylindraceis, inferne inflatis, stipite breviusculo, ut plurimum obliquo, 60-66 \times 15-16; sporidiis di- vel (inferne) tristichis,

- 684 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, **Melanopsamma**.
fusoideo-oblongatis, rectis, uniseptatis, ad septum constrictis, loculis uniguttulatis, hyalinis, 18-24 \approx 5-8; paraphysibus coalitis.
Hab. ad corticem vetustum Betulæ in ditione Mustialensi, Fenniæ (K. STARBAECK).
6. **Melanopsamma salicaria** (Karst.) Sacc. *Sph. salicaria* Karst. Myc. 2807
Fenn. II, p. 95. — Peritheciis gregariis, cortici plus minus immersis, peridermio disjecto, vertice denudatis, sphæroideis vel sphæroideo-depressis, ostiolo minuto, papillato, opacis, tenacellis, glabris, nigris, circiter 0,6 mm. lat.; ascis cylindraceo-subclavatis, 160-170 \approx 12-15, octosporis; sporidiis monostichis, ellipsoideis, uniseptatis, ad septum leviter constrictis, chlorino-hyalinis, 18-22 μ . longis; paraphysibus coalescentibus.
Hab. in ramis aridis *Salicis fragilis* prope Abo Fenniæ. — ***M. fallax** Sacc. et Br. Miscell. Myc. II, pag. 22, n. 93: A typo recedit sporidiis minoribus, nempe 12-14 \approx 6-8, et matrice superficie dealbata. In cortice *Salicis albae*, Troyes Galliæ (BRIARD).
7. **Melanopsamma syringica** Karst. Fragm. M. F. XXIII, pag. 1. — 2808
Peritheciis aggregatis, primitus cortice tectis, dein, ea divulsa, nudatis, superficialibus, ovoideis, conoideis, rotundatis vel mutua pressione forma varia, poro pertusis vel rarius ostiolo subpapillato, atris, 0,3-0,4 mm. diam.; ascis clavatis, 105 \approx 18, 8-sporis; sporidiis distichis, ovoideis vel fusoideo-oblongatis, rectis, 1-septatis, 2-guttulatis, ad septum constrictis, chlorino-hyalinis, 15-20 \approx 7-8; paraphysibus ascos superantibus, 2 μ . cr.
Hab. ad ramos exsiccos *Syringæ vulgaris* in horto Mustialensi Fenniæ.
8. **Melanopsamma incrustans** Pass. Diagn. Fung. nuovi, I, n. 41. — 2809
Peritheciis gregariis, contiguis, atro-incrustantibus, subconoideis, opacis, ostiolo simplici pertuso; ascis oblongo-clavatis, paraphysibus tenuibus obvallatis, 8-sporis, 82-90 \approx 18,5; sporidiis distichis fusiformibus, medio septatis et arcte constrictis, loculis biguttulatis, hyalinis, 22-27 \approx 5.
Hab. in ligno sicco denudato *Persicæ*, Vigheffio pr. Parma. — Sporidia ut in *Zignoella Campi-Siliæ* sed majora.
9. **Melanopsamma graopsis** (Ellis) Sacc. *Sphæria graopsis* Ellis Torr. 2810
Bot. Cl. 1882, p. 73. — Peritheciis sparsis, erumpentibus, membranaceis, globosis vel subelongatis, denique subcollabascentibus, 250-370 μ . diam. rugulosis fibrillisque ligneis albicantibus conspersis, ostiolo papillato, leniter circum sulcato; ascis oblongis, subgibbis, sessilibus, 50-58 \approx 20-25, aparaphysatis; sporidiis octonis, inor-

Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Melanopsamma. 685

dinatis, oblongo-navicularibus, hyalinis dein pallide flavidο-fuscelis, plasmate bipartito, 20-25 \approx 8.

Hab. in ramis decorticatis Pini, New Jersey Amer. bor. —

Ad *Amphisphæriam* forte nutat.

10. **Melanopsamma hypophlœa** Pass. Fung. nuovi, IV, p. 4, n. 9. — 2811

Peritheciis superficialibus, sparsis, carbonaceis, globoso-conoideis, ostiolo papillari nitidulo; ascis leniter clavatis, attenuato-stipitatis, paraphysatis, 8-sporis, 75-80 \approx 10; sporidiis distichis, naviculari-elongatis, hyalinis, medio septatis et constrictis, utrinque acutis, loculis biguttulatis, 18-20 \approx 5.

Hab. in cortice detracto ramorum Pini austriacæ, Vigheffio prope Parma.

11. **Melanopsamma cupressina** Ell. et Ev. in Journ. of Myc. 1886, 2812

p. 103. — Peritheciis superficialibus, gregariis, $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam. vel elongatis et subhysterioideis, rugosis, carbonaceo-membranaceis, sursum poro late pertusis (verisimiliter primo ostiolo prominulo donatis); ascis clavato-cylindraceis, 70-75 \approx 10-12, paraphysibus filiformibus obvallatis, octosporis; sporidiis hyalino-luteolis, clavato-oblongis, 1-septatis, circiter 12 \approx 4.

Hab. in ligno dealbato Cupressi in Amer. bor. (LANGLOIS).

12. **Melanopsamma australis** Pass. Diagn. Fung. nuovi, I, n. 42. — 2813

Peritheciis superficialibus, subglobosis, astomis? atris; ascis per-brevibus sacculatis, membrana crassiuscula, paraphysibus parcis, æquantibus cinctis, 8-poris, 25 \approx 7,5; sporidiis lagenæformibus, medio septatis et constrictis, loculo altero rotundato, altero angusto, subacutis, hyalinis, 9-12 \approx 5.

Hab. in ramulis siccis Juniperi, Vigheffio pr. Parma.

13. **Melanopsamma Amphisphæria** Schulz. et Sacc. in Rev. Myc. 1884, 2814

Schulz. Ill. Fung. Slav. n. 110. — Peritheciis dense gregariis, superficialibus nigris, levissimis, globosis vel globoso-conicis, $\frac{1}{5}$ - $\frac{1}{3}$ mm. diam., ostiolo obtuse conico; ascis cylindricis, paraphysatis, brevissime stipitatis, octosporis; sporidiis recte vel oblique monostichis, ellipsoideis, constricto 1-septatis, 17 \approx 8, hyalinis (extra ascos subinde 3-septatis).

Hab. in ramis decorticatis Cydoniæ vulgaris prope Vinkovce Slavoniæ.

14. **Melanopsamma hyalodidyma** Cooke Trans. Roy. Soc. Edin. 1887, 2815

p. 391. — Sparsa, immersa; peritheciis subglobosis, atris, ostiolo punctiformi emergente; ascis cylindraceis; sporidiis ellipticis, uniseptatis, 12-14 \approx 7 $\frac{1}{2}$.

Hab. in ligno, Socotra.

15. **Melanopsamma borealis** Karst. — Syll. I, p. 578. — Var. **minor** 2816

Cooke Grev. XVII, p. 79. — Peritheciis sparsis vel gregariis, minutissimis, basi insculptis, supra convexis nigris, levibus, subpapillatis; ascis cylindraceis; sporidiis monostichis, anguste ellipsoideis, 2-guttulatis, dein tenuiter 1-septatis, hyalinis, $6 \approx 2,5$.

Hab. in ligno emortuo, Shere Britanniæ (CAPRON).

16. **Melanopsamma cubigena** Berk. Cooke Grev. XVI, pag. 92, *Sph.*

obtusissima B. et C. — Syll. II, p. 411. — Sporidia lanceolata, 1-septata, medio constricta, nucleata, hyalina, $45-50 \approx 6-8$.

17. **Melanopsamma nipæcola** Cooke et Mass. Grev. XVI, pag. 92. — 2817

Peritheciis gregariis, magnis $1-1\frac{1}{2}$ mm. diam., atris conoideis, papillatis, basi applanatis; ascis clavatis, stipitatis; sporidiis biseriatis, lanceolatis, medio uniseptatis, hyalinis, $35 \approx 7$ (potius immaturis).

Hab. in cortice Nipæ, Java (KURZ).

18. **Melanopsamma emergens** Schulz. et Sacc. in Rev. Myc. 1884, 2818

Schulz. Ill. Fung. Slav. n. 616. — Peritheciis laxe gregariis, ligno semiimmersis globulosis, sursum conicis, minutis, 1-2 mm. diam., carbonaceis, minute verruculosis, atris; ostiolo obtusiusculo, perforato; ascis tereti-clavulatis, subsessilibus, filiformi-paraphysatis, octosporis; sporidiis distichis oblongis, sæpe curvulis, utrinque rotundatis, medio constrictis, 4-guttulatis, $14-21 \approx 4,5$, hyalinis.

Hab. ad ligna quercina prope Vinkovce Slavoniæ.

19. **Melanopsamma numerosa** Fautrey Rev. Myc. 1891, pag. 76. — 2819

Peritheciis copiosis, superficialibus, sparsis v. confertis, iuitio sphæricis, nigris, rugulosis, dein collabascendo concavo-cupularibus; ascis fusoideis, tunica fragili; sporidiis cylindraceis, rotundatis, 1-septatis, 4-guttatis, $15-17 \approx 3$, hyalinis.

Hab. in ligno decorticato Carpin i ad sepes, Noidan Galliæ.

— *Habitus fere M. pomiformis.*

20. **Melanopsamma Romelliana** Sacc. Add. Syll. p. 97. — Peritheciis spar-

sis v. subgregariis, primum sphæroideis, demum collapsis, subastomis, superficialibus, nitidis, levibus, $\frac{1}{3}-\frac{1}{4}$ mm. diam.; ascis clavatis, apice nonnihil attenuatis, sessilibus vel breviter pedicellatis, 33-48 \approx 7-9, octosporis; sporidiis subdistichis vel in medio congregatis, spuriæ 1-septatis, primo 4-guttulatis, medio vix constrictis, $7-10 \approx 3-4\frac{1}{2}$ (plerumque $8-9 \approx 3\frac{1}{2}$), hyalinis.

Hab. in corbis Orchidearum in calidario Horti botanici Upsaliensis Sueciæ (ROMELL).

21. **Melanopsamma subfasciculata** (Schw.) Cooke Grev. XV, p. 83, 2821

Sphæria Schw. — Sacc. Syll. 4309. — Sporidia ellipsoidea, uniseptata, hyalina, $20 \div 8$.

22. **Melanopsamma conospora** B. et C. in Herb. Berk. 9727, Cooke Grev. XV, p. 82. — Sporidia piriformia, inæqualiter uniseptata, pallide fusca, $8 \div 4$.

THAXTERIA Sacc. (Etym. a cl. mycologo boreali-americano R. THAXTER), *Bizozzeria* Speg. F. Puigg. n. 277 (1889) nec Sacc. (1885). — Perithecia superficialia, globosa, coriacea, atra. Asci octospori paraphysati. Sporidia botuliformia, 1-septata, ex hyalino chlorina. — An melius *Allantosporis*, una cum *Cacosphæria* Speg., adscribendum genus?

1. **Thaxteria didyma** (Speg.) Sacc. *Bizozzeria didyma* Speg. F. Puigg. 2822 n. 277. — Peritheciis dense constipatis, superficialibus, atris, glabris, 0,5-0,7 mm. diam., breviter crasseque stipitulatis, mox profunde umbilicato-collapsis; ascis clavulatis, longe stipitatis, octosporis, paraphysatis, $65-70 \div 10-12$ parte spor., stip. 110-130 $\div 2-5$; sporidiis allantoideis, 1-septatis, fumoso-chlorinis, $20 \div 6$.

Hab. ad corticem crassiorem putrescentem arborum in silvis prope Apiah Brasiliæ. — Jodi ops nulla! Species nobilissima habitu fere lichenino, primo obtutu, *Trichosphæriæ acanthostromæ* (Mont.) Sacc. (*Sphæria foveolata* B. et R.) simillima sed mox subiculi setularumque defectu dignoscenda.

BERTIA De Not. — Syll. I, p. 481.

1. **Bertia collapsa** Rom. F. aliq. Suec. p. 24. — Peritheciis vulgo 2823 dense gregariis, subiculo fusco, hyphis ramosis intricatis, 6 μ . cr. composito, sed sæpe deficiente insidentibus, sub lente evidenter et amœne rugulosis, glabris, obconicis, convexis, papillatis, demum collapso-cupulatis, nigris, $2/3-3/4$ mm. diam.; ascis clavatis vel piriformibus, apice rotundatis, tenuissime tunicatis, 8-sporis, 35-50 $\div 10-12$; sporidiis congregatis vel subtristichis, oblongis, hyalinis, rectis, biguttulatis, demum 1-septatis, 12-18 $\div 4-5$; paraphysibus.... non visis.

Hab. in cortice interiore relaxato truncorum prostratorum Sorbi Aucupariæ prope Upsala Sueciæ.

VENTURIA Ces. et De Not. — Sacc. Syll. I, p. 587.

1. **Venturia chlorospora** (Ces.) Karst. — Sacc. Syll. I, p. 586, — Var. **canescens** Karst. Myc. fenn. II, pag. 190, *Sphærella canescens*

Karst. Fung. Fenn. exsicc. 895. — A typo differt sporidiis dilutissime luteolis, angustioribus, 14-20 \approx 6. — In foliis Salicis acutifoliæ vetustis, Mustiala Fenniæ.

2. **Venturia Genistæ** (Fuck.) Wint. in Hedwigia 1885, p. 26, *Didymella Genistæ* Sacc. Syll. I, p. 546. — Perithecia ostiolo brevi, apice setarum fasciculo brevissimarum, strictarum, atrarum coronato prædita; ascii 80-95 \approx 8-9; sporidia 13-14 \approx 4-5.
3. **Venturia antarctica** Speg. F. Fueg. 207. — Maculis nullis; peritheciis superficialibus subglobosis, 60-90 μ . diam., sparsis v. laxè gregariis, atris, levibus, ad partem superam pilis brevissimis carbonaceis, atris, opacis, rigidulis, acutis, basi non v. vix incrassatis, 40-45 \approx 3-4, laxè vestitis, coriacellis, contextu sinuoso-parenchymatico, densissimo parum distincto, fuligineo, ostiolo minuto pertuso; ascis subfusoideis antice truncatis, crassiuscule tunicatis, bifoveolatis, postice attenuatis ac pedicello crassiusculo suffultis, 25-27 \approx 4-5, octosporis, aparaphysatis; sporidiis recte vel oblique distichis nonnihil inæquilateralibus vel subcurvulis, medio 1-septatis ac leniter constrictis, loculis subæqualibus vel supero non vel vix magiore, utrinque obtusiusculis, 6-7 \approx 2-2,5 minute 4-guttulatis, hyalinis.

Hab. ad hypophyllum foliorum putrescentium Mayteni prope Shammacus Fuegiæ.

4. **Venturia? fuegiana** Speg. F. Fueg. n. 209. — Maculis nullis; peritheciis hinc inde laxè gregariis, ligno denudato insidentibus, subglobosis (siccis collapsis), parvulis 110-130 μ . d., pilis carbonaceis aculeiformibus acutis, opacis, basi non incrassatis, 25-40 \approx 5, præcipue in parte supera laxa vestitis, membranaceo-coriaceis, contextu subindistincto atro, ostiolo impresso perforato; ascis obclavatis, antice rotundato-subtruncatis, crassiuscule tunicatis, 1-foveolatis, deorsum subcoarctato-rotundatis, brevissime minuteque stipitatis, 40-45 \approx 5, octosporis, aparaphysatis; sporidiis recte v. oblique distichis, elliptico-elongatis, quandoque non vel vix obovatis, utrinque obtuse-rotundatis 11-12 \approx 3, rectis vel leniter inæquilateralibus, diblastis, hyalinis.

Hab. ad ramos dejectos decorticatos putrescentes Fagi betuloidis in Staten island Fuegiæ. — *Eriosphæriæ exiguae* Sacc. peraffinis; generi *Venturiæ* præcipue paraphysibus deficientibus adscribenda.

5. **Venturia elastica** Pass. Diagn. Fung. nuovi, III, n. 13. — Peritheciis superficialibus, minutis, setulosis, atris, hypophyllis, in se-

ries lineares transversim digestis vel sparsis; ascis cylindricis, 8-sporis, elongatis; sporidiis biserialibus, ovalibus, medio septatis, hyalinis, 7,5-8 \times 3.

Hab. in foliis putridis Fici elasticæ, Parma in Horto botanico.

6. **Venturia Straussii** Sacc. et Roum. in Rev. Myc. 1884, pag. 95, 2827 tab. LXVII, fig. 2, Patouill. Tab. Anal. Fung. III, p. 133, fig. 299, *Chætomium pusillum* Strauss in Sturm Deut. Pilze III, 34 (nec Fries). — Peritheciis hinc inde congestis, superficialibus, subiculo setoso, fusco, subrepente insidentibus setisque rigidulis, fuligineis, subrectis, septatis undique vestitis, ostiolo obsoleto; ascis teret-clavatis, brevissime stipitatis, 90 \times 16, non vel spurie paraphysatis, octosporis; sporidiis subdistichis, obovato-oblongis, constricto-l-septatis, loculo infero tenuiore, minute quadriguttatis, lenissime chlorinis, 16-20 \times 8-9.

Hab. in ramulis foliisque vivis Ericæ scopariæ in silvis pineis, Bordeaux Galliæ.

7. **Venturia Gaultheriæ** Ell. et Ev. in Journ. of Myc. 1885, p. 155, 2828 — Maculis orbicularibus, brunneis, $1/3$ mm. diam., centro sæpius griseis; peritheciis sparsis, orbicularibus, 75 μ . diam., membranaceis, contextu crasse parenchymatico, sursum setis paucis, nigris, continuis, rectis, divergentibus, 35 \times 3, præditis; ascis ovato-oblongis, 30-35 \times 8-11, basi incrassatis curvulisque, sessilibus, a paraphysatis; sporidiis biseriatis, subhyalinis, ovato-oblongis, 3-4-nucleatis, l-septatis, ad septum parum constrictis.

Hab. in foliis vivis Gaultheriæ procumbentis, Newfield Amer. bor.

8. **Venturia Cassandræ** Peck 38 Rep. St. Mus. p. 104, t. 2, fig. 11-14, 2829 — Maculis rufo-brunneis v. brunneis, sæpe centro griseis; peritheciis amphigenis, minutis, 70-80 μ . diam., nigris, setis minutis, rectis, paucis, divergentibus, 30-40 μ . longis sursum auctis; ascis oblongis, sursum leniter constrictis, 40-45 \times 8-10; sporidiis distichis, oblongis, quadrinucleatis, medio uniseptato-constrictis, 12 \times 5, hyalinis.

Hab. in foliis vivis Cassandriæ calyculatæ, Caroga Americæ borealis.

9. **Venturia ilicifolia** Cooke. — Sacc. Syll. I, p. 588. Diagnosis nova a Winter relata (confr. Hedwigia 1885, p. 98) a nostra non differt nisi setulis peritheciis longioribus, nempe 100-120 \times 15; hinc species nostra ut typi varietas **breviseta** censenda est.

10. **Venturia Grossulariæ** (Auersw. et Fl.) Sacc. Syll. I, pag. 589,

Winter in Hedwigia 1885, pag. 99. — Ascii plerumque obclavati i. e. e basi latiori sursum parum attenuati, 35-40 \approx 9.

11. **Venturia curviseta** Peck in 35 Rep. St. Mus. Bot. 1884, p. 145. 2830

— Peritheciis numerosis, minutis 75-100 μ . diam., coacervatis vel subgregariis, nigris, apice 5-8 setis nigris, divergentibus, basi abrupte curvatis, 75-125 μ . longis, auctis; ascis oblongis, sursum attenuatis, saepe leniter curvulis, 40-50 μ . l.; sporidiis stipatis v. distichis, oblongis, 10,5-12,5 \approx 4-5, hyalinis, parte superiore crassiore.

Hab. in foliis languidis Nemopanthis canadensis Fenter Amer. bor.

12. **Venturia Arctostaphyli** Cooke et Harkness in Grev. XIII, p. 20. 2831

— Amphigena, sparsa, atra; peritheciis subglobosis, superficialibus, strigosis, 120-150 μ . diam., pilis rigidis, acicularibus, 80 μ . long.; ascis obclavatis, sessilibus, octosporis; sporidiis ellipticis, utrinque rotundatis vix constrictis, uniseptatis, flavidis, 12-15 \approx 5.

Hab. in foliis emortuis Arctostaphyli in California.

13. **Venturia elegantula** Rehm in Hedwigia 1885, pag. 241. — Peritheciis ut in *V. Myrtilli*, sed majoribus, sparsis vel confertis, sessilibus aut primitus immersis: vertice setis simplicibus, acutis, fuscis, 60-120 \approx 4 praeditis, parenchymate fusco contextis; ascis pseudopiriformibus, latis, basi sessilibus, 8-sporis, 75-90 \approx 15-21; sporidiis distichis, oblongis, obtusis, 2-cellularibus, cellulis aequaliter longis, saepe 2-nucleolatis, supera latiore, medio vix constrictis, viridulis, 15-21, plerumque 24-27 \approx 6-7; paraphysibus paucis, intricatis.

Hab. ad folia putrida Vaccinii Myrtilli supra Gampenhöfe prope Sulden ad montem Ortler in Tirolia (DR. REHM).

14. **Venturia ditricha** (Fr.) Karst. — Syll. I, p. 587. — Var. **fraxinicola** Rehm Ascom. n. 792. — Peritheciis parenchymaticis, brunneis, apice setulas circ. 12, simplices, obtusulas, bruneas 45-60 μ , long. gerentibus; ascis piriformibus sessilibus, sursum attenuatis, 8-sporis, 50 \approx 9; sporidiis distichis, clavatis 1-septatis, loculo sup. crassiore longiore, flavidis, 10 \approx 4,5; paraphysibus ramulosis.

Hab. in foliis Fraxini in Germania.

15. **Venturia aggregata** Winter in Hedw. 1884, p. 172. — Peritheciis amphigenis, superficialibus, 2-20 (rarius pluribus) densissime aggregatis, stromatis minuti, rotundati, verruciformis, extus atri, intus cinnabarinii superficie insidentibus, depresso globosis, demum cupuliformibus, parcissime setulosis, ostiolo vix conspicuo perforatis, atris; ascis copiosis, clavatis, longe pedicellatis, apicem versus

parum attenuatis, 8-sporis, paraphysibus filiformibus, numerosis, obvallatis, 60-80 \approx 19-21; sporidiis inordinate subdistichis, elliptico-oblongis, plerumque inæquilateralibus, medio uniseptatis et parum constrictis, demum pallidissime fuscidulis, 16-17 \approx 8-9.

Hab. in foliis..... prope S. Francisco Brasiliæ (E. ULE).

16. **Venturia socia** Sacc. et Berl. in Rev. Myc. tab. LIV, f. 4, bis. 2835

— Peritheciis globosis, 200-300 μ . diam., membranaceis, subastomis, contextu laxo, olivaceo, setuligeris; setis erectis, subulatis, quandoque apice recurvatis, continuis, subhyalinis; ascis subclavatis v. subfusoideis, 30-40 \approx 10, octosporis, sessilibus, aparaphysatis; sporidiis distichis, oblongis, utrinque rotundatis, guttuligeris vel spuriæ uniseptatis, 10 \approx 4, hyalinis.

Hab. in pag. infer. foliorum..... in subiculo Asterinæ cū-jusdam, Santos in Brasilia. — *V. nobili* propinqua, a qua setulis subhyalinis etc. differt.

17. **Venturia cupulata** Ell. et Mart. Am. Nat. Bot. 1884, p. 69. — 2836

Peritheciis hypophyllis, sparsis, astomis, 300 μ . diam., basi setosis, apice calvis collapsisque; setis brunneo-nigris, 210 \approx 6; ascis oblongis, 48 \approx 12, octosporis; sporidiis obovato-hyalinis, distichis, uniseptatis, 12-13 \approx 4-4½.

Hab. in foliis vivis Quercus laurifoliæ, New Florida Americae borealis.

18. **Venturia appianata** Ell. et Mart. in Am. Nat. Bot. 1884, p. 69. 2837

— Peritheciis hypophyllis, lentiformibus, 100 μ . diam., centro pertusis, contextu celluloso, subradiato, setis brunneis, parce septatis, 12-15, erectis, 70-100 μ . longis, præditis; ascis oblongis, 40 \approx 10-12; sporidiis di-tristichis, ovato-oblongis, 1-septatis, flaveolis, 10-12 \approx 3.

Hab. in foliis vivis Magnolie glaucæ, New Florida Americae borealis.

19. **Venturia Petasitidis** (Fuck.) Sacc. Syll. I, p. 592, Winter in Hedwigia 1885, p. 98. — Asci oblongi, utrinque parum attenuati sessiles, 8-spori, 43-50 \approx 9-10.

20. **Venturia Balansæ** Speg. Fungi Guar. Pug. I, n. 220. — Macu-

lis nullis; peritheciis hypophyllis, sparsis, globosis, parvulis, 80-90 μ . diam., atris, setulis erectis, raris, cylindraceis, rectis, pluriseptatis, apice obtusis, basi subincrassatulis, 30-120 \approx 4-5, dilute fusco-fuligineis vestitis, vertice minutissime impresso-ostiolatis, in sicco sæpe cupulato-collapsis, tenuiter membranaceis, contextu obscure parenchymatico-fuligineo, subiculo radiante, parcissimo, ex

hyphis tenuissimis, continuis, fuscescentibus, simplicibus composito insidentibus; ascis cylindraceis, apice rotundatis, crassiuscule tunicatis, basi breve attenuato-stipitatis, 45-50 \approx 4,5-5, paraphysibus filiformibus, superne ramulosis, laxe obvallatis, 8-sporis; sporidiis hyalinis, recte et alterne distichis, elliptico-fusoideis, utrinque acutiuscule rotundatis, 10-12 \approx 2-2,5, rectis v. subinæquilateralis-curvulis, medio 1-septatis, constrictulis, loculis minute 2-guttulatis, supero sæpius paullulo breviore, crassiore obtusioreque.

Hab. ad folia viva *Dichondræ repens* in umbrosis silvæ Caa-guazu Brasiliæ.

21. **Venturia palustris** Sacc. Bomm. et Rouss. in Bull. Soc. R. Bot. Belg. 1886, p. 172. — Peritheciis epiphyllis, sparsis, subsuperficialibus, sphæroideis vel subsphæroideis, $1/10$ mm. diam., nigris, pilis variis, brevibus ornatis, macula brunneo-purpurea insidentibus; ascis subsessilibus, curvatis, 45-51 \approx 9, centro inflatis, utrinque attenuatis; sporidiis hyalinis, irregulariter didymis, subfusoideis, 4-guttulatis uni- vel subbiseriatis, septo leniter constrictis, 12-15 \approx 5.

Hab. in foliis languescentibus *Comari palustris*, Ottignies Belgii. — A *V. Potentillæ* macula brunneo-purpurea, ascis fusiformibus, sporidiis hyalinis præcipue differt.

22. **Venturia Potentillæ** (Fries) Sacc. Syll. I, pag. 594, Winter in Hedwigia 1885, p. 98. — Asci oblongo-cylindraceas, 40-45 \approx 8-10; sporidia subdisticha, oblongo-ovoidea seu piriformia, infra medium uniseptata, pallide viridula, 12-14 \approx 4-5.

23. **Venturia islandica** Johans. Svamp. fran Island. p. 168, t. XXIX, 2840 fig. 10. — Peritheciis plerumque hypophyllis, sparsis, superficialibus, nigris, subsphæroideis vel ovoideis, in superiore parte setulis nigris, paucis (7-10), 100-135 μ . longis ornatis; ascis ovoido-elongatis, plerumque inæquilateralibus, paraphysatis, sessilibus, octosporis, 54-57 \approx 18-20; sporidiis distichis vel congregatis, ovoideis vel ovoido-elongatis, prope basim septatis, ad septum non constrictis, hyalinis, 19-24 \approx 6-8, loculo majore 15-20 \approx 3-5.

Hab. in foliis emortuis *Dryadis octopetalæ*, Eskifjordur Islandiæ orientalis.

24. **Venturia nubigena** Speg. Rev. Myc. 14, p. 78. — Peritheciis am- 2841 phigenis, atris, minutis 80-100 μ . diam., innato-erumpentibus, hemisphærico-subglobosis, ubique laxe pilosis; pilis erectis, rigidulis, continuis vel 1-2-septatis, 30-50 \approx 5, fuligineis, apice obtusis; contextu peritheciorum parenchymatico, coriaceo-membranaceo, fuligineo-atro; ascis non visis; sporidiis subclavato-ovoideis, utrinque

obtusiusculis, medio 1-pseudoseptatis, non constrictis, 12 ± 4 , hyalinis.

Hab. in foliis emortuis Arenariæ tetrapetetræ in summis montibus pr. Vigan Galliæ, socio *Helminthosporio nubigeno* Speg.

25. **Venturia glomerata** Cooke — Syll. I, p. 592. — Var. *dissemi-* 2842
nata Schröt. Nord. Pilze p. 10. — Peritheciis hypophyllis, sparsis, infossis, $60-80 \mu$. diam., globulosis, collapsis, vertice setulis nigris, 33 ± 4 , copiosis coronatis; ascis sessilibus, $26-30 \pm 5-7$, octosporis; sporidiis elongato-clavatis, $8-9 \pm 3$, bilocularibus, e hyalino olivaceis.

Hab. in foliis emortuis Geranii silvatici, Magerö Nordcap Norvegiæ.

26. **Venturia Niesslpii** Sacc. — Syll. I, p. 593 — **nigella* Sacc. Mich. 2843 II, p. 607. — Peritheciis dense gregariis erumpentibus globulosis, $110-120 \mu$. diam., ostiolo setulis brevissimis obtusiusculis munito; ascis tereti-obclavatis, $50-60 \pm 14-15$, octosporis; sporidiis cuneatis, $16-18 \pm 6$, loculo infero multo minori, olivaceis.

Hab. in caule infuscato Gentianæ luteæ, Le Sapey (Isère) Galliæ, socia *Phoma Niesslpii* (THERRY).

27. **Venturia circinans** (Fr.) Sacc. Syll. I, p. 592. Huc spectant, teste cl. Winter (Hedwigia 1885, p. 98), *Venturia glomerata* et *Dothidea Robergei*. — Sporidia $4,5 \mu$. tantum lata.

28. **Venturia anthophila** Passer. Erb. critt. Ital. Ser. II, p. 1369. — 2844 Peritheciis sparsis, superficialibus, prominulis, globoso-conicis, atris, opacis parce setulosis; ascis aparaphysatis, rectiusculis, ovato-oblongis, circiter 60 ± 12 ; sporidiis oblongis, didymis, medio subconstrictis, loculo altero angustiore, hyalinis, vase minuteque guttulatis, $23-25 \pm 7-7,5$.

Hab. in floribus exsiccatis Hydrangeæ hortensis in horto bot. Parmensi (PASSEURINI).

29. **Venturia cincinnata** (Fr.) Rostr. F. Groenl. p. 556, *Sphæria* Fr. Syll. II, p. 406. — Asci e basi ampliati, angustati, $85-95 \pm 20-25$; sporidia ovoideo-oblongata, 1-septata, constricta, $22-24 \pm 8-10$, primo hyalina dein olivacea. — Nascitur quoque in foliis *Oxycocci palustris*, Tasermiut Groenlandiæ.

30. **Venturia sabalicola** Ell. et Ev. Proc. Phil. 1890, pag. 233. — 2845 Sparsa vel subgregaria; peritheciis globosis $125-135 \mu$. diam., pertusis, setulis rectis, obtusis, nigris, $50-80 \pm 6-8$; ascis oblongo-clavatis, $50-60 \pm 7-8$, aparaphysatis; sporidiis fusideo-oblongis, hyalinis 4-nucleatis (demum 1-septatis?) $10-13 \pm 2,5-5$, stipato-distichis.

Hab. in foliis emortuis *Sabal Palmetto*, Bayon Chene La.
Amer. bor. (LANGLOIS).

31. **Venturia microchæta** Pat. in Rev. Myc. 1886, p. 82 (*V. micro-* 2846
seta). — Peritheciis minutis, nigris, globulosis vel depressis, gla-
bris, vertice setis acutis, brevibus, 36-45 μ . longis, nigris coro-
nato; ascis cylindraceis, breviter pedicellatis, 75 μ . longis; sporidiis
cylindraceis, 1-septatis, hyalinis vel vix virescentibus, 21 \approx 2-3.

Hab. in foliis emortuis *Caricis*, Tibet oriental. (AB. DAVID).

32. **Venturia erysiphoides** E. et E. Journ. Myc. 1887, p. 128. — Pe- 2847
ritheciis gregariis, nigris, globosis, 100 μ . diam., late perforatis,
setulis rigidis, nigris, continuis, 60-70 \approx 5-6 conspersis, ascis ob-
longis, sessilibus, paraphysatis, 40-45 \approx 7-8; sporidiis stipatis, fu-
soideis, curvulis, 5-9-nucleatis, 20 \approx 2,5, hyalinis.

Hab. in culmis v. vaginis emortuis *Panici Curtisii*, Pointe
à la Hâche La. Amer. bor. (LANGLOIS). — Affinis *V. graminicolæ*.

33. **Venturia microspora** Speg. F. Fueg. n. 208. — Maculis nullis; 2848
peritheciis sparsis, v. gregariis, subseriatis globosis, 120-150 μ . d., in
sicco collabascentibus, pilis opacis atris acutis basi non v. vix di-
latatis, 20-30 \approx 2-3, undique densiuscule vestitis, tenui-membrana-
ceis; contextu indistincto atro opaco, ostiolo vix papillulato obtu-
so; ascis subclavulato-fusoideis, antice late truncato-rotundatis, non
v. vix crassiuscule tunicatis, deorsum breviter attenuato-pedicel-
latis, 24-25 \approx 2-2,5, octosporis, paraphysatis; sporidiis elliptico-
elongatis, non v. leniter inæquilateralibus v. subcurvulis, medio 1-
septatis non constrictis, utrinque obtuse rotundatis v. subtruncatis,
5 \approx 1, loculis grosse biguttulatis, hyalinis.

Hab. ad folia dejecta v. languida *Festucæ magellanicae* in
Staten island Fuegiæ.

34. **Venturia Haglundi** Starb. Skand. Pyr. p. 9 (*Niesslia*). — Peri- 2849
theciis superficialibus vel subgregariis, sæpe 2-3-connatis, 100 μ .
diam., setulis 40 \approx 5, sparsis, firmis, rectis, basi subincrassatis ve-
stitis; ascis paraphysatis, cylindraceis, sursum sensim attenuatis,
subtruncatis, sessilibus, 25-30 \approx 2,5-3,5, 8-sporis; sporidiis oblique
monostichis, fusoideis, continuis vel demum 1-septatis, 10-13 \approx
1,5-2,5.

Hab. in foliis subvivis *Lycopodiæ complanati* pr. Dufhult
in parœcia Simonstorp Ostrogothiæ.

35. **Venturia Turfosorum** Mouton in Bull. Soc. R. Bot. Belg. 1886, 2850
p. 149. — Peritheciis sparsis, superficialibus, subovatis, 80-100 μ .
diam., membranaceis, contextu fuligineo, ostiolo minuto, obtuso ob-

scuriori, setulis plerumque curvatis, basi incrassatis, continuis, 55-70 \approx 3,5, laxe vestitis; ascis obpiriformibus, breve stipitatis, octosporis, 22 μ . latis; sporidiis ovatis, bilocularibus, ad septum constrictis, hyalinis, 12-14 \approx 6-7.

Hab. supra Sphagna siccitate emortua Tilff in Belgio.

36. **Venturia fimiseda** Mout. Ascom. Lièg. II, pag. 5, tab. I, fig. 14. 2851

— Peritheciis dense sparsis vel gregariis, superficialibus, globulosis, haud collabentibus, 160 μ . diam., ostiolo punctiformi, contextu membranaceo fusco, setulis plerumque continuis, 4 μ . cr. vestitis; ascis e basi dilatata oblongis, sessilibus, octosporis, 42 \approx 10; sporidiis inordinate distichis, longe ovatis v. cuneatis, chlorino-luteolis, medio septatis lenissimeque constrictis, 4-guttulatis, 12-15 \approx 4.

Hab. in fimo vetusto Leporum, Forêt Belgii. — *Venturiæ Myrtilli* proxima.

37. **Venturia pezizoidea** Sacc. et Ell. Mich. II, p. 567. — Peritheciis 2852

laxe gregariis hypophyllis, punctiformibus, subsuperficialibus, globulosis, dein collabentibus, setulis cuspidatis, 40-60 \approx 5, fuligineis vestitis; perithecii contextu tenuissimo, pallidulo; ascis teretico-clavatis, 35 \approx 8, octosporis, apophysatis; sporidiis (immaturis visis) teretiusculis, continuis, hyalinis, 8-10 \approx 2.

Hab. in pag. inf. foliorum languidorum Andromedæ? race-mosæ, Newfield N. J. Amer. bor. (ELLIS).

38. **Venturia parasitica** Ell. et Ev. Proc. Phil. 1890, p. 233 (*Venturiola?*). — Peritheciis dense gregariis, globosis, 90-100 μ . diam., collabascentibus supra spinulis continuis, rigidis, rectis, 30 \approx 5, sparse vestitis; ascis clavato-cylindraceis, 25 \approx 5, apophysatis; sporidiis oblique monostichis vel distichis, oblongis, 3-4-nucleatis, 6-8 \approx 2, hyalinis. 2853

Hab. parasitica in Hypoxylon (perforato?) ad corticem Magnoliæ pr. St. Martinsville La. Amer. bor. (LANGLOIS).

Observatio

Species sequentes, in Sylloge sub aliis generibus militantes, sec. cl.

Winter (Confr. Hedwigia 1836, p. 27) ad hoc genus pertinent, nam perithecia ostiolo apice fasciculo setarum brevium prædita sunt sporidiaque uniseptata.

39. **Venturia Geranii** (Fr.) Winter, *Stigmata Geranii* Sacc. Syll. I, pag. 541.

40. **Venturia epilobiana** (Sacc.) Winter, *Læstadia epilobiana* Sacc. Syll. I, p. 429.

41. **Venturia Systema-solare** (Fuck.) Winter, *Læstadia Systema-solare* Sacc. Syll. I, p. 423.

ARCANGELIA Sacc. Bull. Soc. myc., 1890, V, pag. 115, t. XIV, f. 1
 (Etym. a cl. botanico italicico J. ARCANGELI, in Universitate Pisana Botanices professore). — Perithecia frondibus vivis (Hepaticarum) immersa, subglobosa, membranacea, hyphis prælongis remotis conspersa, nigricantia, ostiolo rotundo vix papillato pertusa. Ascii cylindracei, octospori. Sporidia didyma, hyalina. Paraphyses distinctæ, ramulosæ. — A gen. *Lizonia* differt peritheciis immersis, membranaceis, sporidiis achrois; a gen. *Didymella* peritheciis biogenis, longe piligeris; a gen. *Stigmataea* demum peritheciis longe piligeris, paraphysibus copiosis, ramulosis. Gen. *Lizonia*, olim uti paraphysatum descriptum, teste cl. Piotta in N. G. bot. ital. 1889, p. 317, paraphyses genuinas gerit.

1. **Arcangelia Hepaticarum** Sacc. loc. cit. XIV. — Peritheciis sparsis v. 2-4 approximatis, epiphyllis, immersis dein vix erumpentibus, e globoso obtuse conoideis, 150-200 μ . diam., superne magis pachydermaticis, nigris, infra molioribus pallidioribus, hyphis seu pilis longis, ramulosis, septatis fuligineis, remotiusculis, matricem intrantibus ubique, præcipue basi, cinctis; ostiolo brevissime obtuse papillato, latiuscule pertuso, margine (sub microscopio) tenuissime denticulato; ascis fasciculatis, cylindraceis, breve stipitatis, apice rotundatis, octosporis, 120 \pm 12-15; sporidiis monostichis vel apice subdistichis ovato-oblongis, constricto 1-septatis, hyalinis, 20-22 \pm 8-10, loculo superiore breviore, rotundiore; paraphysibus filiformibus simplicibus furcatisve.

Hab. in frondibus omnino vivis Ricciae tumidae ad Poggio S. Romolo prope Florentiam. — Legit cl. E. LEVIER et communicauit cl. U. Martelli.

ERIOSPHÆRIA Sacc. Syll. I, p. 397.

1. **Eriosphæria vermicularioides** Sacc. et Roum. in Rev. Myc. 1883, tab. 39, fig. 3. — Peritheciis dense gregariis superficialibus, globoso-papillatis, nigris, minutis, $1\frac{1}{6}$ - $1\frac{1}{5}$ mm. diam., subcarbonaceis, setulis filiformibus, rigidis, septatis, fuligineis, basi cinctis parceque supra conspersis; ascis cylindraceis, breve stipitatis, 70-80 \pm 5, octosporis; sporidiis oblique monostichis, ellipsoideis, 1-septatis constrictisque, utrinque rotundatis, 7-8 \pm 4, hyalinis.

Hab. in lignis putrescentibus in Gallia. — Ab *E. Vermicula-*

ria differt ascis cylindraceis, longioribus et angustioribus; sporidiis monostichis; peritheciis supra sæpius glabratiss.

2. **Eriosphæria corylina** Mout. Ascon. Lièg. II, p. 6, t. I, f. 10. — 2856
 Peritheciis globosis, ligno duro insidentibus, media magnitudine, pilis molibus pachydermaticis subhyalinis, continuis parce vestitis, contextu coriaceo fuscescente; ascis subcylindraceis, in stipitem attenuatis, 4-sporis, 120-130 \approx 10-13, p. sp. 90. μ . long.; paraphysibus filiformibus; sporidiis oblique monostichis, oblongo-cylindraceis, obtusis rectis, medio septatis, vix constrictis, plasmate, granuloso, dilute olivaceo-fuscescente, 28-32 \approx 8,5.

Hab. ad caudices sicclos Coryli Avellanæ, Beaufays Belgii.

3. **Eriosphæria? vulgaris** Speg. F. Fueg. n. 211. — Peritheciis sparsis, superficialibus, subglobosis 350-400 μ . d. atris, canescenti-villosis, pilis rectis acutis, 1-cellularibus basi fuscescentibus apice hyalinis tenuissime canaliculatis 500-900 \approx 20, rigidulis dense vestitis, coriaceis, contextu opaco atro indistincto, ostiolo minute umbonato; ascis....?; sporidiis....?, non visis.

Hab. vulgatissima ad ramos dejectos decorticatos putrescentes Fagi betuloidis per totam Fuegiā (semper sterilis).

4. **Eriosphæria australis** Speg. F. Fueg. n. 210. — Peritheciis sparsis 2858 globosis 300-450 μ . d. peridermio sublevato velatis, eoque sæpius sedente liberis, superficialibus, atris, molliter rufesceni-villosis, pilis cylindraceis longiusculis obtusis, basi non incrassatis 100-150 \approx 5 rectis v. flexuosulis, pluriseptatis, pallide rufescensibus vestitis, coriaceo-membranaceis, contextu parenchymatico, fuligineo; ostiolo impresso pertuso; ascis fusoideo-subclavatis 120-130 \approx 12-13, antice truncato-rotundatis, deorsum breviter crasseque attenuato-pedicellatis, octosporis, subdiffluentibus, paraphysibus parcis mucosis; sporidiis recte distichis, elliptico-elongatis 24-28 \approx 6-7, leniter inæqualiteribus v. subcurvulis, utrinque obtusiusculis, primo continuis hyalinis, dein grosse 2-guttulatis, postremo diblastis, dilute fuscescentibus.

Hab. ad ramos emortuos Berberidis ilicifoliæ prope Port Cook in Staten island Fuegiæ.

5. **Eriosphæria inæqualis** Grove in Add. Syll. p. 103. — Peritheciis ovoides, circiter 150 μ . diam., confertis, superficialibus, parte inferiori pilis brevibus, rigidis, radiantibus obsitis, superiori nudis prominulis, rugosiusculis, nitidis; ostiolo conico, pertuso; ascis stipatis, cylindricis, 50 μ . long., subinde basi attenuatis et deinde 70-80 \approx 3-4; sporidiis monostichis, oblongis, 7-8 \approx 2 $\frac{1}{2}$ -3, inæqualibus, 2-3-

guttulatis, hyalinis, maturis tenuiter 1-septatis; loculo superiori ovoideo 1-guttulato, inferiori angustiori et longiori, attenuato, conico, 2-guttulato, loculis demum facile secedentibus.

Hab. in ligno putri, una cum *Gonytricho cæsio*, Barnt Green prope Birmingham, Brit. — *Trichosphæriæ pilosæ* manifeste affinis et ab iconе Curreyana (Linn. Trans. XXII, t. 57, f. 21) vix nisi sporidiis inæqualibus differens. Cum *Trichosphæriæ pilosæ* var. *nitidula* Sacc. comparanda, quæ probabiliter status junior videretur.

6. **Eriosphæria pulchriseta** (Peck.) Sacc. *Acanthostigma pulchrisetum* (Peck) Sacc. Syll. II, n. 3605. — Sporidia brevia 6,2-7,5 μ . longa et medio partita, ergo potius *Eriosphæria* quam *Acanthostigma*.
7. **Eriosphæria calospora** Speg. Fungi Guar. Pug. I, n. 211. — Pe- 2860 ritheciis dense sparsis v. laxe gregariis in matrice immutata v. vix dealbata superficialibus, udis, globosis, 200-250 μ . diam., sic- cis cupulato-collapsis, setulis pluriseptatis, fuligineis, breviusculis, 50-100 \pm 5, subcylindraceis, apice acutatis v. subclavulato-hyalinis, basi non incrassatis laxissime vestitis, membranaceo-coriacellis, atris, levibus, contextu dense subimperspicueque parenchymatico, oliva- ceo, subopaco, ostiolo minute umbonatulo ornatis; ascis ellipticis v. elliptico-subfusoideis, antice truncato-rotundatis, postice brevissime crasseque attenuato-stipitatis, 45-55 \pm 13-20, aparaphysatis, 8-sporis; sporidiis hyalinis, recte v. oblique distichis, elliptico-elongatis, utrinque acutiuscule rotundatis, non v. vix inæquilateralibus, rarius subsigmoideis, medio 1-septatis, non v. leniter constrictis, loculis subæqualibus non v. 2-guttulatis, episporio primo levi, dein transverse rugoso, rugis minute tuberculosis.

Hab. ad culmos emortuos Bambusaceæ scandentis cuiusdam in silva subvirginea, Caa-guazù Brasiliæ.

LENTOMITA Niessl — Sacc. Syll. I, p. 585.

1. **Lentomita longicollis** Karst. in Hedw. 1884, n. 6, p. 85 et Rev. 2861 atque Synop. Ascom. fenn. p. 58. — Peritheciis gregariis, innatis dein semierumpentibus, rarissime liberis, sphæroideis, nudis, gla- bris, fragilibus, atris, rostro cylindraceo, recto, vulgo perithecium æquante, 4 μ . lat.; sporidiis octonis, monostichis, oblongato-elli- psoideis, ultra medium uniseptatis, junioribus 1-2-guttatis, ad septum non constrictis, chlorino-hyalinis, 11-16 \pm 5-6.

Hab. in ligno vetusto Betulæ in Mustiala Fenniæ. — Perithe- ciis majoribus, basi haud applanatis, rostro longiore; sporidiis ma-

boribus, ultra medium spurie uniseptatis etc., ab affini *Lent. brevicolle* differt.

2. **Lentomita acuum** Mout. Ascom. Lièg. II, pag. 5. — Peritheciis superficialibus, minimis, 100 μ . d., contextu membranaceo, fuligineo, parenchymatico, in collum capillare, nigrum, prosenchymaticum, 400 -700 μ . long., 25 μ . cr., apice obtusum subhyalinum productis; ascis paraphysatis (?), clavatis, fugacibus, octosporis, apice membrana non incrassata, 60 \approx 10; sporidiis biseriatis, elongato-oblongis, hyalinis, 25 \approx 6, tenuiter 1-septatis, 4-guttatis, non constrictis.

Hab. in foliis putrescentibus Pini silvestris, Tilff Belgii.

GIBBERA Fr. — Sacc. Syll. I, p. 599.

1. **Gibbera guaranitica** Speg. Fungi Guar. Pug. I, n. 222. — Peritheciis superficialibus, dense lateque cæspitoso-stipatis, plagulas 1-10 cm. diam., granulosas, sordide rubiginosas efficientibus, ob-ovatis, nigro-fuscis, levibus, parvulis, 600 \approx 300-350, primo ferrugineo-pruinulosis, dein glabratis, ad basim tantum subtomentosulis, hyphis repentibus, parce ramosis, septulatis, gracilibus 3-4 μ . crass., fuligineis donatis, ostiolo minuto, umbilicato, radiato-ruguloso, coriacellis; contextu parenchymatico-indistincto, fuligineo-olivaceo, subopaco; ascis cylindraceo-clavatis, apice rotundatis vel acutatis, basi longiuscule attenuato-pedicellatis p. sp. 65-70 \approx 10-12, ped. 25-30 \approx 3; paraphysibus filiformibus, longioribus, simplicibus v. parce ramulosis, densis obvallatis, 8-sporis; sporidiis oblique monostichis v. recte distichis, ellipticis, saepius inæquilateribus, medio 1-septatis, constrictulis, utrinque acutato-rotundatis, 15-18 \approx 6-7, loculis crasse 1-guttulatis, perdilute olivascentibus.

Hab. ad trunco dejectos putrescentes in silvis prope Guarapí Brasiliæ.

2. **Gibbera pezizoidea** Pat. Champ. N. Caled. p. 9. — Peritheciis aggregatis tuberculum subgloboso-oblongum, 5-15 mm. long. formantibus, stromate rufo-nigro insertis, distinctis, globulosis, vertice depresso, infra villosis, supra glabrescentibus; ascis cylindraceis, 100-10, octosporis, monostichis; paraphysibus coalescentibus, ramosis; sporidiis oblongo-ellipsoideis, pallide olivaceis, 1-septatis constrictisque, subinde 2-guttulatis, 16-17 \approx 6.

Hab. in cortice arborum, Bonrail Novo-Caledoniæ.-Stroma pilis brunneis, simplicibus ramosisque tomentosum.

CACOSPHÆRIA Speg. F. Fueg. n. 218 (1887) (Etym. *cacos* malus et

700 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Melanconis.
Sphæria). — Perithecia *Calosphæriæ*; asci clavati octospori, paraphysati; sporidia botuliformia, 1-septata, hyalina.

1. **Cacosphæria antarctica** Speg. l. c. — Acervulis sparsis v. laxe 2865 gariis, discoideis 1-1,5 mm. diam., pustuliformibus, in cortice nidulatibus, peridermio parum elevato tectis ac eum disculo perforantibus; peritheciis paucis 4-9 in quoque acervulo valsiformiter laxe aggregatis, nudis, atris, nitidulis, glabris, parvulis, 280-320 μ . d. coriacellis, contextu opaco indistincto, ostiolis breviusculis, obtusis, crassiusculis, conniventibus, disculum non v. vix exsertum sæpe griseo-pulverulentum efformantibus; ascis clavatis, antice obtusissime rotundatis crassiuscule tunicatis, deorsum in pedicellum longiusculum attenuatis, p. sp. 22-25 \approx 4-5, ped. 18-25 \approx 2, octosporis; paraphysibus duplo longioribus, parcis, crassiusculis; sporidiis di-tristichis, botuliformibus utrinque obtuse rotundatis 9-11 \approx 1,5, primo diblastis, dein distinete medio 1-septatis, non constrictis, hyalinis.

Hab. ad ramos dejectos emortuos Ribis magellanici prope Ushuvaia, Fuegia.

ENDOTHIA Fr. — Syll. I, p. 601.

1. **Endothia Parryi** (Farlow) Cooke in Grevillea XIII, p. 102. *Do-* 2866 *thidea Parryi* Farlow in litteris. — Stromatibus pulvinate erumpentibus; peritheciis numerosis, concentrica aggregatis, fuscis, epidermide cinnabarina cinctis; ascis linear-clavatis, 76-110 \approx 16-20, a paraphysatis; sporidiis octonis, irregulariter subdistichis, hyalinis, 20-26 \approx 5-7, uniseptatis, ellipsoideis, subacutis, parce constrictis.

Hab. in Agave Shawii, S. W. States Americæ bor.

MELANCONIS Tul. — Sacc. Syll. I, p. 602.

1. **Melanconis Aceris** Phill. et Plowr. in Grevillea XIII, p. 77. — 2867 Stromate valsoideo, atro, suborbiculari vel conico-truncato, 2-5 mm. diam., corti-vel lignicolo, linea atra circumscripto; peritheciis $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., 5-15 coacervatis, ostiolis minutis; ascis clavatis 65-70 \approx 10; sporidiis hyalinis, uniseptatis, nucleatis, ad septum leniter constrictis, ellipsoideis, 12-15 \approx 4-5, muticis.

Hab. in Acere in Britannia (W. B. GROVE).

2. **Melanconis Everhartii** Ell. in Bull. Torr. bot. Club. 1883, p. 117. 2868 — Peritheciis globosis, 0,25 mm. diam., in stromata compactis, 4-18; ascis clavato-cylindraceis, 114-120 \approx 19; paraphysibus crassiusculis, remote septatis, granulosis; sporidiis distichis, oblongo-ellipsoideis, uniseptatis, subhyalinis, 34-38 \approx 11.

Hab. in ramulis delapsis Aceris, West Chester in Amer. bor.

3. **Melanconis dasycarpa** Ellis et Kellerm. in Journ. of Myc. 1886, 2869 p. 3. — Stromate corticali, 1 mm. diam., vel paullo latiore, linea nigra circumscripto; peritheciis 3-8 cortice interiore immersis, $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam., ostiolis asperulis, cylindraceo-conoideis, epidermidem stellato-ruptam sæpe perforantibus, fasciculatis et in discum protrusis; ascis oblongo-cylindraceis, subsessilibus, 100-115 \pm 18-20; paraphysibus imperfecte evolutis; sporidiis distichis, oblongo-cylindraceis, leniter curvulis, uniseptatis, subhyalinis, septo non vel vix constrictis, 25-30 \pm 8-10, appendicula crassiuscula, leniter obliqua, hyalina, 16-20 \pm 3, utrinque præditis.

Hab. in ramulis putridis Aceris dasycarpi Manhattan Kansas Amer. bor. — Affinis *M. Alni*.

4. **Melanconis antarctica** Speg. F. Pat. n. 128 — Stromate valso, elliptico, 5-6 mm. long., 2-3 mm. lat., 0,3-0,5 mm. crass., disco excepto, peridermio tecto, e substantia corticis fere immutata efformato, in parte centrali albescenti-flavescente, vix protuberante; peritheciis in singulo stromate 15-25, eoque immersis, circinantibus, sphæroideis 250-300 μ . diam., glabris levibus, ostiolis in disco corticem perforante sed vix prominulo exsertis, conoideis, obtusiusculis, rigidulis, nigris, constipatis, subconfluentibus; ascis cylindraceo-subclavatis, superne rotundato-subtruncatis, minute bifoveolatis, in pedicellum mox fluxilem productis, p. sp. 80-85 \pm 10-11, octosporis, a paraphysatis?; sporidiis hyalinis v. fumosis distichis, elongato-ellipticis vel subfusoideo-ellipticis 25-26 \pm 5, medio 1-septatis leniterque constrictis, non v. vix subinæquilateralibus utrinque obtusiuscula acutatis et caudula gracili subflexuosa, 20-25 \pm 1-1,5 hyalina auctis.

Hab. ad ramos emortuos Fagi betuloidis in silvis prope Punta Arenas Patagoniæ. — Species *M. Alni* Tul. valde affinis sed certe diversa!

5. **Melanconis salicina** Ell. et Ev. Proc. Phil. Academ. 1890, p. 236. 2871 — Stromate plano, tenui, orbiculari, 2 mm. d., e corticis substantia leniter mutata et vix elevata formato, linea nigra parum profunda circumscripto; peritheciis 3-6, rarius singulo, in quoque stromate, $\frac{3}{4}$ mm. d., globosis membranaceis in collum breve ostiolo latiusculo, concavo epidermidem perforante concontractis; ascis late fusoideis 90-110 \pm 12-16 p. sp. copiose paraphysatis; sporidiis stipato-distichis oblongo-fusoideis, 1-septatis leniter constrictis curvulis, 40-60 \pm 8-10, flavidо-hyalinis, utrinque breve obtuseque apiculatis.

Hab. in ramis emortuis Salicis, London Canada Amer. bor.-
Spermogonia adsunt; perithecia ligno basi insculpta.

HERCOSPORA Tul. — Syll. I, p. 605.

1. **Hercospora tiliacea** (Ell.) Sacc. *Diatrype tiliacea* Ellis Amer. Nat. 2872
1883, Febr. p. 195. — Peritheciis subelongatis, $\frac{1}{2}$ - $\frac{1}{3}$ mm. diam.,
cortice parum mutato immersis, 5-10 aggregatis, ostiolis fascicu-
latis rugulosis, conico-cylindricis, erumpentibus, in discum vix co-
hærentibus; ascis late oblongis, 80-90 \approx 18-22, paraphysatis (?),
octosporis; sporidiis 2-3-stichis, oblongo-cylindraceis, leniter cur-
vis, obtusis, constricto-1-septatis, 2-nucleatis, 22-30 \approx 7-8, hyalinis.

Hab. in cortice Tiliæ americanæ, Ames Iowa Amer. bor.
(J. C. ARTHUR). — « An mera forma *H. Tiliæ?* » — Ostiola $\frac{1}{2}$ -1
mm. longa, denique apice irregulariter dehiscentia; acervuli peri-
theciorum lineola stromatica superficiali circumscripti.

DIAPORTHE Nits. — Sacc. Syll. I, p. 607.

I. Chorostate Nits.

1. **D. Chorostate Bloxami** (Cooke) Berl. et Vogl. Add. Syll. p. 105. 2873
Valsa Bloxami Cooke in Grevillea XIV, p. 47. — Pustulis con-
vexis, emergentibus; peritheciis 12-20 congestis, piriformibus, in
collum elongatum productis, pulvere aureo-fulvo obductis; disco
orbiculari, fusco; ascis cylindricis, octosporis; sporidiis uniseriali-
bus, ellipticis vel sublanceolatis, medio uniseptatis, leniter constri-
ctis, hyalinis, 22 \approx 6.

Hab. in ramis Fagi? Twycross Britanniæ (BLOXAM).

2. **D. Chorostate punctata** (Cooke) Berl. et Volg. Add. Syll. p. 108 2874
Valsa punctata Cooke in Grevillea XIV, p. 47. — Pustulis con-
vexis, emergentibus, ostiolis brevibus, in discum orbicularem con-
gestis; ascis clavatis, breviter stipitatis, octosporis; sporidiis bise-
rialibus, ellipticis, rotundatis, medio uniseptatis, nec constrictis,
hyalinis, 20 \approx 7.

Hab. in ramis Tiliæ et Salicis in Sydenham Britanniæ.

3. **D. Chorostate farinosa** Peck 40 Rep. p. 69. — Stromate sub. 2875
pulverulento sordide alutaceo, e cortice parum mutato formato, in
discum parum exsertum minutum erumpente; peritheciis valsoideo-
circinantibus, 4-10 in quoque circulo, ostiolis atris discum alutaceum
v. demum brunneum punctatum reddentibus; ascis subcylindricis, 60-
70 \approx 8-10; sporidiis confertis v. biseriatis, oblongis v. subfusifor-
mibus, 1-septatis, vulgo 4-nucleatis, 15-20 \approx 4-5.

Hab. in ramis emortuis *Tiliæ americanæ* Argusville America borealis.

4. **D. Chorostate concrescens** (Schw.) Cooke in Grevillæ XIII, pag. 38. *Sphæria concrescens* Schw. Syn. Amer. bor. 1301, et Sacc. Syll. II, p. 378. — Sporidia fusiformia, quadrinucleata, demum uniseptata, 12 μ . longa.
5. **D. Chorostate fuscidula** (Cooke) Berl. et Vogl. Add. Syll. p. 106 2876
Valsa fuscidula Cooke in Grevillea XIV, pag. 48. — Pustulata, fusca; pustulis erumpentibus, convexis, subpulveraceis; peritheciis paucis, immersis, collis elongatis, convergentibus, demum punctato-emergentibus; ascis clavatis, breviter stipitatis, octosporis; sporidiis inordinatis, arcte fusoideis, binucleatis, demum uniseptatis, hyalinis, 18-20 \approx 3,5.
Hab. in ramulis *Quercus* verisimiliter in Britannia.
6. **D. Chorostate crinigera** Ell. et Ev. Proc. Phil. 1890, p. 234. — 87 7
 Stromate corticali, corticem interiore late infuscante; peritheciis cortice interiore immersis, in pustulas 2 mm. diam., subcircinatis, 6-20 congregatis, zona nulla limitatis ovato-globosis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. d., ostiolis brevibus convergentibus, apice rotundatis, levibus v. quadrifurcatis, fasciculatim erumpentibus, subinde longioribus cylindraceis 1-2 mm. long.; ascis 45 \approx 7-8, paraphysatis (?); sporidiis biseriatis, oblongo-fusoideis, 4-nucleatis, medio leniter constrictis, utrinque subobtusis, 10-13 \approx 3-4.
Hab. in ramis emortuis *Quercus*, London, Canada (DEARNES) Delaware (COMMONS). Affinis *D. Woolworthii*.
7. **D. Chorostate leiphæma** (Fr.) Sacc. Syll. I, p. 615. — Var. **major** Brun. in Rev. Myc. 1886, p. 205. — Ascis 110-120 \approx 10-12; sporidiis 20-22 \approx 4-5. In lignis vetustis *Quercus* pr. Saintes Galliæ. *Sphæria Micheliana* Fr. — Syll. II, n. 4217 — secund. Cooke et ad exempl. Berkel. est eadem ac *D. leiphæma*.
8. **D. Chorostate Briardiana** Sacc. Miscell. Myc. I, pag. 3. — Peritheciis subcutaneo-prominulis, in acervulos valseos, exiguo, gregarios dense congestis, globulosis, $\frac{1}{3}$ mm. diam., ostiolis breviusculis, vix convergentibus, peridermum vix perforantibus; ascis oblongo-fusoideis, subinæqualibus, 44-52 \approx 6-8, octosporis; sporidiis monostichis vel distichis, oblongo-ellipsoideis, utrinque obtusiusculis, 12-14 \approx 4, medio leniter constrictis tenuiterque 1-septatis, 4-guttatis, hyalinis.
Hab. in ramis emortuis salicinis pr. Troyes Galliæ (BRIARD).

9. **D. Chorostate nigro-cincta** Pass. in Brun. Nouv. Fragm. p. 27. 2879

— Stromate valseo, zona nigra circumscripto; peritheciis subcircinantibus, parcis; ostiolis fasciculatis per peridermum fissum erumpentibus; ascis cylindraceis, aparaphysatis (?), 8-sporis, $110 \approx 10$; sporidiis monostichis, ellipsoideis, medio 1-septatis, non constrictis, grosse 2-guttatis, hyalinis, $12-20 \approx 7,5$.

Hab. in ramulis emortuis *Castaneæ* vescæ, Saintes Galliæ.

10. **D. Chorostate marginalis** Peck 39 Rep. p. 52, t. II, f. 1-5. — 2880

Pustulis numerosis, epidermide elevata tectis; peritheciis 8-15 in pustulas valsoideo-aggregatis, cortice interiore nidulantibus zona atra nulla cinctis, ostiolis circa disculum albo-pulvereum breve erumpentibus, atris; ascis subcylindraceis, $65-75 \approx 10-13$; sporidiis confertis v. distichis, 1-septatis, utrinque obscure apiculatis, $20-23 \approx 5-6$.

Hab. in ramis emortuis *Alni viridis*, Elizabethtown Am. bor.

11. **D. Chorostate nivosa** Ell. et Ev. Proc. Phil. 1890, p. 222. — 2881

Peritheciis saepius 8-12 subcircinatis, $\frac{3}{4}$ mm. diam., cortice immutato sed pustulatim elevato immersis, linea nulla circumscriptis; ostiolis brevibus apice subhemispherico-papillatis circa discum nivem 1 mm. diam., erumpentibus; ascis 60 \approx 12 p. sporif.; sporidiis subseriatis, oblongis, 4-nucleatis, 1-septatis, constrictis, $12-16 \approx 3-4$, rectis curvulisve.

Hab. in ramis *Alni* emortuis, Isle Royale Lake Sup. Amer. bor. (HOLWAY) « An forma *Diap. marginalis* Peck? »

12. **D. Chorostate populea** Sacc. in Mout. Ascom. Lièg. II, pag. 2882

6. — Peritheciis hinc inde dense acervulatis, globosis, atris, crasse coriaceis, majusculis, 600μ . d., ostiolis subconvergentibus, apice coalitis, vulgo non exsertis, ascis oblongis, sursum rotundato-truncatis, bifoveolatis, basi attenuato-stipitatis, octosporis, $80 \approx 17-18$; sporidiis distichis, ellipsoideis, 1-septatis, non v. vix constrictis, eguttulatis, hyalinis, $15-18 \approx 6$.

Hab. in ramulis delapsis *Populi nigræ*, Angleur Beaufays Belgii. *D. salicellæ* vicina.

13. **D. Chorostate pulchella** Sacc. et Br. Misc. Myc. I, pag. 3. — 2883

Pustulis gregariis, subcutaneo-erumpentibus, $1\frac{1}{3}-1\frac{1}{2}$ mm. diam.; peritheciis paucis, circinantibus, $\frac{1}{3}-\frac{1}{2}$ mm. diam., globulosis, ostiolis convergentibus, brevibus, non exertis; ascis fusoideis, subsessilibus, aparaphysatis, $60-76 \approx 12-16$, lumine apice minute bifoveolato, octosporis; sporidiis subdistichis, oblongo-fusoideis, constricto-1-septatis, $16-22 \approx 7-9$, utrinque acutiusculis, rectis curvulisve, hyalinis.

Hab. in ramis corticatis *Populi pyramidalis*, Troyes Galliæ (BRIARD). — *Melanconi dolosæ* affinis videtur.

14. **D. Chorostate Trecassum** Sacc. et Briard Misc. Myc. I, p. 4. 2884

— Peritheciis corticolis, in acervulos valseos, conico-discoideo, regulares arcte aggregatis, globosis, atris, $\frac{1}{3}$ mm. diam., stromate albido, ligneo exceptis circuloque nigro eximie circumscriptis, ostiolis in conum atrum, obtusiusculum, parum extans coalescentibus; ascis cylindraceis, breve stipitatis, apice lumine coarctato, rotundatis, p. sporif. 95-110 \approx 9-11, octosporis; sporidiis monostichis, ellipticis utrinque subrotundatis, constricto 1-septatis, 18-22 \approx 7-9, biguttatis granulosive, hyalinis.

Hab. in ramis corticatis Platani orientalis prope Troyes (Trecasses) Galliæ (BRIARD). — Ob sporidia elliptica et ascos tere-tes ad *D. fibrosam* nutat.

15. **D. Chorostate Rehmiana** Starb. Bidr. Sver. Ascom. pag. 5, tab. 2885 I, fig. 2. — Stromatibus sparsim gregariis, valseis, totos ramulos ambientibus, e basi lata ligno paullum innata zonaque nigra circumscripta disciformi-conoideis pustulatim primo corticem, deinde irregulariter fissam, elevantibus, ad basim 3-5 mm. diam.; peritheciis in singulo stromate 10-20-25, globosis vel mutata pressione angularibus, majusculis, ostiolis longioribus, cylindricis epidermidem, cujus laciniæ acervulis adnatæ, disciformiter erumpentibus; ascis tereti-clavatis, sessilibus, 60-70 \approx 7-9, octosporis; sporidiis ellipsoideis, e fusoideo ellipsoideis, 1-septatis, vix constrictis, subdistichis, 15-20 \approx 3,5-5,5, loculo inf. angustiore.

Hab. in ramulis Ulmi montanæ ad Köping Oelandiæ.-Subaf-finis *D. Saccardianæ*.

16. **D. Chorostate leucopis** (Fr.) Sacc. *Valsa leucopis* (Fr.) Quél. Jura 2886 506, t. IV, f. 9, *Sphaeria leucopis* Fr. S. M. II, 402. — Pustulata; peritheciis ovatis, erectis (inclinatis sec. Quél.), subquinis, intus albissimis, ostiolis conjunctis in discum minutum, globulo centrali albo; sporidiis (teste Quél.) lanceolatis, 4-guttatis, 20 μ . longis, utrinque aristatis.

Hab. in ramis Coryli (KUNZE), Ulmi et Coryli (QUÉLET), in Germania et Gallia. — Perithecia conferta, subcircinata, pauca, majuscula, tuberculo minuto a cortice interiori formato obvallata, dissecta cava albidissima; colla brevissima, apicibus in discum minutum inæqualem nigrum, globulum centrale minutum expellentibus.

17. **D. Chorostate Comptoniæ** Ell. et Ev. Proc. Phill. 1890, p. 234. 2887

— Peritheciis subcircinatis, pustulas minutas dein rimosas formantibus, cortice interiori immersis, $\frac{1}{5}$ mm. diam., 10-20 in quoque

- 706 Pyrenomyceteæ, Sphæriaceæ, Hyalodidymæ, Diaporthe.
 acervulo, globosis, ostiolis breve cylindraceis, obtusis, perforatis, vix emergentibus; nucleo albido, dein nigricante; ascis oblongis, 35-40 \approx 7; sporidiis biseriatis, oblongis, subinæquilateralibus, 1-septatis leniterque constrictis, 10-12 \approx 3-3,5, flavidо-hyalinis.
- Hab.* in ramis Comptoniæ asplenifoliæ, Newfield N. J. Amer. bor. — Spermogonia adsunt, multilocularia, griseo-atra, sporulis allantoideis minutis referta.
18. **D. Chorostate tecta** (Cooke) Sacc., *Valsa tecta* Cooke Grev. XI, 2888 p. 109. — Tecta, inconspicua; peritheciis circinatis, paucis, in cortice nidulantibus; ostiolis minimis, brevibus; ascis clavatis; sporidiis inordinatis, sublanceolatis, quadrinucleatis, demum uniseptatis, hyalinis, 18 \approx 5.
- Hab.* in cortice Myricæ, Darien Georgia Amer. bor. (RAVENEL).
19. **D. Chorostate carpinigera** (B. et C.) Sacc., *Diatrype carpinigera* 2889 B. et C. North Amer. Fungi n. 854. — Pustulis parvis, nigris; peritheciis abditis; sporidiis oblongis, uniseptatis, 14-15 μ . longis.
- Hab.* in ramis Carpini, Pennsylvania Amer. bor. (MICHERER).
20. **D. Chorostate Ellisii** Rehm in litt. Ellis et Ev. in Torr. of Bot. Club. 1883, p. 89. — Stromate corticali, non linea nigra circumscripto; peritheciis numerosis, 15-20 in quoque stromate, 0,5 mm. diam. stipatis, subcircinatis, corticem in pustulas parvas elevantibus, disco convexo, brunneo, margine ostiolis obtusis perforato, centrum pustularum occupante; ascis subclavatis, 57 \approx 8; sporidiis distichis v. stipatis, acute ellipsoideis, 4-nucleatis, subinde medio leniter constrictis, 11-15 \approx 3,5-5,5.
- Hab.* in ramis Carpini americanæ. — Adest simul *Cytospora* sporulis oblongis, hyalinis, 4-nucleatis, 8-11 \approx 2,5.
21. **D. Chorostate neglecta** (Duby) Berl. et Vogl. Add. Syll. p. 108, *Valsa* 2891 Duby in Herb. Currey, Cooke in Grevillea XIV, p. 47. — Stromate epidermide tecto, nigro nigracenteve, ovato aut globoso, in ramis junioribus nullo; peritheciis 6, compresso-ovatis, verticalibus, dense congestis, basi applanatis; ostiolis exsertis, fragilibus, aterrimis, cylindricis, plus minus elongatis, truncato-papillatis, interdum inflexis, sæpe lateralibus; ascis cylindricis, angustis; sporidiis ovoides, hyalinis, uniseriatis, nucleolatis, rarissime uniseptatis.
- Hab.* ad caules sicclos Genistæ tinctoriæ in Britannia.
22. **D. Chorostate sparsa** Niessl Contr. IV, Fl. Myc. Lus. p. 17. — 2892 Stromate late effuso, ambiente, toto immerso, diatrypeo, corticis interioris superficiem nigrificante, ligni substantiam non mutante, sed linea nigra limitato; peritheciis minutis, sparsis, in cortice in-

teriori, depresso globosis, ostiolo crasso, brevi, vix erumpente; ascis lanceolatis, sessilibus, 8-sporis, 52-60 \approx 8-10; sporidiis laxe distichis, oblongis, obtusiusculis, rectis, medio septatis, hyalinis, 10-12 \approx 3-4.

Hab. ad *Glycines violacea* ramulos aridos, Coimbra Lusitaniæ (MOLLER). — Adsunt spermogonia sparsa, sub peridermio immutato in corticis superficie nidulantia, minuta, ellipsoideo-globosa, ostiolo papilliformi, sporulis oblongis, plerumque rectis, 2-4-guttulatis, hyalinis, 12-15 \approx 4-5.

23. **D. Chorostate microcarpa** Rehm in Voss Myc. Carn. III, p. 173. 2893

— Stromate ambiente, corticem interiore nigrificante; peritheciis in acervulos valseos in cortice interiore nidulantes, circ. 8 monostiche congregatis, globosis, nigris, 0,3 mm. diam., ostiolis brevibus, in disco rotundo, plano, pallide subconice erumpente punctiformiter emergentibus; ascis fusoideis, apice rotundatis, 50 \approx 8, octosporis; sporidiis fusoideis, rectis, medio haud constrictis, 4-guttulatis, utroque apice brevissime filiformiter appendiculatis, hyalinis, 15 \approx 4.

Hab. ad ramos emortuos *Cytisi nigricantis* pr. Zirlach Carnioliae.

24. **D. Chorostate Hippophaës** Bomm. Rouss. Sacc. sp. nov. — Acer-

vulis compactis, subvalseis, sparsis; peritheciis globulosis, 4-10 aggregatis, rarius solitariis v. geminatis, corticolis, linea nigra circumscriptis et epidermide erupta cinctis, ostiolis longis, cylindraceis, emergentibus, rugulosis, saepe apice curvis, divergentibus; ascis cylindraceis vel subfusoideis, 48-64 \approx 9, octosporis; sporidiis biseriatis, cylindraceis v. fusoideis, obtusis, brevissime mucronatis, 1-septatis, leviter constrictis, 12-16 \approx 4-5,5, 4-guttatis, hyalinis, plerumque rectis.

Hab. in ramis emortuis *Hippophaës rhamnoidis*, Knocke Belgii. — Adest simul *Phoma consocia* B. R. S. (probabile spermogonium): peritheciis tectis, minutis; sporulis ellipsoideis v. ova-to-acuminatis, 2-guttatis, hyalinis, 6-8 \approx 2-3, basidiis hamatis.

25. **D. Chorostate magnispora** (E. et E.) Sacc. *Valsa magnispora* Ell. 2895

et Ev. Journ. Myc. 1887, p. 42. — Peritheciis cortice interiore immersis, ligum non penetrantibus nec linea atra circumscriptis, 6-10 acervulatis, globoso-ovatis, circ. $1/4$ mm. diam., supra in colla brevia fasciculatim erumpentia et parum emergentia contractis, ostiolis hemisphaericis, nitidis, nigris, levibus, pertusis et subumbilicatis; ascis subsessilibus, tereti-oblongis, 100-120 \approx 18-22; sporidiis

biseriatis, oblongo-fusoideis, hyalinis, 1-septatis, leniter constrictis, 25-35 = 9-11.

Hab. in ramis emortuis *Aceris*, Plainfield N. J. Amer. bor. (MESCHUTT). — A *D. Aceris* ascis sporidiisque multo majoribus differt.

26. ***D. Chorostate Berlesiana*** Sacc. et Roum. in Rev. Myc. 1883, 2896 p. 234, t. 39, fig. 4. — Pustulis gregariis, discoideis, subcutaneis, mediocribus, linea stromatica nulla v. obsoleta; peritheciis circinantibus, minutis, globosis, $\frac{1}{4}$ mm. diam.; ostiolis in disculum erumpentem, atrum conjunctis, obtusiusculis vix emergentibus; ascis fusoideo-clavatis, 50 = 8, apophysatis, octosporis; sporidiis distichis cylindraceis, utrinque obtusiusculis, 13-17 = 3,3-3,5, 1-septatis, non vel vix constrictis, muticis, 4-guttatis, hyalinis.

Hab. in ramis corticatis *Rhamni Frangulæ* in Arduennis (LIBERT). — A *D. syngenesia* sporidiis muticis, stromatibus haud concentrica striatis etc. differt.

27. ***D. Chorostate cercophora*** (Ellis) Sacc. *Valsa cercophora* Ellis 2897 Torr. Bot. Cl. 1882, p. 99. — Peritheciis paucis, 4-6 aggregatis, majusculis, stromate subcarbonaceo, linea nigra lignum penetrante circumscripto immersis, ostiolis obtusis, validis, late hiantibus, in discum conicum prominulum, erumpens conjunctis; ascis clavato-cylindraceis, 75-84 = 10-12; sporidiis distichis, oblongo-ellipsoideis, medio constrictis, utrinque appendiculatis, 10-15 = 3, hyalinis, pluriguttulatis.

Hab. in ramis emortuis *Ilicis opacæ*, New Jersey Amer. bor. — Appendiculæ, sporidiorum longitudinem æquantes, mox dilabuntur; superior apice curvula.

28. ***D. Chorostate Eucalypti*** Harkn. New Calif. Fungi p. 24. — Peritheciis in acervulos valsoideos, linea atra circumscriptis, aggregatis, rostris 1 mm. vel amplius longis, laxis, sæpe agglutinatis; ascis oblongo-fusoideis, 33 = 7; sporidiis fusiformibus, vix curvulis, hyalinis, 1-septatis, 4-guttatis, acuminatis, 15 = 4.

Hab. in foliis emortuis *Eucalypti Globuli*, S. Francisco in California.

29. ***D. Chorostate binoculata*** (Ellis) Sacc. *Valsa binoculata* Ellis 2899 Torr. Bot. Cl. 1882, p. 111. — Peritheciis 3-6 aggregatis, majusculis, stromate e cortice formato lineaque nigra, lignum non penetrante, circumscripto immersis; ostiolis erumpentibus, subglobosis, latiuscule hiantibus; ascis clavato-cylindraceis, 125-150 = 15-18; paraphysibus obtusis granulosis obvallatis, octosporis; sporidiis mono-

stichis, late ellipsoideis, subhyalinis, 1-septatis constrictisque, 2-nucleatis, 20-25 = 12-14.

Hab. in truncis emortuis *Magnoliae glaucæ*, New Jersey Amer. bor. — Ostiola subinde delitescentia et tunc fungi vix indicium.

30. **D. Chorostate æsculicola** (Cooke) Berl. et Vogl. Add. Syll. p. 105, 2900
Valsa Cooke in Grevillea XIV, p. 47. — Pustulis orbicularibus, in lineas breves dispositis; peritheciis subglobosis, atris, congestis, ostiolis cylindricis, convergentibus, levibus, emergentibus; ascis clavatis; sporidiis biserialibus, fusoideis, quadrinucleatis, demum uniseptatis, hyalinis, 20 = 4.

Hab. in ramulis *Æsculi*, Sydenham Britanniæ.

31. **D. Chorostate Hippocastani** (Cooke) Berl. et Vogl. Add. Syll. 2901 p. 105, *Valsa* Cooke in Grevillea XIII, p. 98. — Pustulis paucis; peritheciis (6-8) in cortice immersis et cuticulam e compositis ostiolis convergentibus perforatam elevantibus; peritheciis subglobosis, depresso-plerumque gregariis, atris; ascis clavatis, subsessilibus, 8-sporis; sporidiis fusiformibus, biseriatis, uniseptatis, biguttulatis, ad septa parum constrictis, aliquando utrinque appendicula hyalina præditis, hyalinis, 25-28 = 5-7.

Hab. in ramulis *Æsculi Hippocastani*, Kew, Bustey Park Hampton Court et Honnington Britann.

32. **D. Chorostate stictostoma** Ellis Amer. Nat. 1883, pag. 316 (D. 2902
punctostoma). — Stromate corticali, e cortice immutato formato; peritheciis 8-12, monostichis, $\frac{1}{2}$ mm. diam., ostiolis brevibus cylindraceis, apice in disculum olivaceum, leniter elevatum, $\frac{3}{4}$ mm. diam. junctis, disco erumpente, epidermide arcte cincto ostiolisque perforato; ascis clavato-cylindraceis, 55 = 8-9; sporidiis biseriatis oblongo-ellipticis, 4-nucleatis, medio leniter constrictis, hyalinis, 11-13 = 4-4,5.

Hab. in ramis emortuis *Amelanchieris canadensis*, Decorah Iowa Amer. bor. (E. W. HOLWAY).

33. **D. Chorostate tuberculosa** (Ellis) Sacc. *Valsa tuberculosa* Ellis 2903 Bull. Torr. Bot. Cl. 1882, p. 112. — Peritheciis 8-10 aggregatis, globosis, 350 μ . diam., stromate e cortice immutato nisi solidiore et nigro-circumscripto formato immersis, ostiolis brevibus cylindraceis, obtusis, irregulariter pertusis, fasciculatim erumpentibus; ascis late lanceolatis, 75 = 15, octosporis, filiformi-paraphysatis (?); sporidiis distichis, ellipsoideis, 2-nucleatis (1-septatisque?), 12-13 = 7,5.

Hab. in ramis emortuis *Amelanchieris canadensis*, New Jersey America borali.

34. **D. Chorostate cylindrospora** Peck 38 Rep. St. Mus. p. 104. — 2904

Stromatibus valsoideis, prominulis, erumpentibus sparsis; peritheciis numerosis, 15-30 in quoque stromate, stipatis, epidermide intus nigrescente tectis, ostiolis longis, stipatis, exsertis, fere æquilibus, nigris; ascis acutis, subfusiformibus, 45-55 \approx 6-8; sporidiis subcylindraceis, stipatis vel distichis, quadrinucleatis, 13-16 \approx 3-4.

Hab. in ramis emortuis *Pruni pensylvanicæ* in montibus Adirondack Amer. bor.

35. **D. Chorostate Cratægi** (Curr.) Fuck. — Sacc. Syll. I, pag. 204. 2905

Incertum an hoc spectet *Pseudovalsa* (*Valsaria*) *Cratægi* Winter, Cooke in Grevillea XIV, pog. 48. — Stromate valsoideo; sporidiis subuniserialibus, ellipticis, uniseptatis, pallide fuscis, 16-18 \approx 6.

Hab. in *Cratægo* in Germania.

36. **D. Chorostate Cydoniæ** Pass. Diagn. Fung. nnovi, III, n. 14. — 2906

Acervulis valseis, laxe vel crebre erumpentibus, pulvinatis, parenchymati corticali insidentibus; peritheciis subglobosis intus albidis, ostiolo obtuso vix emergentibus; ascis cylindraceo-clavatis, curvis, 8-sporis, 100-112 \approx 12; sporidiis oblique uniseriatis vel subdistichis, oblongo-ovatis, prope medium septatis, leniter constrictis, loculis inæqualibus haud guttulatis, altero angustiore, hyalinis, 17,5 \approx 7,5.

Hab. in ramulis *Cydoniæ*, sociâ *Phoma Cydoniæ*, Parma.

37. **D. Chorostate elæostroma** (Cooke) Berl. et Vogl. Add. Syll. p. 107, 2907

Valsa olivæstroma Cooke in Grevillea XIV, p. 48. — Erumpens; pustulis sub epidermide nidulantibus; peritheciis 6-10 in circulos dispositis, stromate olivaceo, pulveraceo insidentibus, collis subelongatis convergentibus; ascis clavatis, octosporis; sporidiis subuniserialibus, lanceolatis, medio uniseptatis, leniter constrictis, hyalinis, 26-28 \approx 8.

Hab. in ramis *Cerasi Avium*, Jedburgh Britanniæ:

38. **D. Chorostate columbiensis** Ell. et Ev. Proc. Phil. 1890, p. 233. 2908

— Peritheciis in acervulos subcircinantibus, cortice interiore immersis basique ligno insculptis, 3-6 in quoque acervulo, $3/4$ -1 mm. diam., infra collabentibus, in colla brevia ostiolo subquadrisulcato coronata erumpentia contractis, crusta tenui nigra, stromaticæ tectis, stromate corticali intus expallente, ellipsoideo, 2-5 mm. d.; zona atra circumscripta; ascis oblongo-lanceolatis, 100 \approx 12; spo-

ridiis biseriatis, oblongis, 20-22 \approx 7-8, constricto 1-septatis, 2-nucleatis, hyalinis.

Hab. in truncis emortuis arboris ignotæ, in Columbia britannica Amer. bor. (MACOUN).

39. **D. Chorostate elephantina** Cooke et Harkn. in Grevillea XIV, 2909 pag. 8. — Pustulis valsæformibus, rarissime dispersis; peritheciis cortice immersis vel ligno adnatis, globosis, atris, ostiolis tenuibus, elongatis, flexuosis, cylindraceis, emergentibus; ascis clavatis, octosporis; sporidiis lanceolatis quadrinucleatis, mox uniseptatis, hyalinis, 12 \approx 3,5.

Hab. in caulis Geranii in California (HARKNESS).

40. **D. Chorostate Gladioli** Ell. et Ev. in Journ. of Myc. 1886, p. 101. 2910 — Peritheciis matrice immersis, epidermide nigrescente tectis, maculas albas, definite limitatas, 2-3 mm. long., subinde confluentes et tum 1 cm. vel amplius longas occupantibus, circiter $\frac{1}{4}$ mm. diam., parcis, saepe 1-2 in quaque macula, subinde etiam 6-8, ostiolo setiformi donatis; ascis stipitatis, p. sp. circiter 40 \approx 6-7; sporidiis distichis, subfusoideis, 7-10 \approx 2,5, binucleatis, demum 1-septatis, hyalinis.

Hab. in caulis emortuis Gladioli, Louisiana Amer. bor. (LANGLOIS).

II. *Euporthe* Nits.

41. **D. Euporthe ilicina** Cooke Fungi Brit. 490, Grev. XVIII, p. 74. 2911 — Stromate ramulos cortice relaxato, in ligni superficie nigricante limitato, peridermio decolorato, linea nigra circumscripto; peritheciis globulosis, singulis vel gregariis, ostiolo leniter emergente; ascis subfusoideis; sporidiis lanceolatis, quadrinucleatis, dein uniseptatis, 16-18 \approx 3.

Hab. in ramis Ilicis Aquifolii in Britannia.

42. **D. Euporthe leucosarca** Ell. et Ev. Proc. Phil. 1890, p. 233. — 2912 Peritheciis dense sparsis, cortice immutato immersis et epidermide pustulatim bullata ostiolisque minutis papillatis perforata tectis; ascis oblongo-cylindraceis, 80-110 \approx 20-22, apophysatis (?); sporidiis biseriatis, late fusoideis, hyalinis, 1-septato-constrictis, 2-guttatis, 22 \approx 7-8, 1-septatis constrictisque, 2-nucleatis.

Hab. in ramis emortuis Carpinii americanæ, London Canada Amer. bor. (DEARNESS).

43. **D. Euporthe silvestris** Sacc. et Berl. Miscell. Myc. II, p. 27. 2913 n. 121, tab. VIII, fig. 7. — Peritheciis laxe gregariis, ligno semi-

infossis et peridermio tectis, linea angusta, endoxyla, atra, hinc inde circumscriptis, globulosis, $\frac{1}{2}$ mm. diam., ostiolis vix emergentibus; ascis tereti-fusoideis, 45-50 \approx 6-8; sporidiis octonis, distichis v. oblique monostichis, fusoideis, constricto 1-septatis, 14-15 \approx 5, 4-guttatis, hyalinis, utrinque initio minute hyalino-apiculatis.

Hab. in sarmenis *Vitis viniferæ silvestris*, Cervarese, Padova Italiæ borealis.

44. **D. Euporthe megalospora** Ell. et Ev. Proc. Phil. 1890, p. 235. — 2914

Peritheciis globosis, $\frac{1}{2}$ - $\frac{3}{4}$ mm. diam., sparsis, ligno intus expallente superficie atrato immersis, in ostiola elongata, rugulosa, 2-3 mm. long., subflexuosa abrupte contractis; ascis p. spor. 70-90 \approx 10-12; sporidiis distichis oblongo-fusoideis, leniter curvis, 1-septatis constrictisque, 2-4-nucleatis, utrinque acutis, 25-35 \approx 4,5-5,5.

Hab. in ligno emortuo *Sambuci canadensis*, Manchester Mass. Amer. hor. (W. C. STURGIS). — Angustior quam *Diap. leucosarca*.

45. **D. Euporthe hypoxylloides** Rehm Ascom. Ill. n. 874, — Stroma- 2915

tibus rotundis vel oblongis, saepe confluentibus, $\frac{1}{3}$ -3 cm. longis, $\frac{1}{2}$ -1 cm. latis, 3-4 mm. altis, hypoxylloides, intus suberoideis, fuscis, extus nigris; peritheciis stromate nidulantibus et circ. 1 mm. profunde immersis, solitariis, globosis, 0,3-0,4 mm. diam.; peritheciis globosis vel more Lophiostomatum plus minusve lineariter compresse protuberantibus, collis conoideis, obtusis, poro perspicuo pertusis; ascis fusoideis, teneris, 8-sporis, 60-66 \approx 7-8; sporidiis cylindraceis, obtusis, rectis, nucleis 2-4 magnis præditis, demum 2-cellularibus et medio vix constrictis, utrinque apice appendicula 3 μ . longa auctis, 12-15 \approx 3, distichis.

Hab. ad corticem interiore Aceris emortui in silva montis Arber Böhmerwald. — Habitu hypoxyleo dignoscenda species.

46. **D. Euporthe Chionanthi** Brun. Champ. Nouv. obs. env. Saint. 2916

II, p. 4. — Stromate parvo, oblongo vel elongato, in ramis decorticatis effuso, nigro superficiem maculante, linea atra in lignum penetrante limitato; peritheciis nigris, gregariis, ligno immersis, globulosis, ostiolo vix erumpente donatis; ascis fusoideis, 52-70 \approx 7-8, octosporis; sporidiis distichis, oblongo-fusoideis, 3-septatis, («an genuine?») non vel vix medio constrictis, hyalinis, 12 \approx 3.

Hab. in ramis emortuis *Chionanthi virginicæ*, Saintes Galliæ.

47. **D. Euporthe scobinoides** Schulz. et Sacc. in Rev. Myc. 1884, 2917 Schulz. Ill. Fung. Slav. n. 803. — Peritheciis laxe gregariis, ligno

subimmutato vel superficie leniter infuscato penitus immersis, linea stromatica nigra, per lignum sinuose excurrente, circumscriptis, globulosis 0,3-0,4 mm. diam., collo teretiusculo, in ostiolum erumpens, obtusiusculum desinente; ascis fusoideis, sessilibus, octosporis, 15 \approx 8; sporidiis distichis, oblongo-fusoideis, utrinque obtusiusculis 4-guttatis, medio-constrictis, 10-14 \approx 3, hyalinis.

Hab. in ramis dein decorticatis *Fraxini excelsioris* pr.

Vinkovce Slavoniæ

48. **D. Euporthe gorgonoidea** Cooke et Hark. in Grevillea XIII, 2918 p. 18. — Stromate effuso, cortice facile solubili tecto, ligni superficiem nigrificante v. crustam interruptam sistente, intus nigro-limitato; peritheciis globosis, ligno immersis, plerumque dense stipatis; ostiolis cylindricis, gracilibus, flexuosis, maxime elongatis; ascis cylindrico-clavatis, octosporis; sporidiis fusoideis, biseriatis, rectis, quadrinucleatis, demum 2-4-cellularibus, hyalinis, 15-17 \approx 3.

Hab. in ramis emortuis *Acaciæ* in California (HARKNESS). —

Diversa a *D. medusæa* et *D. fasciculata*.

49. **D. Euporthe ceuthosporioides** (Berk.) Sacc. Syll. I, pag. 645. — Forma **Lauri nobilis** Pass. in litt. ad Brunaud. — A typo differt 2919 stromate minus nitente, ceterum simillimo et piceo (forte initio rufescente ut mavult cl. Berkeley), ascis minoribus, 60 \approx 7-8, sporidiisque quoque minoribus, 11-13 \approx 3,5-4.

Hab. in foliis *Lauri nobilis*, Saintes Galliæ (BRUNAUD). —

In utraque forma sporidia sunt utrinque obtusiuscula, sed videntur mucronulo mucoso, peregrino aucta.

50. **D. Euporthe Malbranchei** Sacc. Syll. I, p. 361. — Var. **abscondita** Berl. Exc. Frioul p. 7, t. 9, f. 3. — A typo differt peritheciis pro- 2920 funde ligno immersis, 300-400 μ . diam.; sporidiis crassioribus, 12-14 \approx 5-7,2- raro 4-guttatis, utrinque crassiuscule appendiculatis.

Hab. in ramis decorticatis *Fici Caricæ*, Osoppo Ital. bor.

51. **D. Euporthe Peckii** Sacc. *D. sparsa* Peck 39 Rep. p. 52 (1886), nec Niessl (1883). — Peritheciis parcis, minutis, sparsis, ligno superficie atrato immersis; ascis clavatis v. subcylindricis, 75-85 \approx 7-10; sporidiis confertis, oblongis vel subfusoideis, hyalinis, ad septum constrictis, 4-nucleatis, 20-30 \approx 5-6,5.

Hab. in ramis *Rhois Toxicodendri*, Saugerties Amer. bor.

52. **D. Euporthe Orobanches** Berl. Exc. Frioul p. 8, t. 9, f. 2. — 2922 Stromate late effuso, maculas nigras magnitudine varias saepe formante vel epidermidem in nigrum tingente, intus linea atra circumscripto; peritheciis aggregatis, sub epidermide offuscata nidu-

lantibus, minutis, globulosis, caule immersis, in collum longum cylindraceum, sinuosum abeuntibus; ascis clavatis, 1-2-foveolatis, 70 \approx 7, 8-sporis; sporidiis subdistichis, oblongo-ovoideis, medio 1-septatis constrictisque, 4-guttulatis, 10-12 \approx 3, hyalinis.

Hab. in caulibus emortuis Orobanches, Osoppo agri Forojulensis Ital. bor.

53. **D. Euporthe Characiæ** H. Fab. Sphér. Vaucl. p. 45. — Stromate 2923
superficiali, late effuso, lignum denudatum fuligineo inquinante;
peritheciis sparsis, ligno immutato infossis, globosis, matricem leni-
ter inflantibus, $\frac{1}{2}$ mm. latis, ostiolo elongato, stricto, recto v. de-
flexo, corticem secedentem perforante; ascis lanceolatis, sessilibus,
8-sporis, 55-70 \approx 7-9; sporidiis distichis, hyalinis, oblongis, utrin-
que obtusiusculis, medio constrictis et tenuissime septatis, 4-gut-
tatis, 15-18 \approx 4.

Hab. ad basim putrescentem Euphorbiæ Characiæ, Vau-
cluse Galliæ.

54. **D. Euporthe Rubiæ** H. Fab. Sphér. Vaucl. p. 46. — Peritheciis 2924
sparsis aut parcis acervulatis, ligno immutato infossis, $\frac{1}{4}$ mm. la-
tis, ostiolo epidermidem erumpente, longe rostellato, flexuoso, dia-
metrum perithecii bis et quatuor superante; ascis lanceolatis,
sessilibus, 8-sporis, 50-54 \approx 7-8; sporidiis distichis, oblongo-lanceo-
latis, medio constrictis et subtiliter septatis, 4-guttatis, 15 \approx 4.

Hab. ad caules exsiccatos Rubiæ peregrinæ, Vaucluse
Galliæ.

55. **D. Euporthe hemicrypta** (Dur, et Mont.) Sacc. *Sphæria hemicry- 2925*
pta Dur. et Mont. Syll. Crypt. n. 837 et Fl. Alg. I, pag. 517. —
Peritheciis tectis simplicibus aggregatisque subcircinantibus, globo-
sis, atris, ligno semiimmersis collabentibus levibus, in ostiolum lon-
gum angustatis, ostiolis erumpentibus solitariis aut junctis cylin-
dricis subdivergentibus; ascis oblongo-subclavatis, 50 \approx 12-13; spo-
ridiis cymbiformibus, hyalinis, triseptatis (an potius 4-nucleatis?),
10 \approx 2,5.

Hab. in ramis Physalidis somniferæ, Rhois Coriariae
et Anagyridis fœtidæ, Alger (DURIEU).

56. **D. Euporthe discrepans** Sacc. Mich. II, pag. 568. — Peritheciis 2926
hinc inde dense gregariis, ligno superficie late infuscato v. deni-
grato immersis, globosis, $\frac{1}{3}$ mm. diam., ostiolis erumpentibus breve
rostellatis; ascis tereti-fusoideis, utrinque obtusiusculis, 65 \approx 8-9,
octosporis, apophysatis; sporidiis oblique monostichis v. distichis,

cylindraceis v. subfusoideis, utrinque obtusis, rectis curvulisve, medio lenissime constrictis, $10-12 \approx 3,5-4$, 4-guttatis, hyalinis.

Hab. in caulis R umicis emortuis, Newfield N. J. (ELLIS). — A *D. discorde*, cui affinis, ostiolis elongatis, sporidiis minoribus mox dignoscitur.

57. **D. Euporthe apiculosa** Ellis Torr. Bot. Cl. 1882, p. 19. — Peritheciis caule immersis, ostiolis obtusiusculis, prominentibus superficiemque asperulam reddentibus; ascis oblongis, p. sporif. $40 \approx 7-8$; sporidiis fusoideis, utrinque breve apiculatis, $8-10 \approx 4,5$.

Hab. in parte ima caulum Erigerontis canadensis, dejectorum, New Jersey Amer. bor.

58. **D. Euporthe immutabilis** Cooke et Harkn. in Grevillea XIV, pag. 9. — Stromate varie effuso interruptoque, caulis superficiem vix decolorante, linea nigra in ligno limitato; peritheciis sparsis globosis, ostiolis brevibus, punctiformibus; ascis clavatis, octosporis; sporidiis sublanceolatis, biserialibus, rectis curvulisve, 2-4-nucleatis, uniseptatis, hyalinis, $12-14 \approx 4$.

Hab. in caule Scrophulariæ in California (HARKNESS).

59. **D. Euporthe Tulasnei** Niessl. — Sacc. Syll. I, pag. 657, forma 2929 **Galegæ** Wint. Contr. Fl. Myc. Lus. V, pag. 18. — Differt a forma typica; ostiolis interdum valde elongatis, nodulosis, flexuosis; sporidiis parum crassioribus, semper medio constrictis.

Hab. in caulis putridis Galegæ officinalis cum spermogoniis, Coimbra in Lusitania (MOLLER). — An species distincta?

60. **D. Euporthe brachystoma** Sacc. et Malbr. Fung. Gall. ser. V, n. 2163. — Stromate effuso, epidermidem decolorante lineaque sinuosa, nigra circumscripto; peritheciis gregariis, cortice tectis lignoque adnatis, globoso-depressis, $\frac{1}{5}-\frac{1}{4}$ mm. diam., nigris; ostiolo brevissimo, vix emergente; ascis fusoideis, sessilibus, a paraphysatis, octosporis, $70 \approx 7$; sporidiis distichis, ovato-fusoideis, $12 \approx 3,5$, rectis, utrinque obtusiusculis, 4-guttatis, dein 1-septatis, hyalinis. — Spermogoniis *Phomam Dianthi* sistentibus, peritheciis in macula picea plerumque innatis, epidermide velatis, globoso-depressis; sporulis ovato-oblongis, $7 \approx 3$, biguttatis, hyalinis, stipitellis filiformibus, curvis, $10-12 \approx 1-2$ suffultis.

Hab. in caulis emortuis Dianthi barbati, Brionne Galliæ. — Hinc *Diaporthi intermediæ*, illinc *D. Tulasnei* affinis species.

61. **D. Euporthe Kellermanniana** Winter in Torr. bot. Club. 1885, 2931 p. 49. — Stromate nullo; peritheciis profunde immersis, depresso

globosis, membranaceis, atris, 210-260 μ . latis, ostiolo erumpente præditis; ascis oblongo-fusoideis, 8-sporis, 35-40 \approx 5-7; sporidiis oblongis, sæpe inæquilateralibus, utrinque rotundatis vel parum acutiusculis, medio uniseptatis, non constrictis, 4-guttulatis, 9-11 \approx 3-3,5.

Hab. ad culmos putridos Zeæ Maydis pr. Lexington Amer. bor. (W. A. KELLERMANN).

62. **D. Euporthe fuegiana** Speg. F. Fueg. n. 217. — Stromatibus plus 2932 minusve late effusis, in ramis decorticatis, maculas extus atras crустosas undulatas irregulariter lineares, 1-5 cm. long., 1-3 mm. lat., flexuosas intus albas linea nigra tenui limitatas efficientibes; peritheciis densiuscule constipatis v. subseriatis ligno immersis, contextu parenchymatico-sinuoso subindistincto, ostiolo carbonaceo elongato longiuscule exerto crassiusculo, recto v. flexuoso; ascis fusoideis, apice stricte rotundato subtruncatis, postice acutato-rotundatis, 60-65 \approx 10, octosporis, a paraphysatis; sporidiis oblique distichis, elliptico-elongatis, utrinque acutis ac strictissime rotundato-subtruncatis, 18-20 \approx 5-6, medio 1-septatis, non constrictis, loculis sæpe parce coarctatulis non v. grosse 2-guttulatis, hyalinis.

Hab. non rara in ramis emortuis Chilitrichi amelloidis per Fuegiam totam.

63. **D. Euporthe antarctica** Speg. F. Fueg. n. 215. — Stromatibus 2933 plus minusve late effusis, in ramis denudatis maculas extus irregulariter elongatis, strictis, 1-2 cm. long., 1-2 mm. lat., tumidulas atras subcrustosas, intus pallescentes v. albas, linea tenuissima candida limitatas efficientibus; peritheciis densiuscule constipatis, subseriatis ligno immersis, subprominulis, globulosis, 400-450 μ . diam., tenui-membranaceis, fuscis, contextu parenchymatico oblongo subimperspicuo, ostiolo minuto obtuse rotundatis, postice attenuato-acutatis, 40-60 \approx 10-14, octosporis, a paraphysatis; sporidiis oblique distichis, ellipticis, utrinque obtuse acutatis, 14-15 \approx 5-6, medio 1-septatis, lenissime constrictis, loculis non v. grosse 2-guttulatis, hyalinis.

Hab. ad ramos emortuos Pernettiæ mucronatae in dumetis in Staten island Fuegiæ.

II. *Tetrastaga* Nits.

64. **D. Tetrastaga didymelloides** Sacc. et Malbr. Mich. II, p. 594. — 2934 Peritheciis laxe v. dense gregariis cortice immersis, globoso-depressis, ostiolo minuto vix erumpente, linea stromatica nulla vel ob-

soleta; ascis fusoideo-clavulatis, subsessilibus, 55-60 \approx 7-8, octosporis, aparaphysatis; sporidiis distichis, fusoideis, 14-15 \approx 4,4-guttatis 1-septatis vix constrictis, hyalinis, rectiusculis.

Hab. in ramis corticatis *Lonicerae*, Elbeuf Galliæ (MALBRANCHE). — Stromate obsoleto ad *Didymellam* nutat, sed paraphyses nullæ et sporidia prorsus diaportheoidea.

65. **D. Tetrastaga incompta** Sacc. Mich. II, p. 565. — Peritheciis 2935 sparsis vel hinc inde subaggregatis, grandiusculis, cortice intus infuscato nidulantibus; ostiolis vix erumpentibus, punctiformibus; ascis tereti-fusoideis, 60 \approx 8-10, apice subtruncatis, deorsum attenuatis, octosporis, aparaphysatis; sporidiis distichis, fusoideis, 1-septatis, lenissime constrictis, 14-16 \approx 4, rectis curvulisve, 4-guttulatis, utrinque acutiusculis, hyalinis.

Hab. in ramis corticatis *Ampelopsis hederacea*, Ecully (Rhône) Galliæ (THERRY).

66. **D. Tetrastaga santonensis** Sacc. Fung. Gall. ser. V, n. 2162. 2936 — Stromate obsoleto; peritheciis epidermide translucida tectis, globoso-depressis, nigris, $\frac{1}{3}$ mm. diam.; ostiolo brevissimo, obtuso perforante; ascis fusoideis, sessilibus, aparaphysatis, 55-60 \approx 12, octosporis, apice minuto, 2-foveolato; sporidiis distichis vel oblique monostichis, fusoideis, inaequilateralibus, 1-septatis, leniter constrictis 14-18 \approx 5, 4- dein 2-guttatis, hyalinis, utrinque appendiculâ brevi, acuta, hyalina ornatis.

Hab. in cortice ramulorum *Salicis vitellinæ*, Saintes Galliæ (BRUNAUD). — Quum stroma sit indistinctum, ad *Gnomoniæ* genus nutat, cui fructificatione quoque accedit; at et cum *Diaporthe salicella* et *D. Spina* affinitas manifesta.

67. **D. Tetrastaga exasperans** Nitschke. — Sacc Syll. I, pag. 686. 2937
Var. **santonensis** Pass. in litt. Brun. in Rev. Myc. 1886, p. 205.
— Peritheciis corticolis, in stromate pustuliformi, 2-3-aggregatis, vix erumpentibus; ascis oblongo-claviformibus, sœpe basi incurvatis, attenuatis, paraphysibus gracilibus, longioribus cinctis, 50-72 \approx 8-5, octosporis; sporidiis distichis, fusoideis, 1-septatis, vix constrictis, hyalinis, loculis biguttulatis, obscure septatis, 15 \approx 5.

Hab. in ramulis emortuis *Betulae albæ*, Saintes Galliæ.

68. **D. Tetrastaga Terebinthi** H. Fab. Sphér. Vaucl. p. 46. — Stromate, sub epidermide immutata et secedente, corticem caulum fuligineo late inquinante; peritheciis cortici infossis, sparsis vel gregariis et matricem inflantibus, globosis, $\frac{1}{3}$ mm. latis; ostiolis nunc papillatis, nunc elongatis, cylindraceis, curvulis et fascicula-

tim erumpentibus; ascis lanceolatis, sessilibus, 8-sporis, 45-50 \approx 7; sporidiis lanceolatis, medio constrictis, et subtile septatis, subdistichis, 4-guttatis, hyalinis, 12 \approx 3-4.

Hab. ad basim caulinum emortuorum Pistaciæ Terebinthi, Vaucluse Galliæ.

69. **D. Tetrastaga dolosa** Sacc. et Roum. in Rev. Myc. 1883, tab. 2939
39, fig. 5. — Stromate vix ullo manifesto; peritheciis dense et æqualiter gregariis, cortice immersis, globulosis, vix $\frac{1}{2}$ mm. diam., intus griseis, ostiolo brevissimo, obtuso, perforante; ascis fusoideis, 60-70 \approx 10-12, octosporis; sporidiis distichis, fusoideis, utrinque obtusiusculis, medio constricto-1-septatis, 4-guttatis, hyalinis, 10-12 \approx 4.

Hab. in ramis junioribus Robiniae Pseudacaciae in Ardennis (LIBERT). — A *D. oncostoma* diversissima.

70. **D. Tetrastaga punctulata** Sacc. Bomm. Rouss. Mycol. Belg. II, 2940
p. 11. — Peritheciis globulosis, sparsis, 250-260 μ . diam., epidermide atrata tectis, stromate corticali nigro-limitato immersis, ostiolo erumpenti, punctiformi; ascis tereti-clavatis, 60-70 \approx 7-8, subsessilibus, 8-sporis; sporidiis distichis, ellipsoideo-fusoideis, obtusis, 1-septatis, constrictis, 4-guttulatis, 12-15 \approx 4-4,5, hyalinis.

Hab. in ramis Tecomæ radicantis, Parc de Tervueren Belgii. — A *D. spissa* differt ostiolis minus exsertis, ascis sporidiisque paullo majoribus.

71. **D. Tetrastaga priva** Sacc. et Roum. in Rev. Myc. 1884. — Stromate effuso sed vix manifesto lineaque nigra privo; peritheciis gregariis cortice immersis, globosis, $\frac{1}{3}$ mm. diam., ostiolo erumpente sed vix exstante; ascis fusoideis, aparaphysatis, 65 \approx 8, octosporis; sporidiis distichis, fusoideis, rectis, utrinque obtuse attenuatis, 16-18 \approx 4,5-5, medio constricto-1-septatis, 4-guttatis, hyalinis.

Hab. in ramis corticatis Fraxini Orni in Ardennis (LIBERT). — A ceteris speciebus fraxinicolis stromate obsoleto, linea nigra deficiente, sporidiis ratione majusculis mox dignoscitur.

72. **D. Tetrastaga interrupta** Niessl Contr. Fl. Myc. Ius. p. 17. — 2942
Stromate maculiformi, diatrypeo, toto immerso, haud expanso, atropurpureo, interdum expallente, linea atra limitato; peritheciis paucis, in singulo stromate saepe solitariis, in corticis parenchymate nidalantibus, depresso globosis; ostiolo minimo abbreviato, vix erumpente; ascis lanceolato-clavatis, stipite brevi, 8-sporis, 50-60 \approx 6-7; sporidiis laxe distichis, fusiformibus, acutiusculis, rectis sed pa-

rum inæquilateralibus, medio septatis, haud constrictis, 4-guttulatis, hyalinis, $9-11 \approx 2-3$.

Hab. ad ramulos emortuos *Tecoma* radicantis, in horto botanico, Coimbra Lusitaniæ. — A *Diaporthe Lebiseyi* differt: stromatibus exiguis, peritheciis erostellatis, ascis magis elongatis.

73. **D. Tetrastaga Vepris** (de Lacr.) Fuck. — Syll. I, p. 667. — Var. 2943 **Rosarum** Pass. in Brun. Champ. nouv. VI, pag. 1. — Peritheciis cortice interiore immersis, globulosis, nigris, ostiolis punctiformibus; ascis subfusoideis, $25-30 \approx 8$; sporidiis distichis, fusoideis, 1-septatis, non constrictis, rectis, $10 \approx 2,5$, obscure guttatis, hyalinis.

Hab. in sarmenis emortuis Rosæ caninæ, Saintes Galliæ. — Ad *D. Vepris* var. *appendiculatum* Rehm — Syll. I, p. 668 — spectare videtur *Sphæria dichætospora* Ripart Bull. Soc. bot. France 1876, p. 225.

74. **D. Tetrastaga Neilliæ** Peck 39 Rep. p. 52, t. II, f. 6-9. — Peritheciis numerosis, 0,5-0,6 mm. diam., late, laxe et inæqualiter aggregatis, cortice interiore immersis lignoque saepe leviter insculptis, epidermide ostiolis conicis v. rostellatis nigris perforata tectis, infra subinde concavis; ascis subcylindricis, $60 \approx 7-10$; sporidiis confertis v. biseriatis, oblongis v. subfusoideis, ad septum leniter constrictis, 2-4-nucleatis, $13-16 \approx 5-6$.

Hab. in ramulis emortuis Spirææ (Neilliæ) opulifoliæ, Albany Amer. bor. — Perithecia raro subvalseo-aggregata.

75. **D. Tetrastaga oligocarpoides** Rehm Ascom. Ill. n. 875. — Stroma latissime effuso, ambiente, linea atra tenui, profunde in ligno immersa cincto, peridermii et corticis colore haud mutato; peritheciis 6-8 grægarie in interiore cortice nidulantibus, globosis, stroma valseum haud æmulantibus sed peridermum longitudinaliter dirumpentibus, 6-8 plerumque lineariter seriatis, collis plerumque brevibus, rarissime usque 2 mm. longis et flexuosis, haud perspicue pertusis, nigris prominentibus; ascis fusoideis, teneris, 8-sporis, $45-50 \approx 6-7$; sporidiis elliptico-fusoideis, rectis, haud appendiculatis, 2-cellularibus, utraque cellula 1-2 nucleata, medio saepe subconstricta, hyalina, $9-12 \approx 3$, distichis.

Hab. ad ramos siccos Rosæ caninæ in Belgio (Mad. BOMMER et ROUSSEAU). — A *D. incarcerated* ascis sporidiisque minoribus recedit.

76. **D. Tetrastaga magellanica** Speg. F. Pat. n. 129. — Stromate effuso, obsoleto, corticis interioris et ligni superficiem nigrificante; peritheciis laxe grægariis v. sparsis cortice nidulantibus, vix tumi-

dulis, lenticulari-globosis, 300-400 μ . diam., ostiolis brevissimis, subconoideis epidermidem perforantibus sed vix exsertis; ascis fusoides, 55-60 \approx 5-6, apice subtruncatis, inferne acutatis, a paraphysatis, octosporis; sporidiis distichis, elliptico-elongatis, 10-14 \approx 2,5-3, utrinque obtusiusculis, subinæquilateralibus, medio 1-septato-constrictis grosse 1-guttulatis, hyalinis.

Hab. ad ramos emortuos Ribis magellanici prope Cabo Negro Patagoniæ. — *D. miti* peraffinis, sporidiis tamen minoribus obtusioribusque recedens; an tantum ejusdem varietas?.

77. **D. Tetrastaga Winteri** Speg. F. Fueg. n. 214. — Stromatibus 2947 parum manifestis late per corticem excurrentibus; peritheciis hinc inde sparsis v. plus minusve dense aggregatis peridermio non v. vix fisso ac pustulato-elevato tectis, globulosis, 300-350 μ . d., coriaceo-membranaceis atris glabris, contextu indistincto opaco, ostiolo carbonaceo vix papillulato; ascis fusoideo-clavatis, antice obtuse rotundatis vel subtruncatis crasse tunicatis bifoveolatis, postice attenuatis acutato-rotundatis sessilibus, 60-65 \approx 8-9, octosporis a paraphysatis; sporidiis oblique mono- v. distichis ellipticis non v. subovatis utrinque obtuse rotundatis, 12-14 \approx 3,5-5 rectis v. leniter inæquilateralibus medio 1-septatis constrictulis, loculis (supero non v. vix crassiore) granuloso-farctis sæpe minute biguttulatis, hyalinis.

Hab. ad corticem ramorum emortuorum Drymidis Winteri in Staten island Fuegiæ.

78. **D. Tetrastaga densissima** Ellis Amer. Nat. 1883, p. 316. — Peritheciis minutis, $\frac{1}{4}$ mm. diam., nigris, cortice interiore immersis, sæpius 15-30 congregatis; ostiolis brevibus, subulatis, leniter convergentibus, nec cohærentibus, peridermum mox decadens perforantibus; ascis clavato-cylindraceis, 40-45 \approx 5-6; sporidiis biseriatis, fusiformibus, 4-nucleatis, 11-15 \approx 2-2,5, initio utrinque minute setaceo-appendiculatis.

Hab. in ramis Quercus coccineæ, Newfield New Jersey Amer. bor. — Linea stromatica nigra desideratur.

79. **D. Tetrastaga Conradi** Ellis Amer. Nat. 1883, p. 316. — Peritheciis sparsis, minutis, depresso-sphæricis, epidermide tectis, ligno non immersis lineaque nigra haud circumscriptis; ostiolis cylindraceis, rectis, rugulosis, nigris, apice abrupte acutatis; ascis subcylindraceis, 35-40 \approx 6-7, octosporis; sporidiis biseriatis, ovato-ellipticis, 1-septatis, hyalinis, vix constrictis, 6-8 \approx 2 $\frac{1}{2}$ -3.

Hab. in ramis emortuis Corematis Conradi, Willow Grove, New Jersey Amer. bor.

80. **D. Tetrastaga digitifera** Mout. Not. Ascom. p. 3. — Stromate 2950
sub epidermide non mutata ambiente, corticis superficiem nigrificante et in ligno linea nigra limitato; peritheciis cortice immersis, subglobosis, $\frac{1}{2}$ mm. diam., collis subtriplo longioribus, erumpentibus a basi leviter attenuatis, curvulis; ascis clavatis, 8-sporis, apice truncatis, 110 ± 15 ; sporidiis tereti-oblongis, medio septatis et vix constrictis, minute guttulatis, $25-35 \pm 8.5$.

Hab. in ramis Sarothamni scoparii prope Liège Belgii (MOUTON). — Var. **lignicola**: peritheciis ligno superficie atrato leviter immersis, ostiolis modo crassiusculis, rugulosis, modo elongatis decumbentibus. In ramis decorticatis Sarothamni scoparii, Poix Galliae (M. ROUSSEAU).

81. **D. Tetrastaga Mori** Berl. F. moric. f. V, n. 9, t. 25, f. 1-6. — 2951
Stromate late effuso, epidermidem leviter nigrificante in cortice v. in ligno nigro-limitato; peritheciis solitariis vel 3-4 coacervatis, cortice immersis globoso-conoideis, minutis $180-200 \mu$. diam., subinde subtus cupulatis, ostiolo brevissimo obtuso; ascis clavatis, brevissime stipitatis, apice 1-2 foveolatis, $36-40 \pm 7$, octosporis; sporidiis irregulariter distichis, oblongis, utrinque rotundatis, muticis, 4-5-guttulatis, primo continuis, medio constrictis, demum 1-septatis, 10 ± 3 , hyalinis.

Hab. in ramulis corticatis emortuis Mori albæ, Padova in Italia boreali.

82. **D. Tetrastaga delitescens** Bomm. Rouss. Sacc. sp. nov. — Pe- 2952
ritheciis corticolis, peridermio tectis, globulosis, nunc 5-15 subcircinatibus stromateque albo-furfuraceo linea nigra cincto exceptis, nunc inæqualiter sparsis et stromate subnullo; ostiolis convergentibus; ascis clavatis, $45-57 \pm 9-10$, p. sporif., octosporis, subsessilibus, apophysatis; sporidiis distichis, ellipsoideis mucronulatis, 1-septatis, constrictis, $12-15 \pm 4-5$.

Hab. in ramis Liriodendri Tulipiferæ, Tervueren Belgii.

— Extus vix visibilis. Ad *Chorostaten* nutat.

83. **D. Tetrastaga prominula** Bomm. Rouss. Sacc. Mycol. Belg. II, 2953
p. 11. — Peritheciis globulosis, cortice non atrato immersis, sparsis vel 2-8 aggregatis, pustulas minutis corticem vix perforantes formantibus; ostiolo tereti-conoideo, non exerto; ascis tereti-clavatis, $57-70 \pm 12-15$, octosporis; sporidiis distichis, tereti-fusoideis, obtusis, rectis curvulis, 1-septatis, constrictis, utrinque ciliatis, $17-18 \pm 5-6$, 4-guttulatis, hyalinis.

Hab. in ramulis junioribus Myricæ Gales, Westmalle Belgii.

84. **D. Tetrastaga Ligustri** Allesch. Suelbay. Pilze p. 194, t. II, f. 3. 2954

— Stromate sub epidermide late effuso, linea nigra limitato, ligni superficie tandem atrata; peritheciis subsparsis peridermio immersis vel ligno quoque insculptis, globoso-ellipsoideis, nigricantibus; ostiolis elongatis, basi incrassatis, rectis curvulisve, saepe fasciculatis; ascis clavatis, stipitatis, 8-sporis, 45-50 μ . long.; sporidiis distichis fusoideis, obtusulis, 1-septatis vix constrictis, 4-guttulatis, hyalinis, 12-14 \approx 3.

Hab. in ramis siccis *Ligustri vulgaris* pr. München Bavariæ. — Affinis *D. crassicollis*.

85. **D. Tetrastaga Baccharidis** Cooke. — Syll. I, n. 692. — Var. 2955
gallica Brun. Champ. Saint. VII, p. 2. — Stromate minuto subovato, epidermidem infuscante, linea nigra circumscripto; peritheciis globulosis, tectis; ostiolo brevi tantum erumpente; ascis fusoideis, 45-60 \approx 6-7; sporidiis subdistichis fusoideis, rectis curvulisve, 4-guttulatis, 1-septatis, non constrictis, 12-15 \approx 3-3,5, hyalinis.

Hab. in ramis tenellis emortuis *Baccharidis halimifoliae* Fouras Galliæ.

86. **D. Tetrastaga Asclepiadis** Ell. et Ev. in Bull. Torr. bot. Club. 2956
1883, p. 98. — Stromatibus e maculis nigris superficiem caulum occupantibus formatis, 0,5 cm. longis v. confluentibus, intus linea nigra limitatis; peritheciis sparsis, globosis, 0,25 mm. diam. in substantia caulis non omnino immersis; ostiolis obtusiusculis, cylindraceis, crassis, 250-300 μ . longis; ascis 35-40 \approx 9, p. sp.; sporidiis distichis, ellipsoideis, medio constrictis, 3-4-nucleatis, subobtusis, 10-12 \approx 3-4.

Hab. in caulinis emortuis *Asclepiadis tuberosæ*, Newfield Americæ borealis.

87. **D. Tetrastaga Lupini** Harkn. New Calif. Fungi p. 24. — Peritheciis aggregatis; ostiolis brevibus crassiusculis; ascis delicatis, fusoideis, 55-60 \approx 9.; sporidiis fusoideis, 1-septatis, 4-guttatis, leniter constrictis; loculis saepius inæqualibus, mox septo divisis, 15 \approx 4.

Hab. in ramis *Lupini arborei*, San Francisco Californiæ.

88. **D. Tetrastaga Geranii** Cooke et Harkn. in Grevillea XIV, p. 8. 2958
— Peritheciis gregariis, cortice immersis, saepe nigro-limitatis, subglobosis, atris, initio tectis, dein rimoso-fissis; ostiolis brevibus, emergentibus; ascis clavatis; sporidiis lanceolatis, quadrinucleatis, hyalinis, 15-16 \approx 4.

Hab. in caulinis *Geranii* in California (HARKNESS).

89. **D. Tetrastaga Therryana** Sacc. et Penz. Mich. II, p. 593. — 2959

Stromate late effuso epidermidemque longe lateque denigrante lineaque nigra infra limitato; peritheciis gregariis cortice immersis, globulosis, mediocribus, nigris; ostiolis erumpentibus, filiformibus, flexuosis, caulem setulosum efficientibus; ascis fusoideis, 36-46 \times 6-7, apophysatis, octosporis; sporidiis distichis fusoideis, 10-11 \times 2,5-3, 4-guttatis, obsolete 1-septatis, hyalinis.

Hab. in caule Hellebori fœtidi, St. Bonnet le froid (Rhône) Galliæ (THERRY).

90. **D. Tetrastaga mucronulata** Sacc. Mich. II, pag. 568. — Perithe-

ciis cortice superficie infuscato immersis lignoque adnatis, seriatim gregariis, globulosis, $\frac{1}{4}$ mm. diam., ostiolo breve rostellato, emergente; linea stromatica nulla manifesta; ascis fusoideis, utrinque obtusiusculis, 60 \times 10, apophysatis, octosporis; sporidiis distichis v. oblique monostichis, crassiuscule fusoideis, rectis vel curvulis, utrinque acutis atque mucronulatis, 14-15 \times 6, medio septulatis leviterque constrictis, 4-guttatis, hyalinis.

Hab. in caulis emortuis Asteris, Newfield New Jersey Amer. bor. (ELLIS).

Sect. 5. Phæodidymæ.

PHÆOSPHÆRELLA Karst. Symb. Myc. Fenn. XXVI, p. 28 (Etym. *phæos* fucus et *Sphærella*). — Characteres *Sphærellæ* sed sporidia colorata.

1. **Phæosphærella maculosa** (Sacc.) Karst. l. c., *Sphærella macularis* Karst. Myc. Fenn. II, p. 174, *Sphærella maculosa* Sacc. Syll. I, p. 487. — Sporidia demum flavo-fuliginea, 14-24 \times 5-7.

TICHOThECIUM Flotow in Wint. Hedw. 1886, p. 15 (Etym. *teichos* paries et *thece* ascus, quia sec. Flot. asci parietales). — Perithecia plus minusve immersa, deum emergentia, minuta, corneo-carbonacea, atra, poro simplici pertusa. Paraphyses mucosæ, interdum indistinctæ v. nullæ. Asci 4-8- vel polyspori. Sporidia bi-vel quadricellularia, fusca. — Meo sensu sectiones, in quas genus dividitur, genera distincta sistunt.

I. Asci 4-8-spori; sporidia uniseptata.

1. **Ticthothecium gelidarium** (Mudd.) Berl. et Vogl. Add. Syll. p. 118, 2961 *Sphæria gelidaria* Mudd Manual of Brit. Lichens p. 130. — Pe-

ritheciis nigris, maculiformibus, sparsis, nucleo albo farctis; paraphysibus distinctis, laxis, flexuosis mucosisque; ascis brevibus, cylindraceis, 4-sporis, ad utramque extremitatem sporidiorum constrictis; sporidiis subrotundis vel late oblongis, irregularibus, protoplasmate obscure brunneo v. fuligineo farctis, indistincte bilocularibus, 10 \approx 6.

Hab. in thallo *Squamariæ* gelidæ in Britannia.

2. **Tichothecium perpusillum** (Nylander) Arnold Lichenol. Fragm. XVI, 2962 pag. 27 et in Wint. Hedw. 1886, pag 17, *Endococcus perpusillus* Nyl. Exposit. Pyrenoc. p. 64. — Peritheciis minutis, sessilibus, fere globosis, ostiolo haud impresso, indistincte pertuso, atris, opacis, scabris, 0,1-2 mm. diam.; ascis elongato-clavatis vel fusiformibus, 8-sporis; sporidiis ellipticis, medio haud constrictis, sæpe parum curvatis, utrinque attenuatis rotundatisque, dilute fuscis, 15-21 \approx 6-7.

Hab. in *Aspicilia cinerea* et *A. gibbosa* in Germania et Britannia. — Teste Cl. Leighton, Lichen. of Great Brit. Edit. III, pag. 496, perithecia sunt minutissima, immersa, depresso globosa, atra; sporidia octona, oblonga, fusca, bicellularia, 14-19 \approx 6-7.

3. **Tichothecium Stigma** Körber Parerga Lichenol. p. 468 et Wint. in 2963 Hedw. 1886, pag. 16. — Peritheciis minutissimis, punctiformibus, immersis, demum parum prominulis, atris, opacis, ostiolo indistincto, supra thallum dispersis; ascis clavatis vel fusiformi-clavatis, 8-sporis; sporidiis ellipticis, utrinque acutiusculis, medio sæpe subconstrictis, bicellularibus, interdum cellula superiori parum majori, fuligineis vel dilute rufo-fuscis, 14-20 \approx 5-6.

Hab. in thallo *Rhizocarpi geographici* et *Psoræ lamprophoræ* in Germania.

4. **Tichothecium macrosporum** Hepp in litt. sec. Arnold Lichenol. 2964 Ausfl. III, p. 960 et Wint. in Hedw. 1886, p. 17. — Peritheciis initio omnino immersis, demum vertice prominulis, globosis, atris, poro distinto pertusis, circ. $\frac{1}{10}$ mm. diam., plerumque gregariis; ascis oblongo-clavatis, deorsum longe attenuatis, 8-sporis, 60-65 \approx 17-21; sporidiis subdistichis, elliptico-oblongis, utrinque attenuato-rotundatis, uniseptatis, haud v. vix constrictis, fuscis, 16-24 \approx 6-8.

Hab. in thallo *Rhizocarpi geographici* et *R. alpicoli* in Germania.

5. **Tichothecium Arnoldi** (Hepp) Körber Par. Lichenol. p. 469, Wint. 2965 Hedw. 1886, pag. 16. — Peritheciis minutissimis, punctiformibus, confertissime stipatis, immersis, globosis, ostiolo indistincto, atris,

substantia molliuscula, viridi-brunnea farctis; ascis breviter clavatis; sporidiis 6-3, soleæformibus, bicellularibus, cellula superiore majore, viridi-fuscis, 8-11 \approx 3-4.

Hab. in thallo Urceolariæ scruposæ in Germania.

6. **Tichothecium calcaricolum** (Mudd) Arnold Lichenolog. Ausfl. XI, 2966 p. 521 et Wint. in Hedw. 1886, p. 16, *Microthelia calcaricola* Mudd Manual p. 306. — Peritheciis media magnitudine irregulariter sparsis, sessilibus vel basi immersis, globosis, poro submagni pertusis, atris; ascis clavato-subventricosis, 8-sporis; sporidiis elliptico-oblongis, bicellularibus, fuscis, 19-20 \approx 6; paraphysibus gelatinosis.

Hab. in thallo Aspiciliæ calcareæ, A. cinereæ et A. gibbosæ in Germania et Britannia. — Sec. Arnold sporidia 15-18 μ . tantum longa sunt. — Var. **Sendtneri** Arnold Lichenol. Ausfl. XI, p. 37: peritheciis hemisphæricis, prominulis, atris; sporidiis utrinque obtusis, fuscis, bicellularibus, 15-18 \approx 6-8, octonis. In thallo Polyblastiæ Sendtneri in Germania.

7. **Tichothecium gemmiferum** (Tayl.) Körb. Parerga Lichen. p. 468, 2967 et Wint. in Hedw. 1886, pag. 16. *Verrucaria gemmifera* Taylor in Mackay, Flora hibern. II, p. 95, *Microthelia propinqua* Körb. System. p. 374. — Peritheciis sparsis, immersis, minutissimis, globosis, vertice umbilicato, demum collapso, poro simplici pertusis, opacè atris, 0,1 mm. diam.; ascis clavatis, crasse tunicatis, 8-sporis, 30-36 \approx 10-12; sporidiis ellipticis vel elliptico-oblongis, obtusis bicellularibus, plerumque haud vel perparum constrictis, fuscis, 8-12 \approx 3-4; paraphysibus coalitis.

Hab. in thallo crustaceo lichenum variorum in Germania.

8. **Tichothecium Complanatæ** (Arnold) Wint. in Hedw. 1886, p. 17, 2968 *Endococcus Complanatæ* Arnold Lichenol. Ausfl. in Tirol. V, p. 13. — Peritheciis minutissimis, punctiformibus, parum prominulis, atris; sporidiis octonis, oblongis, fusiformibus, utrinque plus minus attenuatis, rectis vel parum curvatis, bicellularibus, fuscis, 15-17 \approx 4-5; paraphysibus nullis.

Hab. in thallo Lecanoræ complanatae in Germania.

9. **Tichothecium squamarioides** (Mudd) Wint. in Hedw. 1886, p. 17, 2969 *Sphæria squamarioides* Mudd Manual of Brit. Lich. p. 130, *Polyccum squamarioides* Arnold Lichenol. Fragm. XVI, p. 29. — Peritheciis aggregatis, acervulos minutos formantibus, punctiformibus, atris, nucleo albo-cinereo; ascis elongato-oblongis vel oblongo-subclavatis, 8-sporis; paraphysibus indistincte diffluenti-

bus, mucosis; sporidiis obtuse fusiformibus vel fusiformibus, bicellularibus, interdum medio subconstrictis, obscure olivaceis v. subatratis, 8,5-11 \approx 3-4.

Hab. in thallo *Placodii gelidi* in Germania et Britannia.

II. Asci polyspori; sporidia uniseptata.

10. **Tichothecium nanellum** (Ohlert) Arnold, Lichenol. Fragm. XVI, 2970

p. 27 in Flora 1874 et Wint. in Hedw. 1886, p. 17, *Endococcus nanellus* Ohlert Zusamn. Lich. Preuss. pag. 44. — Peritheciis nigro-fuscis, globosis, poro pertusis, 90-105 μ . diam., inter squamulas thallinas et tomentum *Stereocauli tomentosi* nascentibus; ascis clavatis, apice acuminatis, polysporis, 30 \approx 12; sporidiis oblongis, utrinque acutis, continuis vel bicellularibus, fuscis.

Hab. in *Stereocaulo tomentoso* in Germania.

11. **Tichothecium cerinarium** (Mudd) Berl. et Vogl. Add. Syll. p. 120, 2971

Sphæria cerinaria Mudd Man. of Brit. Lich. p. 136. — Peritheciis minutis, punctiformibus, semiimmersis, atris, solitariis, 10-12, disco apotheciorum insidentibus, nucleo mucoso, pallide farctis; paraphysibus nullis; ascis non visis; sporidiis numerosis, oblongis, uniseptatis, medio leniter constrictis, pallide luteis, 4 \approx 2.

Hab. parasitice in apotheciis *Callopismatis cerini* in Britannia.

12. **Tichothecium erraticum** Mass. Symmicta Lich. p. 94 et Wint. in 2972

Hedw. 1886, p. 15, *Endococcus erraticus* Nyland. Lichenes Scandinaviae p. 283. — Peritheciis semiimmersis, globosis, ostiolo indistincto, impresso, nitide atris, circ. 0,05 mm. diam.; ascis initio oblongis, subcylindrico-clavulatis, demum in parte inferiori ventricosis, fere obovoideis, polysporis; sporidiis late ellipticis v. subglobosis, medio haud constrictis, bicellularibus, fuscis, 3,5 \approx 2,5.

Hab. in thallo crustaceo *Lichenum variorum* in Germania.

13. **Tichothecium pygmæum** Körb. Parerga Lichen. p. 467 et Wint. 2973

in Hedw. 1886, p. 15. — Peritheciis sparsis, totis v. pro maxima parte immersis, plerumque vertice parum impresso, pertuso tantum emergentibus, globosis, atris, subnitidis, 0,1 mm. diam.; ascis late clavatis, saepe parum ventricosis, deorsum stipitiformibus, attenuatis, vertice late rotundatis, polysporis, 47-52 \approx 12-14; sporidiis inordinatis, ellipticis, interdum subconstrictis, bicellularibus, fuscis, 6-8 \approx 3-3,5.

Hab. in thallo *Lichenum variorum saxicolarum* in Germania.

— Var. **ventosicola** (Mudd) Wint. in Hedw. 1886, p. 15, *Microthe-*

lia ventosicola Mudd Manual. p. 307: vix diversa a forma normali; peritheciis semiimmersis; sporidiis 8-9 = 5. In Hæmatommate ventoso parasitans in Germania. — Var. **grandiuscula** Arnold Lichen. Ausfl. V, p. 523 et Wint. in Hedw. 1886, p. 15: peritheciis 2-usque fere 3-plo majoribus quam in forma normali; sporidiis 6-9 = 3-4. In speciebus variis Lecideæ, Lecanoræ etc. in Germania. — Var. **microcarpa** Arnold Lichen. Ausfl. XIV, p. 477: peritheciis, ascis sporidiisque minoribus; sporidiis 20 in quoque asco, 5-6 = 2-3. In fructibus Callopismatis aurantiaci in Germania. — Var. **ecatonspora** Anzi Neosymb. p. 16 et Wint. in Hedw. 1886, p. 15: peritheciis numerosis, gregariis; sporidiis 9 = 3. Inter apothecia Rinodinæ Bischoffii in Germania.

III. Asci 4-8-spori; sporidia triseptata.

14. **Tichothecium leucomelarium** (Mudd) Berl. et Vogl. Add. Syll. 2974 p. 121, *Sphæria leucomelia* Mudd Man. of Brit. Lich. p. 105. — Peritheciis minutis, semiimmersis, subsphæroideis; ostiolis leniter depresso, minutis vix prominulis, nucleo pallido; paraphysibus gracilibus, filiformibus; ascis elongato-cylindraceis, medio inflatulis, 4-sporis; sporidiis oblongis vel obtuse fusiformibus, 3-septato-constrictis, primo pallide brunneis, dein obscure brunneis, 18 = 5.

Hab. parasitice in thallo Borreræ leucomelæ et B. ciliaris in Britannia.

15. **Tichothecium rimosicolum** (Leight.) Arnold in Flora 1861, p. 678, 2975 *Xenosphæria rimosicola* Körb. Parerg. Lich. p. 467. — Peritheciis numerosis, sparsis confertis, plus minusve immersis, fere globosis, vertice appanato, ostiolo impresso pertusoque, 0,2-3 mm. diam.; ascis ovato-clavatis vel subventricosis, 8-sporis; paraphysibus indistinctis; sporidiis ellipticis vel elliptico-oblongis, plerumque utrinque acutatis, 4-cellularibus, ad septa subconstrictis, dilute fuscidulis, 13-16 = 5-6.

Hab. in thallo Rhizocarpi subconcentrici, R. petræi, Siegortiæ calcareæ variorumque lichenum aliorum in Germania.

16. **Tichothecium decolorans** Rehm Ascom. Diagnos. n. 490 et Wint. 2976 in Hedw. 1886, pag. 18, *Phæospora decolorans* Rehm Exsicc. n. 490. — Peritheciis gregariis, thalli superficie plus minusve dealbata, insidentibus, punctiformibus, conoideis, atris, glabris; ascis clavatis, 8-sporis, 60 = 8; sporidiis ellipticis, obtusis, 2-4-cellula-

728 Pyrenomyceteæ, Sphæriaceæ, Phæodidymæ, Didymosphaeria.
ribus, cellulis mediis guttulis magnis præditis, ad septa constrictis,
rectis, 12 ± 4 ; paraphysibus nullis.

Hab. in thallo Biatoræ decolorantis in Germania.

17. **Tichothecium peregrinum** (Flotow) Wint. in Hedw. 1886, p. 18, 2977
Verrucaria peregrina Flotow in Botan. Zeitg. 1885, n. 8, *Phæo-*
spora peregrina Arnold in Lichenol. Ausfl. XX, p. 355. — Peritheciis punctiformibus, atris, thallo immersis, vertice tantum pro-
minulis; ascis oblongis, 8-sporis; sporidiis elongato-oblongis, 4-cel-
lularibus, ad septa parum constrictis, sæpe guttulis 4-præditis, fu-
scis, 15 ± 5 ; paraphysibus nullis.

Hab. in thallo Lecanoræ Bockii in Germania.

SOROTHELIA Körb. Parerga Lichenol. p. 471 et Wint. in Hedwigia
1886, pag. 10 (Etym. a *soros* et *thele* papilla). — Perithecia ag-
gregata, soros rotundatos vel irregulares, maculiformes forman-
tia, superficialia, carbonacea, atra; ascis 8-spori, paraphysibus ob-
vallati; sporidia didyma, fusca.

1. **Sorothelia confluens** Körber. Par. Lich. p. 471, Wint. in Hedw. 2978
1886, p. 10. — Peritheciis minutis, sessilibus, globosis v. hemisphæri-
cis, aterrimis, conspurcatis, mox lata basi in soros irregulares con-
fluentibus indeque deformibus; sporidiis in ascis cylindraceo-clava-
tis, octonis, mediocribus, soleæformibus, biguttulatis, $15-25 \pm 5-6$,
fuscis.

Hab. in thallo Phlyctidis argenæ in Silesia.

DIDYMOSENTHIA Fuck. — Sacc. Syll. I, p. 701.

I. Herbicolæ.

1. **Didymosphaeria conoidella** Sacc. et Berl. Misc. Myc. II, n. 26. — 2979
Peritheciis laxe gregariis, epidermide secedente nudatis e globoso
conicis, atris, sæpe concentrice, obsolete plicatis, $\frac{1}{3}-\frac{1}{2}$ mm. diam.,
ostio acutiusculo; ascis clavulatis, modice stipitatis, paraphysa-
tis, octosporis, $80-90 \pm 12$; sporidiis distichis, ellipsoideo-oblongis,
constricto-1-septatis, $15-18 \pm 6-7$, fuligineis.

Hab. in ramulis emortuis Capparidis sarmentosæ, Teviot
Australiæ. — A *Did. conoidea* sporidiis duplo fere longioribus
mox distinguenda.

2. **Didymosphaeria subconoidea** Bomm. Rouss. Sacc. sp. nov. — Pe- 2980
ritheciis subcutaneis, dein erumpentibus, $150-170 \mu$. diam., sub-
carbonaceis, ostio teretiusculo; ascis cylindraceis, $75-100 \pm 6-7$,
octosporis; paraphysibus tortuosis, simplicibus vel ramosis, crebre

septatis, 3,5-4,5 μ . cr., ascum æquantibus; sporidiis monostichis, subfusoideis, olivaceis, 1-septatis, constrictis, 4-guttatis, olivaceis, 9-13 \approx 3,5, loculo sup. crassiore.

Hab. in caulibus emortuis *Digitalis purpureæ*, Poix Gal-liæ. — *D. conoidea* subaffinis.

3. **Didymosphaeria Lycii** (Kalchbr.) Sacc. *Amphisphaeria Lycii* Kalchbr. 2981
Szep. Gomb. II, p. 273, tab. V, f. I, C. — Peritheciis gregariis, epi-
dermide tectis eamque inflantibus, membranaceis, globosis depla-
natis, astomis, dein poro simplici vertice lacerato dehiscentibus; ascis
tenellis, facile diffluentibus, cylindraceis, paraphysatis, octosporis;
sporidiis oblique monostichis, oblongo-ellipsoideis, didymo-constri-
ctis, fuligineis. — St. conid: *Sporodesmium Lycii* Niessl. Pycni-
dium: *Hendersonia Lycii* Hazs.

Hab. in ramulis *Lycii* in Banatu Hungariæ.

4. **Didymosphaeria massarioides** Sacc. et Brunaud Mich. II, pag. 592 2982
— Peritheciis laxe gregariis, subinde duobus, confluentibus, globoso-
depressis, $1\frac{1}{8}$ - $3\frac{1}{4}$ mm. diam., cortice tectis; ostiolo obtuso crassiu-
sculo vix erumpente; contextu perithecii crassiusculo; ascis cylin-
draceis, apice rotundatis, breve stipitatis, 120 \approx 10, copiose filifor-
mi-paraphysatis, octosporis; sporidiis, oblique monostichis, ovato-
oblongis, utrinque obtusiusculis, 20 \approx 8, constricto-1-septatis, flavo-
ochraceis, dein fuligineis, sæpe 2-guttatis.

Hab. in ramulis corticatis *Lycii barbari*, Saintes Galliæ (BRUNAUD).

5. **Didymosphaeria fenestrans** (Duby) Winter in Hedwigia 1886, p.
24, Huc sec. cl. Winter ducendæ sunt *Didymella Epilobii*, *Gno-
monia Epilobii* et *Gnomonia fenestrans*. Syll. I, p. 556, 561, 562.

6. **Didymosphaeria innumerabilis** Winter Grevill. XV, p. 88. — Peri- 2983
theciis in maculis magnis arescendo-candicantibus v. pallide fu-
ligineis, determinatis, angulato-rotundatis seu irregularibus, mar-
gine luteolo angusto et area fuscidula latiori cinctis, usque 12 mm.
latis gregariis, hypophyllis sæpe confertis v. aggregatis, subsuper-
ficialibus, subglobosis, poro simplici pertusis, membranaceis, atris,
98-110 μ . diam., ascis e basi latiori ovata sursum attenuatis ses-
silibus, 8-sporis, 35-40 \approx 11,5-12,5; paraphysibus tenuissimis, filifor-
mibus plerumque mucosis; sporidiis congregatis, bacillari-subcla-
vatis, interdum parum curvatis, medio septatis leniterque constri-
ctis, plerumque utrinque, semper autem deorsum subattenuatis,
rotundatis, hyalinis, loculo superiori parum latiori, 16-18 \approx 4,5.

Hab. in foliis vivis Passifloræ cujusdam in prov. S. Catharina Brasiliæ (ULE).

7. **Didymosphaeria Matthiolæ** (Fuck.) Berl. et Vogl. Add. Syll. p. 114, 2984

Pleospora Fuck. in Reis. nach Nordpolarm. III, et Oud. Contr. Fl. Myc. Now. Seml. p. 152, t. II, fig. 7. — Peritheciis sparsis, punctiformibus, epidermide tectis, atris; ostiolo conico, obtuso, prominulo, atro; ascis maturis ovatis seu subglobosis, $30 \approx 24$, immaturis basi contractis, superiore parte ovato-ventricosis, $55 \approx 24$, octosporis; sporidiis stipatis, utrinque obtusis, antice parum obtusioribus crassioribusque, rectis, uniseptatis, $20 \approx 8$, demum pallide fuscis.

Hab. in foliis Matthiolæ nudicaulis, Nowaja Semlja.

8. **Didymosphaeria anomala** (E. et E.) Sacc. *Leptosphaeria* E. et C. 2985

Journ. Myc. 1887 p. 117. — Peritheciis gregariis, membranaceis, laxe parenchymaticis, $\frac{1}{3}$ mm. diam., nigris, levibus, subsphaericis, demum supra subcollabentibus, initio epidermide tectis, dein erumpentibus; ascis oblongo-cylindraceis, circ. $100 \approx 20$, subsessilibus; paraphysibus filiformibus, evanescentibus; sporidiis distichis, latis, oblongo-fusoideis, 1-septatis, inaequilateris et subincurvis, pallide stramineis, constrictis $30-35 \approx 10-12$, utrisque obtusis.

Hab. in caulis emortuis, sociâ *Cylindrocolla diffluente* in Scofield, Utah Amer. bor. (S. J. HARKNESS).

II. Arboricolæ.

9. **Didymosphaeria populina** Vuill. Malad. Peopl. p. 3. — Peritheciis 2986 peridermio innatis, sphæricis, $\frac{1}{5}$ mm. d., membranaceis, apice pertusis non papillatis; ascis elongatis, $85 \approx 28$, brevi pedicellatis; paraphysibus gracilibus ramosis; sporidiis inaequaliter distichis, didymis, profunde constrictis, $22 \approx 14$, pallide brunneis, loculo sup. majore.

Hab. ad ramulos Populi pyramidalis, Lorraine. — Adsunt perithecia pycnidica subepidermica (*Phoma populina* sp. nov.) sporulis ellipsoideis, hyalinis $5-6 \approx 2-2,5$, 1-guttatis.

10. **Didymosphaeria nana** Rost. Fungi Groenl. p. 557. — Peritheciis 2987 gregariis, minutis; ascis cylindraceo-clavatis, $64 \approx 12$; sporidiis distichis, oblongis, 1-septatis, fuscis, $25 \approx 7-8$.

Hab. in foliis siccis Betulae nanæ, Fiskefjord Groenlandiae.

11. **Didymosphaeria denudata** Ell. et Gall. Journ. Myc. 1890, p. 32. 2988

— Peritheciis sparsis, ovatis, suberumpentibus, minutis, $\frac{1}{4}$ mm. d., vertice emergente ruguloso, nigro, papillato, contextu densiu-

scule membranaceo; ascis cylindraceis, 50 ± 7 , in basim stipitiformem abrupte contractis; sporidiis monostichis, ellipsoideis, 1-septatis, brunneis, $6-7 \pm 4$.

Hab. in cortice epidermide orbato ramorum *Quercus* emortuæ Newfield N. J. Amer. bor. — A *Did. Cupula* peritheciis haud collabentibus, sporidiis minoribus etc. diversa.

12. **Didymosphaeria accedens** Sacc. in Fairm. Contr. Myc. West. N. 2989 Y. p. 48, t. IV, f. 12. — Peritheciis gregariis, tectis, $\frac{1}{4}-\frac{1}{3}$ mm. lat., ostiolo breviter papillato, erumpente, nucleo primo albo; ascis tereti-clavatis, 120 ± 10 ; paraphysibus obsoletis; sporidiis senis octonisve monostichis, ellipsoideis, utrinque rotundatis, 1-septatis, leviter constrictis, $20-22 \pm 11$, fuligineis.

Hab. in ramis emortuis arborum quarumdam (*Fraxini?*), Lyndonville N. Y. Amer. bor. — Affinis *D. Rhamni* et *D. incarcerata*, ostiolo ascisque diversa.

13. **Didymosphaeria buxina** Passer. Diagn. Fung. Nov. IV, p. 4. — Peritheciis sparsis, tectis, per corticem fissum erumpentibus, atris; ascis paraphysatis, clavatis, 8-sporis, $60-75 \pm 15-17$; sporidiis distichis v. oblique monostichis, ovato-oblongis, medio septatis, vehemente constrictis, utrinque subacutis, loculis inæqualibus ample guttulatis, primo luteis, tandem fuscis, $17,5-20 \pm 7,5-8$.

Hab. in ramulis emortuis *Buxi semperfirantis*, Vigheffio, pr. Parma Ital.

14. **Didymosphaeria producta** Sacc. et Paol. in Sacc. Myc. Sib. p. 13, 2991 f. 10. — Peritheciis sparsis gregariisve, initio epidermide velatis, mox subsuperficialibus, e globoso lageniformibus, atris, $\frac{1}{3}$ mm. diam., in ostiolum perithecium subæquans v. brevius, cylindraceo-conoides, apice obtusiusculum productis; ascis fusoideo-clavatis. utrinque obtusiusculis, subsessilibus, $65-70 \pm 9-11$; paraphysibus filiformibus, copiose obvallatis; sporidiis oblongo-fusoideis, inæquilateralibus, utrinque obtusulis, medio constricto-uniseptatis, dilute olivaceis, 4-guttatis, $22-26 \pm 5-6$.

Hab. in caule *Lonicerae cœruleæ* in silvis pr. fluv. Chabyk Sibiria (MARTIANOFF). — Ad genus *Rhynchostoma* vergit.

15. **Didymosphaeria atro-grisea** Cooke et Peck, Grevill. XVII, p. 92. — 2992 Peritheciis sparsis, convexis, cortice immersis, cuticula grisea tectis, demum ostiolo atro erumpentibus; ascis cylindraceis 8-sporis; sporidiis monostichis, ellipticis, uniseptatis, fuscis 15 ± 8 .

Hab. in cortice *Quercus albæ*, New York (PECK) et Poughkeepsie Amer. bor. (GERARD) — « An *Amphisphaeria*? ».

16. **Didymosphaeria scabella** (Quél.) Berl. et Vogl. Add. Syll. p. 342, 2993

Sphæria Quél. 13 suppl. p. 9, t. VIII, f. 24. — Peritheciis corneis, lenticularibus, 400-500 μ . diam., mammiformibus, granulosis v. vix muriculatis, nigris; ostiolo conoideo, umbilicato, vix prominulo; sporidiis lanceolatis, 30 μ . longis.

Hab. in ramis denudatis putrescentibus *Hederæ*, Jura.

17. **Didymosphaeria endoleuca** Passer. Diagn. F. N. III, n. 15. — Peritheciis solitariis, sparsis, atris, sub epidermide pustulatim inflata, integra v. vix fissa nidulantibus, nucleo candido; ascis elongato-clavatis, basi breviter attenuato-stipitatis, paraphysatis, 8-sporis 100-125 \approx 15; sporidiis amygdaloideis v. ellipticis, medio septatis, non constrictis, obscure fuscis, 20-27,5 \approx 8-10.

Hab. in ramis siccis *Cercidis Siliquastri* sociâ *Diplodia Siliquastri* in Horto botanico Parmensi Ital. (PASSEPINI).

18. **Didymosphaeria Ceanothi** (Cooke et Hark.) Berl. et Vogl. Add. Syll. 2995

p. 136, *Sphæria* Cooke et Harkn. in Grevillea XIII, p. 19. — Tecta, sparsa; peritheciis globoso-applanatis, atris, breviter papillatis; ascis amplis, clavatis, octosporis; sporidiis ellipticis, uniseptatis, medio constrictis, atro-fuscis, 35 \approx 15.

Hab. in sarmentis *Ceanothi* in California (HARKNESS),

19. **Didymosphaeria Genistæ** Fuck. — Sacc. Syll. I. p. 705, Winter

in Hedwigia 1886, p. 23. — Ascii 43-48 \approx 12; sporidia 14-16 \approx 4-5.

20. **Didymosphaeria sarmenti** (Cooke et Hark.) Berl. et Vogl. Add. Syll. 2996

p. 113, *Sphæria* Cooke et Harkness in Grevillea XIII, p. 19. — Sparsa; peritheciis tectis, subglobosis, atris, levibus, demum applanatis; ostiolo brevi punctiformi; ascis cylindrico-clavatis; sporidiis ellipticis, uniseptatis, medio non constrictis, fuscis, 12 \approx 5,

Hab. in sarmentis *Vitis* in California (HARKNESS).

21. **Didymosphaeria Persicæ** Pass. in Brun. Champ. Saint. V. p. 3. — 2997

Peritheciis sparsis, conicis, nigris, semper tectis, ostiolo punctiformi, corticem vix perforante; ascis cylindraceis elongatis, 115-125 \approx 8, paraphysatis; sporidiis 1-stichis, oblongo-ellipsoideis, medio 1-septatis, non v. parum constrictis, brunneis, 17,5 \approx 5-6.

Hab. in ramis emortuis *Persicæ vulgaris*, Saintes Galliæ.

22. **Didymosphaeria rubicola** Berl. Pug. F. Fior. p. 11 t. II, f. 5. — 2998

Peritheciis dense sparsis v. subgregariis, epidermide non nigrificata tectis, ostiolo minute eam perforante prominulis, globosis globoso-ve-conoideis, intus nucleo primo candido, dein fusco-atro farctis, 250-300 μ . diam., carbonaceo-membranaceis; ascis cylindraceis, rectis v. curvulis, subsessilibus, crasse tunicatis, paraphysibus [de-

mum coalitis cinctis, 100-120 = 14-16; sporidiis octonis, irregulariter monostichis, ovoideis, utrinque rotundatis, raro acutiusculis, 1-septatis, non v. lenissime constrictis, 20-23 = 10-12, rufo-fuligineis.

Hab. in sarmentis emortuis Rubi fruticosi pr. Firenze Italæ. *D. permutteræ* et *D. gregariæ* affinis.

23. **Didymosphaeria Hakeæ** Winter. Contr. Fl. Myc. Lus. V, p. 14. — Pe. 2999
ritheciis gregariis, sine macula, immersis, depresso-globosis, ostiolo punctiformi, prominulo instructis, membranaceis, fusco-atris, 0,25 mm. diam.; ascis oblongo-cylindraceis, utrinque attenuatis, in stipitem brevem productis, 8-sporis, 55-75 = 9-11; sporidiis distichis, oblongis, supra medium uniseptatis et profunde constrictis, inæquilateralibus, utrinque acuminatis, binucleatis, fuligineis 15-17 = 4,5-5; paraphysibus filiformibus.

Hab. ad Hakeæ salignæ folia arida rarissime prope Coimbra Lusitaniæ (MOLLER).

24. **Didymosphaeria Banksiæ** Cooke Grev. XIX, p. 90. — Epiphylla; 3000
maculis orbicularibus, pallidis, margine nebuloso-brunneo; peritheciis parcis, centralibus, nigris, cuticula diffracta erumpentibus, subglobosis, papillatis; ascis clavatis, sessilibus, octosporis; sporidiis biseriatis, 1-septatis, loculo sup. subgloboso, infero angustiore, brunneis, 10 = 5.

Hab. in foliis vivis Banksiæ, Victoria Australiæ (MARTIN).

25. **Didymosphaeria phyllogena** Winter in Journ. of Myc. 1885, pag. 3001
121. — Peritheciis in macula rotundato-angulata vel irregulari, fusca, lineâ atrâ cincta, determinata, usque 8 mm. diam., hypophylla immersis, epidermidem perforantibus, punctiformibus, tenuissime membranaceis, 100 μ . diam.; ascis clavatis, brevissime pedicellatis, 8-sporis, 50-70 = 9-11; sporidiis ditischis (rarius submonostichis), oblongis, inæquilateralibus, utrinque attenuatis, bicellaribus, ad septum constrictis, fuscidulis, 14-16 = 5; paraphysibus crassis, cylindraceis, septatis.

Hab. ad folia delapsa Liriodendri Tulipiferæ, Missouri.

26. **Didymosphaeria Dryadis** (Fuck.) Berl. et Vogl. Add. Syll. p. 114, 3002
Pleospora Fuck. Die 2.^o Deutsche Nordpolarf. II, 93. Oudem. Contr. Fl. Myc. Now. Seml. p. 152, t. II, fig. 6. — Peritheciis in foliorum aridorum nondum delapsorum pagina superiori sparsis, punctiformibus, globoso-conicis, atris, demum subliberis, ostiolis perforatis; ascis oblongo-ovatis, oblique stipitatis, tunica crassa præditis, octosporis, 88 = 32; sporidiis subdistichis, oblongis, utrinque

obtusis, uniseptatis, ad septum constrictis, loculis uniguttulatis, 30-32 \approx 12, primo hyalinis, demum flavo-fuscis.

Hab. in foliis exsiccatis Dryadis octopetalæ, Nowaja Semlja Maris glacialis.

27. **Didymosphaeria circinata** (Fuck.) Wint. in Hedwigia 1886, p. 23.

Gnomoniella? circinata Sacc. Syll. I, pag. 416. — Sporidia conglobata, oblonga, bicellularia, viridula, 9-10 \approx 3.

28. **Didymosphaeria circinans** Harkn. New Calif. Fungi p. 24. — 3003

Hypophylla; peritheciis gregariis in maculis circularibus, 1-6 mm. diam., tectis, circa ostiola matricem maculantibus; ascis octosporis, cylindraceis, sessilibus, 66 \approx 6, paraphysibus filiformibus cinctis; sporidiis ovoideis vel oblongis, uniseptatis, leniter constrictis, oliveo-brunneis, 5-7 \approx 3.4.

Hab. in foliis Eucalypti Globuli in California.

29. **Didymosphaeria Cupula** Ellis Amer. Nat. 1883, p. 313, *Sphæria Cu-* 3004

pula Ell. et Ev. in Bull. bot. Club. 1883; p. 90. — Peritheciis sparsis innatis, epidermide nigrescente, ostiolo papilliformi excepto, tectis, primum convexo-hemisphæricis dein siccitate collapsi; ascis 75 \approx 7; sporidiis ellipsoideis, brunneis, uniseptatis, monostichis, 9,5-11,5 \approx 4,5-5.

Hab. in foliis adhuc pendulis et in ramulis quercinis, Newfield Amer. bor.

30. **Didymosphaeria euryasca** Ell. et Gall. Journ. Myc. 1889 p. 67. 3005

— Peritheciis sparsis, partim erumpentibus, partim epidermide tectis, minutis, 80-100 μ . diam. supra perforatis...?; ascis inæquilateraliiter ovoideis, sessilibus, 35-40 \approx 12-15; paraphysibus; sporidiis 2-3-stichis, ovato-oblongis, 1-septatis, constrictis, utrinque rotundatis, brunneis, 12-15 \approx 3,5-5.

Hab. in foliis emortuis Pini Murrayanæ, in culmine Mt. Helena, Lewis and Clarke County, Montana Amer. bor.

III. In Monocotyledoneis.

31. **Didymosphaeria serrulata** E. et M. in Journ. of Bot. 1885, p. 99. 3006

— Maculis dealbatis; peritheciis sparsis vel gregariis, epidermide nigrescente, demum ostiolo brevi, obtuso, erumpente perforata, tectis, lenticularibus $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam.; ascis cylindricis, paraphysibus filiformibus obvallatis, 100-112 \approx 10-12; sporidiis monostichis, hyalinis, primum 3-4-nucleatis dein brunneis et 1-septatis, 18-20 \approx 5-6, initio muco obvolutis.

Hab. in petiolis *Sabal*is serrulatæ, maculas pallidas effor-
mans in Amer. bor.

32. **Didymosphaeria longipes** Trabut in Add. p. 411. — Peritheciis spar- 3007
sis, epidermide circa ostiolum nigrescente tectis, prominulis, globoso-
conoideis, atris, uitentibus, majusculis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., vertice
albo-punctatis, poro pertusis; ascis cylindraceis, longissime pedi-
cellatis, apice rotundatis foveolatisque, pedicello subinde curvulo
vel flexuoso, 80-100 μ . longo, paraphysatis, octosporis, 180-200 \approx
12; sporidiis recte vel oblique monostichis, subinde transverse dis-
positis, ovoideis, septo constrictis, loculo superiore sæpe crassiore
rotundatoque, inferiore vero subinde minore, subconoideo, fuligi-
neis, 14-18 \approx 8-9.

Hab. in *Agave americana* in Algeria (TRABUT).

33. **Didymosphaeria Spatharum** Wint. in Hedwigia 1885. — Perithe- 3008
ciis sparsis, immersis, depresso-globosis, ostiolo late conoideo, ob-
tuso præditis, erumpentibus, atris, membranaceis, matricem tegen-
tem parum nigrificantibus, 240-270 μ . latis; ascis cylindraceis, deor-
sum in stipitem brevem attenuatis, 8-sporis, 45-52 \approx 7; sporidiis
oblique mono-vel partim subdistichis, oblongis, utrinque rotundatis,
medio vel supra medium uniseptatis, non vel perparum constrictis,
fuligineis, 9-10 \approx 3,5-4; paraphysibus gracilibus, filiformibus.

Hab. in spathis aridis Bobartiae spathaceaæ, in monte
Tabulari Promont. Bon. Spei (MAC-OWAN).

34. **Didymosphaeria Typhæ** Peck 38 Rep. Stat. Mus. p. 104. — Pe- 3009
ritheciis minutis, punctiformibus, subglobosis, epidermide ostiolis
minute papillatis perforata tectis; ascis cylindraceis, 62-85 \approx 8-10;
sporidiis oblongis vel ellipticis, monostichis uniseptatis, non vel
vix septo constrictis, 10-15 \approx 5-6, coloratis; paraphysibus filifor-
mibus.

Hab. ad basim foliorum emortuorum *Typhæ latifoliæ*, Guil-
derland Amer. bor.

35. **Didymosphaeria typhina** Pass. in Brun. Nouv. Fragm. p. 29 (D. 3010
Typhæ). — Peritheciis sparsis, epidermide non nigrificata tectis,
lenticularibus, nigris, ostiolo v. prominulo v. depresso; ascis cylin-
draceis basi subattenuatis 8-sporis, 60-65 \approx 5-6, copiose paraphy-
satis; sporidiis monostichis, medio 1-septatis, castaneo-brunneis,
7-8 \approx 5.

Hab. in caulis emortuis *Typhæ angustifoliæ* in Ro-
chefort pr. Saintes Galliæ. — «An diversa a præcedente?»

36. **Didymosphaeria crastophila** Niessl in Rehm Ascom. n. 644, Hedw. 3011

1883, pag. 84. — Peritheciis mediocribus, dispersis, interdum subgregariis et confluentibus, strato firmo, clypeiformi, fusco-nigro, nitido, tectis, non erumpentibus, fere globosis, coriaceis; ascis cylindraceis, 8-sporis, $90 \approx 7$; sporidiis obtuse ellipticis medio saepe subconstrictis, 2-cellularibus, utraque cellula nucleo 1 parvulo instructis, rarius cellulis inæqualibus, fuscidulis, $9 \approx 5$ monostichis; paraphysibus tenerrimis.

Hab. ad culmos sicclos Poœ nemoralis prope Brünn Moraviae (NIESSL).

37. **Didymosphaeria Andropogonis** Ell. et Lang. Proc. Phil. 1890 p. 235. 3012

— Stromate e culmi substantia immutata sed pallidiore formato, 3-4 cm. long., 1 cm. lat., linea virescenti-nigra, profunde penetrante cincto, æqualiter ardosiaco-nigro; peritheciis sparsis, subglobosis, $\frac{1}{2}$ - $\frac{3}{4}$, mm. diam., ostiolo erumpente et convexo-discoideo excepto, omnino immersis; ascis cylindraceis circ. $110 \approx 8$ -10; paraphysibus validis sed evanescentibus; sporidiis 1-stichis, oblongo-cylindraceis utrinque rotundatis, subincurvis, 1-septatis hyalinis dein brunneis, 18-22 ≈ 4.5 -5.5.

Hab. in culmis Andropogonis muricati St. Martinsville Louisiana Amer. bor. (LANGLOIS).

38. **Didymosphaeria pardalina** Ell. et Ev. in Journ. of Myc. 1886, 3013

p. 102. — Peritheciis 4-6 gregariis, maculis suborbicularibus, vel ellipsoideis, atris, 2-4 mm. diam., insidentibus vel confluentibus, matrice omnino immersis, $\frac{1}{2}$ mm. diam., coriaceis, ostiolo minuto, punctiformi donatis, epidermide non elevantibus; ascis cylindraceis, 150-200 \approx 12-15; sporidiis monostichis, oblongo-cylindraceis, olivaceo-brunneis, 1-septato-constrictis, vix curvis, utrinque obtusis, 22-30 \approx 8-10.

Hab. in culmis emortuis Spartinae polystachyæ in America boreali (LANGLOIS).

39. **Didymosphaeria arundinicola** Bizz. Fung. Ven. novi v. crit. p. 2, 3014

t. II, f. 2. — Peritheciis minutissimis, subsuperficialibus, ostiolatis; ascis oblongo-clavatis, 100-110 \approx 21-23, octosporis; sporidiis bilocularibus, non vel vix constrictis, oblongo-ellipsoideis, utrinque hyalino-apiculatis, 26-28 \approx 12 (sine apiculo), initio hyalinis, biguttatis, dein fuligineis.

Hab. in culmis exsiccatis et putridis Arundinis Donacis in Horto bot. patavino Ital. bor.

40. **Didymosphaeria arenaria** Mout. Not. Ascom. p. 75. — Peritheciis 3015 sparsis, immersis, membranaceis, $\frac{1}{4}$ mm. diam., minute papillatis,

matrice circa ostiolum fuscescente; ascis tereti-clavatis, apice late rotundatis, 8-sporis, 140-150 \approx 27-30; paraphysibus filiformibus septatis ascos superantibus; sporidiis 1-2-seriatis, obovato-oblongis, medio septatis leviterque constrictis, pallide fuliginosis, pellucidis, 28-30 \approx 12-14.

Hab. in culmis *Ammophilæ arenariæ* pr. Hamburg Germaniæ, sociâ *Leptosph. Ammophilæ* (in Rehm Ascom. n. 691).

41. **Didymosphaeria lusitanica** (Niessl) Berl. et Vogl. Add. Syll. p. 115, 3016

Phorcys lusitanica Niessl Contr. Fl. Myc. Lus. p. 15. — Peritheciis sparsis, omnino immersis, clypeatis, majusculis, 0,4-0,5 mm. diam., globosis, atris, ostiolo minuto; ascis elongato-cylindricis, pedicellatis, membrana interna apice valde incrassata, 8-sporis, 120-140 \approx 10-12; sporidiis monostichis, oblongis, utrinque rotundatis, plerumque curvatis vel quasi medio infractis, uniseptatis et valde constrictis, guttulis binis, atro-fuscis, 18-22 \approx 8-10; paraphysibus tenuibus, apice ramosis; ascis sporidiisque ut in *Delitschia*, sed peritheciis profunde innatis, tectis.

Hab. in culmis aridis *Donacis arundinaceæ*, Galla prope Figueira da Foz Lusitaniæ (MOLLER).

42. **Didymosphaeria rhytidosperma** Speg. Fungi Guar. Pug. I, n. 223. 3017

— Peritheciis sparsis v. laxissime subseriatis, epidermide areolatim nigrefacta tectis eique arcte adnatis, prominulis, hemisphærico-lenticularibus, majusculis, 1-1,2 mm. diam., ostiolo minuto, umbo-natulo, vix epidermidem perforante, non exerto donatis, crassiuscule membranaceo-carnosulis, sordide albescenti-ceraceis, circa ostiolum fuscescensibus sæpeque coriacellis, contextu prosenchymatico-fibrosulo, subradiante, parum distincto, albo-fuscescente; ascis cylindraceis, apice obtusissime rotundatis, postice breve attenuato-stipitatis, 120-130 \approx 7-8, jodii ope nulla, paraphysibus gracilis, obscure septulatis obvallatis, 8-sporis; sporidiis recte monostichis, utrinque acute rotundatis, rectis, medio constrictis; loculis æqualibus, non v. 1-2 crasse guttulatis, episporio pellucido, pulchre fuligineo, longitudinaliter dense rugoso-striatis, rugis simplicibus, rectis v. vix helicoideis.

Hab. ad culmos dejectos emortuos *Bambusaceæ* cujusdam in dumetis prope Carapeguá Bras. — Species pulcherrima ac distinctissima, characteribus peritheciis subnectriacei, facile novum genus inter *Nectriaceas* constituens.

IV. In Acotylodoneis.

43. **Didymosphaeria filicina** Winter Grevill. XV, p. 88. — Peritheciis 3018

gregariis, in foliolis plerumque dealbatis epiphyllis, immersis, globosis, poro simplici pertusis, vertice late papillæformi tantum prominulis atris, coriaceis, 80-90 μ . diam.; ascis fasciculatis, oblongis, utrinque attenuatis, substipitatis, 8-sporis, 44-48 \approx 12,5; paraphysibus parum distinctis, filiformibus, plerumque mucosis; sporidiis di-tristichis, bacilliformibus cylindrico-subclavatis, utrinque rotundatis, medio septatis, haud constrictis, hyalinis, 14-16 \approx 3.

Hab. ad folia subviva Gymnogrammes Colomelani in Brasilia (ULE). — Si sporidia constanter hyalina, potius *Didymellæ* species.

44. **Didymosphaeria pteridina** Sacc., *Didymospharella filicina* Cooke 3019

in Grevill. XIX, p. 3. — Sparsa, tecta; peritheciis globosis, atris, minutis, ostiolis papillatis; ascis cylindraceis; sporidiis uniseriatis, ovalibus, medio septatis, nec constrictis, fuscis, 8 \approx 5.

Hab. in caulis Pteridis, New Zealand (COLENSO).

45. **Didymosphaeria microstictica** (Leighton) Winter in Hedw. 1886, 3020

p. 25, *Verrucaria microstictica* Leight. Lichen. Flora of Great-Britain. Edit. II, p. 493. — Peritheciis minutissimis, plus minusve immersis, globosis, vertice plano-convexo, pertuso, fusco-atris; ascis cylindricis, breve stipitatis, 8-sporis, 78-88 \approx 10; sporidiis mono-vel subdistichis, elliptico-oblongis, bicellularibus, medio constrictis, atris, opacis, 14 \approx 7; paraphysibus numerosis, filiformibus, subramosis.

Hab. in thallo Acarosporæ fuscatae et A. cervinæ in Germania et Britannia.

46. **Didymosphaeria (?) Sauteri** (Körber) Winter in Hedw. 1886, p. 25, 3021

Polycoccum Sauteri Körber Parerga pag. 470. — Peritheciis minutissimis, conferte stipatis, maculas subatras formantibus, demum sæpe confluentibus, semiimmersis, globosis, ostiolo umbonato fusco-atris; ascis cylindraceis, substipitatis, 8-sporis, 100 \approx 15-16; sporidiis monostichis, dacryoideis, cellula superiori multo latiori longiorique, fuscis, 16-19 \approx 9-10,5.

Hab. in Stereocaulo condensato et S. alpino atque in Lecidea fumosa in Germania.

47. **Didymosphaeria Bryonthæ** (Arnold) Winter in Hedw. 1886, p. 25, 3022

Endococcus Bryonthæ Arnold in Lichenol. Fragment. XVI, p. 26, tab. II, fig. 15, in Flora 1874. — Peritheciis gregariis, minutis,

subglobosis, valde convexis, prominulis, vertice pertusis; ascis fere cylindraceis (sec. icon. clavatis, deorsum stipitiformi-attenuatis), 8-sporis; paraphysibus distinctis, tenuibus capillaribus; sporidiis inordinatis, oblongis vel clavatis, bicellularibus, haud vel parum constrictis, pallide fuscidulis, $12\text{-}16 \approx 4\text{-}5$.

Hab. in disco apotheciorum Lecanoræ subfuscæ var. bryonthæ in Germania.

48. **Didymosphaeria Sporastatiæ** (Anzi) Winter in Hedw. 1886, p. 25, 3023

Polycoccum Sporastatiæ Arnold in Flora 1874, p. 144, tab. II, fig. 9. *Tichothecium Sporastatiæ* Anzi Neosymbolæ pag. 17. — Peritheciis sparsis, fere omnino immersis, globosis, vertice tantum parum convexo pertusoque prominulis, atris; ascis oblongis, 17μ . crassis; sporidiis distichis, oblongo-clavatis, inæqualiter didymis, medio parum constrictis, fuscis, $17\text{-}23 \approx 7\text{-}8$.

Hab. in thallo Sporastatiæ cinereæ in Germania.

MASSARIELLA Speg. — Syll. I, p. 716.

1. **Massariella vibratilis** (Fuck.) Sacc. — Syll. I, pag. 716. — var. 3024

Mespili Pass. in Brun. Nouv. Fragm. p. 29. — Peritheciis carbonaceis, cortice tectis, erumpentibus, ostiolo papillato; ascis cylindraceis, paraphysibus brevibus cinctis; sporidiis ovoideis, medio obscure septatis, strato tenui, hyalino involutis, $25 \approx 12$.

Hab. in ramis emortuis Mespili germanicæ, Pessines pr. Saintes Galliæ.

2. **Massariella scoriadea** (Fr.) Cooke Grev. XVII, p. 93. *Anthostoma scoriadeum* (Fr.) Sacc. Syll. 1127. — Sporidia elliptica, uniseptata, $70 \approx 23$, loculo superiore majusculo, medio constricta, episporio crasso, hyalino. Ex Cookei autopsia Scler. Suec. n. 344.

3. **Massariella seriata** Cooke Grevill. XVII, pag. 92. — Peritheciis 3025 depressiusculis, majusculis, seriato-dispositis, peridermio tectis, demum fissis; ascis clavatis; sporidiis ellipticis, $60 \approx 18\text{-}20$, 1-septatis, medio constrictis, fuscis, loculis æqualibus, episporio crasso, hyalino obvolutis.

Hab. in ramis Caryæ, S. Carolina (RAVENEL).

4. **Massariella didymopsis** Mouton in Bull. Soc. R. Bot. Belg. 1886, 3026 p. 151. — Peritheciis sparsis, corticolis, globoso-depressis, $\frac{1}{3}$ mm. diam., peridermio nigrificato connexis, ostiolo brevi erumpente; ascis cylindraceis, breve stipitatis; paraphysibus ramulosis obvallatis, jodii ope non cœrulescentibus, octosporis, $145\text{-}160 \approx 17\text{-}20$; sporidiis monostichis, ellipsoideis, uniseptatis, constrictis, loculis

juxta septum guttulatis, olivaceo-fuscis, strato mucoso obvolutis,
20-25 \approx 11-12.

Hab. in ramulis Sarothamni scoparii, Gomzé Belgii.

5. **Massariella bispora** Curtis Catalogue et Herb., Cooke in Grevill. 3027
XVII p. 93. — Peritheciis corticolis, subgloboso-depressis, tectis,
subsparsis, ostiolo peridermum perforante; sporidiis matricem inqui-
nantibus; ascis clavatis; sporidiis ellipticis, 1-septatis, fuscis, 45
 \approx 18-20, loculis æqualibus, medio constrictis, muco hyalino ob-
volutis.

Hab. in cortice Aceris, in America boreali (Dr. CURTIS).

6. **Massariella Rhododendri** Oud. Contr. Mycol. XIII, p. 29. — Peri- 3028
theciis parvis $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., sparsis, in parte peripherica pa-
renchymatis corticalis nidulantibus, periderma perforantibus, lentis,
subcoriaceis, primitus fere globosis, postremo depresso-depressis, centro per-
foratis; ascis cylindraceis, 170 \approx 15; sporidiis monostichis, badiis,
fere opacis, ovoides v. oblongis, 19-23 \approx 14, 1-septatis, medio levis-
sime constrictis, eguttulatis.

Hab. in ramulis Rhododendri arborei pr. La Haye Hollan-
diæ (Dom.^a DESTRÈE). — Stratum hyalinum in adultis vix con-
spicuum.

GIBELLINA Pass. in Rev. Myc. 1886, pag. 177. *Gibellia* Pass. Una
nuova mal. Frum., nec Sacc. (Etym. a cl. Prof. J. GIBELLI in Uni-
vers. Taurinensi Botanices Prof. eximio). — Stroma vel subiculum
matrici immersum, byssoides, atro-griseum, primitus canescens,
plus minus expansum, ex hyphis tenuibus, fumoso-pellucidis intri-
catis formatum. Perithecia stromate insidentia vel immersa, con-
tigua, contextu fibroso, globosa, in collum subæquilonium, cras-
siusculum, rectum vel subinde flexuosum, erumpens attenuata.
Asci elongato-clavati, paraphysati, octospori. Sporidia oblonga, di-
dyma, fuscescentia. — A genere *Neopeckia*, cui characteribus ac-
cedit, peritheciis innatis et toto habitu differt; ab aliis generibus
phæodidymis vero stromate byssideo et peritheciis fibrosis præ-
cipue diversum.

1. **Gibellina Cerealis** Pass. Nuova mal. Frum. pag. 5. et Rev. Myc. 3029
1886, p. 177. — Stromate intra foliorum vaginam expanso; peri-
theciis in series lineares digestis vel sparsis, rarius cæspitosis, sub-
globosis, ostiolo vaginam perforante, vix emerso, acutiusculo; ascis
paraphysibus tenuibus obvallatis, diffluentibus, 100-110 \approx 22-25;
sporidiis distichis, fusiformi-lanceolatis, primo hyalinis, integris

dein medio septatis, non constrictis, e melleo vel avellanaceo fuscouscentibus, 22-30 \approx 7,5-9.

Hab. ad culmos languidos Tritici vulgaris, quos corruptit, Vigatto et Toschiura prope Parma Ital. bor. — Fungillus satus ab ill. Passerini in olla, una cum Tritico, primo anno quievit, altero vero plantas novas Tritici affecit et vexavit. Nonne, observante Passerinio, sporidia tantum germinant post longam humationem?

PROTOVENTURIA Berl. et Sacc. in Atti Soc. Veneto-Trentina 1886

(Etym. a *protos* primus et *Venturia*), *Venturia* De Not. Atti VI, Riun. Scien. Ital. pag. 484. — Perithecia superficialia, majuscula, carbonacea, fragilia, globoso-depressa, vertice setis rigidis, aterrimis, longiusculis vestita, basi setulis subtilioribus, numerosissimis, pallidis, subtortuosis, septatis cincta, poro rotundo, amplo pertusa. Asci oblongi vel elliptici, basi abrupte attenuati, in pendulum brevissimum desinentes, octospori. Paraphyses nullæ vel (verisimiliter) obsoletæ. Sporidia constricto-didyma, bilocularia, loculis subæqualibus, saturate fuligineis.

1. **Protoventuria Rosæ** (De Not.) Bér. Sacc. l. c. *Venturia Rosæ* De 3030 Notaris Atti VI Riun. Sc. It. p. 484, t. I, f, I, et Giorn. Bot. It. 1864, p. 332, Ces. et De Not. Schema Sfer. It. pag. 51, *Pyrenophora Rosæ* (De Not.) Sacc. Syll. II, p. 285. — Peritheciis sparsis v. laxè gregariis, globosis vel globoso-depressis, majusculis, superficialibus, poro ampio pertusis, sursum setis rigidis, aterrimis apice acutis vestitis basique filamentis mycelicis numerosissimis, 4 μ . diam., longis, sinuosis, simplicibus, sæpe 4-5 coalitis, septatis, pallidioribus cinctis; ascis amplis, saccatis, sursum obtusis, basi abrupte attenuatis atque in stipitem brevissimum desinentibus, tunica crassa præditis, 95-105 \approx 24-27, octosporis; sporidiis inordinatis vel distichis, majusculis, 28-32 \approx 13-15, uniseptatis, ad septum constrictis, loculis subæqualibus vel superiore quidquam majore obtusore, saturate olivaceo-fuligineis,

Hab. in ramulis emortuis Rosæ alpinæ in Monte Cenisio Italiæ borealis.

AMPHISPÆRIA Ces. et De Not. — Sacc. Syll. I, p. 718.

1. **Amphisphæria appianata** (Fr.) Ces. et De Not. Sacc. Syll. I, p. 718, Winter in Hedwigia 1886, p. 102. — Perithecia demum collapsa, concentrica rugulosa; asci e basi ampliata sursum oblongati, breve pedicellati, 8-spori, 85-105 \approx 19-21; sporidia subdisticha, loculo

superiori majori, ad medium valde constricta, utrinque rotundata; paraphyses filiformes.

2. **Amphisphæria decorticata** (Cooke et Hark.) Berl. et Vogl. Add. 3031
Syll. p. 124, *Sphæria* Cooke et Hark. in Grevillea XIII, p. 19. —
Primitus seriata, intra fibrillas corticis nidulans, demum subsuperficialis; peritheciis erumpentibus subserialibus sparsisve, subglobosis, atris, opacis, vix $\frac{1}{2}$ mm. diam.; ascis cylindraceis, octosporis; sporidiis ellipticis, uniseptatis, medio arcte constrictis, ex cellulis subglobosis efformatis, læte fuscis, 18 \approx 9.

Hab. in ramis decorticatis *Quercus* in California.

3. **Amphisphæria quercreti** Cooke et Mass. Grevill. XVI, p. 92. — Pe. 3032
ritheciis gregariis, majusculis, 1 mm. diam., initio immersis dein semi emergentibus, conico-convexis, diu epidermide tectis, papillatis, atris, basi applanatis; ascis cylindraceis 8-sporis; sporidiis uniseriatis, ellipticis, utrinque rotundatis, medio leniter constrictis, 1-septatis, brunneis, 28 \approx 10.

Hab. in cortice *Quercus albæ*, N. Carolina (DR. CURTIS).

4. **Amphisphæria Magnusii** Sacc. Bomm. et Rouss. Fl. Myc. Belg. II, 3033
p. 19. — Peritheciis superficialibus laxe gregariis globuloso-depressionis, nigris, carbonaceis, $\frac{1}{3}$ mm. diam., ostiolo brevissimo obtuso; ascis clavatis, 8-sporis, 100 \approx 16-18, apice obtusis, in stipitem 30-40 μ . long. attenuatis, paraphysatis; sporidiis subdistichis, oblongo-fusoideis, rectis v. curvis, 1-septatis, constrictis, biconicis, 22-28 \approx 8-10, olivaceo-fuligineis.

Hab. in ligno *Quercus* et *Carpini* in Groenendaal Belgii.
Ab *A. Spegazziniana*, cui affinis, differt peritheciis sporidiisque majoribus.

5. **Amphisphæria suecica** (Rehm) Sacc., *Melanopsamma suecica* Rehm 3034
Beitr. Ascom. Hedw. 1882, p. 120. — Peritheciis solitariis aut gregariis, hemisphæricis, dimidiatis, vix papillatis, poro pertusis, 0,5 mm. diam., subscabris, atris, sessilibus in ligni cariosi superficie; ascis piriformibus, crassis, 8-sporis, 90 \approx 21; sporidiis (ut in *Delitschia*) 2-cellularibus, cellula superiore fere duplo angustiore, utraque nucleo 1 magno prædita, fuscis, muco gelatinoso lato obductis, 2-stichis, 25 \approx 8; paraphysibus ramosis.

Hab. ad ramum fagineum putridum prope Krumbad Sueviæ bavaricæ (BRITZELMAYR).

6. **Amphisphæria anceps** Sacc. et Briard Miscell. Myc. I, n. 2207. 3035
— Peritheciis ligno striolatim nigrificato subimmersis, globulosis, compressiusculis, minutis, $\frac{1}{4}$ mm. diam., minute papillatis; ascis

tereti-clavulatis, stipitatis, apice obtusis, octosporis, $63 \approx 10$, p. sporif.; sporidiis oblique monostichis, 1-septatis, lenissime constrictis, $9-11 \approx 5-6$, biguttatis, olivaceis.

Hab. in ramis decorticatis, emortuis Populi, prope Troyes Galliæ (BRIARD). — Hinc *Didymosphæriæ* (Microtheliæ), hinc *Val-sariæ* (Phæospermati) nutat, ideoque anceps *Amphisphæriæ* species.

7. **Amphisphæria subiculosa** Ell. et Ev. in Journ. of. Myc. 1886, 3036 p. 103. — Peritheciis superficialibus, gregariis, depresso globosis, $\frac{1}{3}$ mm. diam., brunneo-atris, rugosis, sursum sublucidis, ostiolo radiato-sulcato, non prominulo præditis, mycelio byssino, brunneo, repente, late effuso insidentibus; ascis cylindraceo-clavatis, $90-100 \approx 12-15$, copiose paraphysatis; sporidiis distichis, oblongis, pallide brunneis, 1-septatis ad septum leniter constrictis, utrinque obtusis, binucleatis, $15-10 \approx 4-5$.

Hab. in Populo decorticata, in America boreali (LANGLOIS).

8. **Amphisphæria salicina** (Rehm) Sacc., *Melanopsamma salicina* 3037 Rehm Beitr. Ascom. Hedw. 1882, p. 120. — Peritheciis carbonaceis, hemisphæricis, vix distincte papillatis poroque pertusis, magnitudine seminis sinapeos, solitariis, hic illic 2-8 confluentibus, in ligno longe lateque denigrato; sporidiis biscoctiformibus, acutatis, medio distincte constrictis, 2-cellularibus, quaque cellula plerumque nucleo 1-rarius pluribus prædita, hyalinis, dein fuscis, $20 \approx 6$, distichis; paraphysibus ramosis.

Hab. ad truncum Salicis retusæ, Waldraſt in alpibus tirolensibus (Doct. ARNOLD).

9. **Amphisphæria heteromera** Briard et Sacc. in Rev. Myc. 1885, 3038 n. 12. — Peritheciis subsparsis e basi insculpta superficialibus, globosis, papillatis, carbonaceis, atris nitidulis; ascis cylindraceis, subsessilibus, $104-108 \approx 18-20$, octosporis, filiformi-paraphysatis; sporidiis subdistichis, oblongo-fusoideis, rectiusculis, utrinque subattenuatis, infra medium septatis constrictisque, fuligineis, loculo infero minore.

Hab. in trunco decorticato Carpini prope Troyes Galliæ (BRIARD).

10. **Amphisphæria Wellingtoniæ** (Cooke et Hark.) Berl. et Vogl. Add. 3039 Syll. p. 124, *Sphæria Wellingtoniæ* Cooke et Hark. in Grevillea XIII, pag. 19. — Gregaria, immersa; peritheciis atris, elongato-compressis, hysterioideis, striatis, opacis, $150-180 \approx 80$, poro pertusis; ascis cylindraceis, 8-sporis; sporidiis ellipticis, uniseptatis, fuscis, vix constrictis, loculo utroque uniguttulato, $12-14 \approx 8$.

Hab. in ligno dealbato Sequojae in California (HARKNESS).

11. **Amphisphæria deformis** E. et Langl. Journ. Myc. 1888 p. 123. 3040

— Peritheciis gregariis, erumpentibus, subglobosis v. subcompressis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., nigris, asperulis, ostiolo irregulari, lacerato-depresso; ascis clavato-cylindraceis, circ. 50 \approx 7, subevanescentibus, obscure paraphysatis; sporidiis monostichis, oblongis v. clavato-oblongis, brunneis, 1-septatis, constrictis; loculis guttulatis.

Hab. in palis vetustis pineis (Juniperi v. Thujæ), Pointe à la Hâche Louisiana Amer. bor. — Habitus fere *Melanopsammæ* compressinæ.

12. **Amphisphæria aquatica** (Ell. et Ev.) Berl. et Vogl. Add. Syll. 3041

p. 125 *Sphæria aquatica* Ell. et Ev. in Torr. bot. Club. 1884, p. 42. — Peritheciis sparsis, membranaceis, 250 μ . diam., semiimmersis, subglobosis vel parum elongatis; ostiolo papilliformi; ascis linearibus, 150 \approx 6-8, copiose paraphysatis; sporidiis recte monostichis, oblongis, uniseptatis, septo leniter constrictis rectis vel leniter curvulis, primum pallidis, crasse binucleatis, demum obscure brunneis, 15-20 \approx 6.

Hab. in fundo situlæ juniperinæ diu aqua irrorato, Newfield Amer. bor.

13. **Amphisphæria perpusilla** Pass. et Beltr. F. Sic. n. 13. — Peri. 3042

theциis peregrinis, superficialibus, carbonaceis, subglobosis; ascis subcylindricis, basi breviter stipitatis, paraphysibus filiformibus sublongioribus obvallatis, 75-90 μ . long., 8-sporis; sporidiis biseriatis vel oblique uniseriatis, elliptico-oblongis, medio septatis fuscidulis, 15 \approx 5.

Hab. in ligno denudato arido Oleæ europææ, Monostalla Siciliæ.

14. **Amphisphæria strichnicola** Pat. Journ. Bot. 1889 p. 168. — Pe. 3043

ritheciis sparsis, nigris punctiformibus, dimidiatis, ostiolatis, ascis cylindraceis, 8-sporis, 110 \approx 16; paraphysibus numerosis, filiformibus, simplicibns v. ramosis; sporidiis monostichis, septo constrictis, brunneo-viridulis, 25-28 \approx 6-7.

Hab. in cortice Strichni Gubleri, Haut-Orénoque Venezuela (A. GAILLARD).

15. **Amphisphæria immersa** Berl. et Vogl. F. Ancon. p. 8, t. I, f. 6. — 3044

Peritheciis ligno profunde immersis, ostiolo papillato, crassiusculo prominulis, globosis, atris, 300 μ . diam., carbonaceis; ascis cylindraceis, basi in stipitem brevem, nodulosum abeuntibus, 8-sporis, 120-130 \approx 12; paraphysibus filiformibus, asco longioribus; sporidiis

oblique monostichis, ovoideis v. ovoideo-oblongis, 1-septatis, ad septum non v. vix constrictis, loculis crasse 1-guttatis, 18-22 \approx 8-9, rufo-fuligineis.

Hab. in ligno sicco Tamaricis cujusdam pr. Ancona Ital.

16. **Amphisphæria umbrinoides** Passer. Diagn. F. N., I, n. 43. — Peritheciis subsparsis, superficialibus v. basi vix insculptis, carbonaceis, subglobosis, ostiolo papillato; ascis paraphysibus tenuibus stipatis; sporidiis oblongo-fusiformibus pr. medium septatis et constrictis, loculis ut plurimum valde inæqualibus, initio hyalinis, biguttulatis, tandem fuligineis v. castaneo-fuscis, 23-28 \approx 7,5-9.

Hab. in cortice vetusto Æsuli Hippocastani, Parma Ital. — Ad *Amphisph. umbrinam* fructificatione accedit, sed perithecia multo minora et sporidia plus minusve, et interdum vehementer, constricta.

17. **Amphisphæria closterophora** B. et Br. Grevill. XVI, pag. 92. — 3046 Peritheciis sparsis, majusculis, 1 $\frac{1}{2}$ mm. diam., semiimmersis, conicis, v. basi applanatis, atris, nitidis, ostiolo papillato; ascis clavatis 8-sporis; sporidiis lanceolatis, 1-septatis, medio constrictis, fuscis 75 \approx 10.

Hab. in ramis denudatis in insula Java (KURZ).

18. **Amphisphæria biocellata** Pass. in Martelli Flor. Bog. pag. 143. 3047 — Peritheciis ligno insculptis, ostiolo obtuso, emerso, oblongis, umbrinis; ascis cylindraceis, basi angustatis, sessilibus, paraphysibus filiformibus etiam duplo superatis, octosporis, 45-60 \approx 7,5; sporidiis oblique monostichis, ellipsoideis, ad septum non vel vix constrictis, loculis guttulatis, 10-12,5 \approx 5, castaneo-fuscis.

Hab. in lignis nudis lichenosis, pr. Keren Abyssiniæ (BECCARI).

19. **Amphisphæria lamprostoma** Pass. in Martelli Flor. Bog. p. 143. 3048 — Peritheciis sparsis, superficialibus vel vix insculptis, subglobosis, atro-opacis, ostiolo papillato, nitido; ascis cylindraceo-subclavatis, paraphysatis, octosporis, 70 \approx 7,5; sporidiis monostichis, ellipsoideo-oblongis, medio septatis, non constrictis, loculis subæqualibus, flavo-fuscis, 10-12 \approx 5.

Hab. in corticibus arborum annosarum, Keren Abyssiniæ (BECCARI).

20. **Amphisphæria Eduardi** Pass. in Mart. Fl. Bog. pag. 144. — 3049 Peritheciis peridermio incidentibus, sparsis vel aggregatis, hemisphericis, papillato-ostiolatis, carbonaceis; ascis oblongis, paraphysatis, attenuato-stipitatis, octosporis, 100 \approx 15; sporidiis distichis fusiformibus, utrinque acutis, medio septatis et constrictis; loculis subinæqualibus sæpe guttulatis, luteo-fuscidulis, 20-22,5 \approx 7,5.

Hab. in corticibus lichenosis, Keren Abyssiniæ (BECCARI).

21. **Amphisphæria oronoensis** (E. et E.) Sacc. *Sphæria* E. et E. Journ. 3050

Myc. 1887, p. 117. — Peritheciis sparsis, subcarbonaceis, nigris, rugosulis, subsuperficialibus, basi tantum ligno subimmersis, minutis, circ. $\frac{1}{6}$ mm. long., globosis v. leniter depresso-globosis, ostiolo papilliformi; ascis linearibus, 75 ± 5 , p. sp. circ. 50μ . long., copiose paraphysatis; sporidiis monostichis, oblongo-ellipsoideis, brunneis, 1-septatis, non constrictis, $6-8 \pm 2,5-3$; loculis æqualibus v. inferiore paullo angustiore.

Hab. ligno putri, Orono Maine Amer. bor. (F. L. HARVEY). — *A. sardoæ*, ut videtur, affinis.

22. **Amphisphæria corticola** Pat. Bull. Soc. Myc. 1888, p. 113. — 3051

Peritheciis sparsis, rotundatis, 1,5-2 mm. diam., levibus, nigris, superficialibus, basi cortice incrusted, intus griseolis, pariete crasso præditis; ascis?; sporidiis brunneis, ovoideis, obtusis, rectis 1-septatis, non constrictis, $23-25 \pm 8-10$.

Hab. in cortice arborum in Atures Venezuelæ.

23. **Amphisphæria Clusiæ** Pat. Bull. Soc. Myc. 1888, p. 113. — Peritheciis 3052

nigris, opacis, superficialibus, 10-15 gregariis, subglobosis, ostiolo mammillato; ascis cylindraceis, 100 ± 10 ; paraphysibus copiosissimis, hyalinis, linearibus; sporidiis octonis, monostichis, brunneo-nigricantibus, ovoido-rotundatis, 1-septatis, constrictis, 2-guttatis, 12 ± 6 .

Hab. in foliis dejectis putridis Clusiæ roseæ in Maipures Venezuela.

24. **Amphisphæria dothideospora** Cooke et Harkn. in Grevillea XIV, 3053

p. 9. — Gregaria; peritheciis emergentibus, epidermide nigrefacta tectis, conico-convexis, nitidis, ostiolo papillato; ascis subclavatis, octosporis; sporidiis subellipticis, inæqualiter uniseptatis, constrictis, fuligineis, 20 ± 12 .

Hab. in caule Mimuli in California (HARKNESS).

25. **Amphisphæria Hypoxylon** E. et E. in Journ. of Myc. 1886, p. 41. 3054

— Peritheciis dense gregariis, superficialibus, globosis, $\frac{1}{4}$ mm. diam., rugosis, atris, pruinosis, membranaceo-carbonaceis; ostiolo vix prominulo; ascis clavato-cylindraceis, $55-65 \pm 10-12$, copiose paraphysatis; sporidiis oblique monostichis, ovato-ellipticis, brunneis, uniseptatis, $8-9 \pm 5$.

Hab. parasitice in Hypoxylī effusi stromatibus, Plaquemines Amer. bor. (REV. A. B. LANGLOIS).

26. **Amphisphæria terricola** Wint. Journ. Bot. 1887, p. 270. — Peri- 3055

theciis gregariis seu subsparsis, primo immersis, demum plus minusve emersis, interdum fere superficialibus, globosis, magnis, vertice rotundato, ostiolo papilliformi, pertuso coronatis, interdum demum umbilicatis et collapsis, atris, tenuissime pilosis, 3-4 mm. lat.; ascis cylindraceis, in stipitem longum attenuatis, apice rotundatis, 8-sporis, 175-210 \pm 26 p. sp.; sporidiis oblique monostichis v. subdistichis, oblongis, utrinque subattenuatis, inæqualiter didymis, sæpe curvatis, loculo supero intense fusco, parum latiore, sæpe cuspidato semperque apiculo brevissimo, hyalino, papilliformi prædicto, loculo infero angustiore, rotundato, dilutius colorato, sine apiculo, 38-54 \pm 14-18.

Hab. ad terram madidam in Villers-Cotterets (QUESTIER) et Falaise Galliæ (DE BRÉBISSON).

27. **Amphisphæria albo-maculans** (Schwz.) Cooke Grevill., *Sphæria* Schw. — Syll. II, pag. 431. — Sporidia fusiflora uniseptata, brunnea, 12 \pm 3.
28. **Amphisphæria conferta** (Schwein.) Cooke Grev. XV, p. 81 *Sphæria* Schw. — Syll. II, 4277, Herb. Berk. 9603. — Sporidia uniseptata, fusca, utrinque obtusa, medio constricta, 12 \pm 4.
29. **Amphisphæria diffusa** (Schwein.) Cooke Grev. XV, p. 81. Sacc. Syll. N. 4283, Herb. Berk. 9602. — Sporidia uniseptata, fusca, utrinque acuminata, medio constricta, 18-20 \pm 6-7.

Ad genus **Amphisphæriam** forte pertinet *Hypoxyylon? gregale* (Syll. I, n. 1469) quod idem est ac *Valsaria gregalis* (Syl!. I, n. 2487), nam perithecia, teste Cooke, non confluentia.

DELITSCHIA Auersw. — Sacc. Syll. I, p. 732.

1. **Delitschia furfuracea** Niessl in litt. Rehm fasc. XV, n. 747. — 3056
Peritheciis parenchymaticis, brunneis; ascis amplis, clavatis, 8-sporis, 300 \pm 50; sporidiis distichis, oblongis, obtusis, medio constrictis, bicellularibus, strato mucoso obvolutis, obscure brunneis, 45-50 \pm 21; paraphysibus filiformibus vel ramosis, guttulatis.

Hab. in fimo leporino, Brünn Moraviæ.

2. **Delitschia Marchalii** Berl. et Vogl. Add. Syll. p. 127, *Delitschia* sp. 3057
(?) March. Fung. coproph. p. 10. — Peritheciis gregariis, minutissimis, subglobosis; ascis cylindraceis, acutiusculis, deorsum brevissime attenuato-stipitatis, 75-85 \pm 11-13, parum et minute paraphysatis, 8-sporis; sporidiis plerumque oblique monostichis, ellipticis, apicibus obtusis, ad septum vix constrictis, primo hyalinis tandem

subfuscis, annulo hyalino, angusto cinctis, in articulos binos haud secedentibus, $10,5-12 \approx 5,5-6$.

Hab. in fimo Cuniculorum inter Limelette et la Baraque in Belgio. — *Delitschiae elephantinæ* affinis sed magnitudine ascorum et forma sporidiorum distinguenda.

3. **Delitschia Niesslii** Oud. Hedw. 1882, p. 163. — Peritheciis sparsis, ad herbarum residua in fimo cuniculorum vetusto superficialibus, minutis, subglobosis, fuscis, glabris, levibus; ascis cylindraceis 70-80 ≈ 8 , paraphysibus subtilissimis ramosissimis obvallatis, 8-sporis; sporidiis nitidissime monostichis, ellipticis, $14 \approx 6$, utrinque obtusis, ad septum vix et ne vix quidem constrictis, maturis non aut saltem difficile in articulos dilabentibus.

Hab. in fimo Cuniculorum prope Harlem Hollandiæ.

4. **Delitschia vaccina** Passer. Fungi Nov. IV, p. 4. lib. — Peritheciis gregariis, minutis semiimmersis, glabris, atris, contextu membranaceo, minute celluloso fuligineo; ascis longis, tubulosis, basi in stipitem longissimum, filiformem attenuatis, apice acutis, 8-sporis; sporidiis oblique uniseriatis, ellipticis 1-septatis, ad septum non constrictis, primo pallide olivaceis, dein intense fuscis, $50 \approx 22-25$.

Hab. in fimo vaccino, sociâ *Hypocopra minima* et *Sporormia intermedia* in pratis Cisa in apennino Parmensi Ital.

5. **Delitschia canina** Mout. Ascom. Lièg. II, pag. 7, t. I, f. 4. — Peritheciis gregariis, immersis, globosis, $500-700 \mu$. diam., glabris, sordide atris, contextu coriaceo opaco, ostiolo nigro, crasso, rudi; ascis clavatis, stipitatis, 8-sporis, $200-200 \approx 26$, p. sp. 170μ . long.; paraphysibus filiformibus; sporidiis subdistichis, oblongis, obtusis, ad septum arctissime constrictis ideoque facillime in articulos binos secedentibus, fusco-opacis, strato mucoso obvolutis, $45-55 \approx 15-16$.

Hab. in stercore canino vetusto, Gomzé Belgii.

6. **Delitschia patagonica** Speg. F. Pat. n. 130. — Peritheciis sparsis, immersis, globosis, $500-700 \mu$. diam., glabris, levibus, ostiolo papillulato-obtuso, atro, subcarbonaceo, superficiem matricis attingente v. vix exerto donatis, atris, crassiuscula membranaceis, contextu indistincto, fuligineo, opaco; ascis cylindraceis sursum obtuse rotundatis, deorsum abrupte, breve noduloseque stipitatis, $200 \approx 20$, paraphysibus crassiusculis, guttulatis, simplicibus parcis obvallatis, 8-sporis; sporidiis oblique monostichis, ellipticis, $35 \approx 16$, utrinque obtuse rotundatis, medio 1-septatis et leniter constrictis, non v. vix subinæquilateralibus, opace fuligineis, tunica mucosa, hyalina, tenui obvolutis.

Hab. in fimo Aucheniae Guanaco pr. Mount Entrance ad ostia Rio S. Cruz Patagoniae.

7. **Delitschia leptospora** Oud. Hedw. 1882, p. 163. — Peritheciis sparsis, ad herbarum residua in fimo cuniculorum vetusto superficialibus, minutis, subgloboso-conicis, atro-fuscis, glabris, levibus; ascis subsessilibus, brevibus, 60-70 \approx 11-12, cylindraceo-oblongis, paraphysibus subtilissimis, ramosissimis obvallatis, 8-sporis; sporidiis tristichis, fusiformibus, utrinque obtusiusculis, 22-23 \approx 4-5, fuscis, ad septum valde constrictis ideoque facillime in articulos dilabentibus.

Hab. in fimo Cuniculorum prope Harlem Hollandiae.

8. **Delitschia consociata** Mout. Ascom. Lièg. II. p. 7, t. I, f. 8. — 3063 Peritheciis immersis, minutis, glabris; ascis oblongis, stipite breve et crasso, 8-sporis, 80 \approx 20; paraphysibus filiformibus, ramosis; sporidiis distichis, oblongis, utrinque obtuse rotundatis, 18-20 \approx 6, -7,5. haud constrictis, strato mucoso obductis.

Hab in fimo Capreolorum, sociâ *Del. Marchalii* et *D. moravica* in Belgio. — *D. leptosporæ* affinis sed sporidiis crassioribus non constrictis dignoscitur.

9. **Delitschia lignicola** Mouton in Bull. Soc. R. Bot. Belg. 1886, 3064 p. 151. — Peritheciis plerumque gregariis, semi vel omnino immersis, atris, globosis, $\frac{2}{3}$ mm. diam., ostiolo papillato, hirsuto; pilis fuscis, translucidis, 4 μ . crassis, septatis obtusis; ascis cylindraceis, stipitatis, apice rotundatis, paraphysibus filiformibus obvallatis, octosporis, 195-210 \approx 12-15, p. sp. 165-170 \approx 12-15; sporidiis monostichis, parallelogrammo-ellipsoideis, uniseptatis, leniter constrictis, opacis, muco non obductis.

Hab. in fragmento ligneo putrescente, Tilff Belgii.

NEOPECKIA Sacc. in Peck Torr. bot. Club. 1883, p. 127 (Etym. a *neos* novus et a cl. C. H. PECK). — Perithecia carbonacea, subfragilia, superficialia sed subiculo copioso, effuso semiimmersa, rarius subiculo parco, perithecia vero pilosa, globosa, papillata, denique latiuncule pertusa; asci octospori, elongati copiose paraphysati. Sporidia didyma, fuliginea. — A genere *Amphisphæria* subiculi copiosi vel pilorum præsentia differt, ab *Enchnosphæria* et *Eriosphæria* sporidiis perfecte didymis, fuligineis recedit.

1. **Neopeckia Coulteri** (Peck) Sacc. in Torr. bot. Club. 1883, p. 127, 3065 *Sphaeria Coulteri* Peck in Hayden's U. S. Geol. Survey, 1872, p. 792, *Enchnosphæria (?) Coulteri* Sacc. Syll. II, p. 207, *Lasio-*

sphæria acicola Cooke in Grevillea 1880, pag. 87. *Amphisphæria* (?) *acicola* Sacc. Syll. vol. I, p. 727. — Peritheciis 0,5 mm. diam., demum papillâ amissâ, perforatis; ascis 130-140 \approx 14-15, p. sp., paraphysibus copiosis, filiformibus; sporidiis utrinque obtusiusculis, 20-28 \approx 9-10, intense fuligineis.

Hab. in foliis Pini in montibus Rocky Amer. bor. — Habitu externo *Enchnosphæriæ Pinetorum* peraffinis, sed fructificatio prorsus aliena.

2. **Neopeckia quercina** G. Delacroix Bull. Soc. Myc. Fr. 1890 f. IV. — 3066

Peritheciis atris, carbonaceis, primum imperforatis, 300-400 μ . lat.; mycelio copioso, sub perithecio effuso, hyphis continua fuscis, 500 \approx 3, apice obtusiusculis; ascis cylindraceis, 180 \approx 12; sporidiis monostichis, primum subhyalinis, uniseptatis, biguttatis, demum intense fuscis, uniseptatis, guttulis non visibilibus, 24 \approx 10; paraphysibus furcatis, gracilibus.

Hab. in ligno quercino sicco, Saone et Loire Galliæ.

3. **Neopeckia fulcita** (Buckn.) Sacc., *Lasiosphæria fulcita* Bucknall 3067

Fungi Bristol X, p. I, f. XIII, f. 8. — Peritheciis confertis, emergenti-superficialibus, ovato-globosis, rugosis, pilis brevibus, septatis, brunneis, laxe conspersis, 0,4 mm. diam., ostiolo lato, profunde radiato-sulcato; ascis linearibus, 120 \approx 5, filiformi-paraphysatis; sporidiis ellipsoideis, mox 1-septatis, brunneis, 1-2-nucleatis, 12-13 \approx 5.

Hab. in ramis emortuis quercinis pr. Bristol Britanniæ.

RHYNCHOSTOMA Karst. — Sacc. Syll. I, p. 730.

1. **Rhynchostoma tinctum** (Ell. et Ev.) Berl. et Vogl. Add. Syll. p. 125, 3068

Ceratostoma Ell. et Ev. in Bull. Washb. Lab. Nat. Hist. 1884, p. 5. — Peritheciis erumpentibus, superficialibus, ovoideo-globosis, $\frac{1}{6}$ mm. diam., rugulosis, ostiolatis; ostiolis filiformibus, $\frac{2}{3}$ - $\frac{3}{4}$ mm. longis, sublevibus, subinde sursum parum dilatatis; apice globulo sporidiorum coronato; ascis subcylindraceis, circiter 30 \approx 7, subsessilibus, paraphysibus, filiformibus obvallatis; sporidiis distichis, oblongo-ellipsoideis, subacutis, uniseptatis, 6-7 \approx 2,5-3.

Hab. in ligno emortuo Aceris macrophylli, Klikitat Co. Am. bor. — Affine *R. minuto*.

2. **Rhynchostoma Beccarianum** Pass. in Martelli For. Bog. osensis, 3069

p. 144. — Peritheciis globosis, atris, basi ligno insculptis, in rostrum cylindricum, æquilongum, rectum, apice truncatum, albo-marginalatum et centro perforatum productis; ascis numerosis, cylin-

draceis, octosporis, 60×5 ; paraphysibus filiformibus, tenuibus, superantibus obvallatis; sporidiis oblique monostichis, interne fuscis, ellipsoideis, obscure septatis, $7,5 \times 4,5-5$.

Hab. in ligno denudato sicco, Keren Abyssiniæ (BECCARI).

3. **Rhynchostoma rubro-cinctum** Karst. Hedw. 1890 p. 178. — Peritheciis subgregariis, innato-erumpentibus conoideo-sphæroideis, basi applanatis, atris, glabris, 0,2 mm. diam., rostro cylindraceo, secto v. saepius subflexuoso, sublevi, brunnescente, sub apice obtuso rubro solitoque marginato, 1-2 mm. long.; ascis clavatis, cito diffluentibus, $40-45 \times 10$; sporidiis octonis, di-tristichis, ovoideis v. fusideo-oblängatis, 1-septatis, vix constrictis, olivaceis, $8-10 \times 3-5$; paraphysibus flexuosis, $1-1,5 \mu$. cr.

Hab. in ligno vetusto betulino, colore subrubro tincto, loco humido umbroso pr. Mustiala Fenniæ.

RHYNCHOMELIOLA Speg. Fungi Guar. Pug. I, p. 283 (Etym. *rhynchos* rostrum et *Meliola*, ob perithecia eximie rostrata). — Perithecia atra, parvula, globulosa, sursum in ostiolum cylindraceum, prælongum, obtusum producta, subiculo concolori, tomentosulo, conidiifero, phyllogeo insidentia; ascis cylindraceo-clavati, aparamphysati, 8-spori; sporidia elliptica, didyma, chlorino-olivacea. — Genus *Dimerosporio* affine, ostioli præsentia recedens, ad *Capnodium* pluribus characteribus vergens; ab utroque vero satis distinctum est, et melius sub *Sphæriaceis* militare videtur.

1. **Rhynchomeliola pulchella** Speg. Fungi Guar. Pug. I, n. 224. — Mycelio hypophyllo, cum tomento foliorum intertexto, fusco v. olivascenti-atro, maculas plus minusve elatas, $1-15$ mm. diam., indeterminatas, saepe confluenti-evanescentes, irregulares efficiente, ex hyphis sterilibus, repentibus, dense intricatis, fuligineis, parce septatis, subcrassiusculis, $4-6 \mu$. crass., hinc inde hyphis fertilibus, conidiiferis, erectis ornatis composito; peritheciis in mycelio nidulantibus et absconditis, globulosis, parvulis, $100-110 \mu$. d., membranaceis, contextu minute parenchymatico, fuligineo-atro, glabris, levibus, superne in ostiolum prælongum, subcylindraceum, $400-520 \times 25-30$, merenchymaticum, apice truncato-rotundatum, in senectute fimbriato-dehiscens, pallidius productis; ascis cylindraceo-clavatis, apice rotundatis, basi brevissime attenuato-stipitatis, $24-28 \times 5$, subdiffluentibus, aparamphysatis, 8-sporis; sporidiis distichis, ellipticis, rectis v. vix inæquilateralibus, utrinque acutato-obtusatis, medio

1-septatis non constrictis, 6-7 \approx 2 primo hyalinis, dein dilute oliveis.

Hab. ad folia viva Feijoæ Sellovianæ in silva subvirginæ Caà-guazù Bras. — Hyphæ conidiiferæ erectæ, rectæ, longiusculæ, 240-458 \approx 6-6,5, parce septatæ, fuligineæ, apice pallidiores acrosporæ, monospermæ, simplices; conidia fusoidea v. elliptico-v. clavulato-elongata, recta v. leniter inæquilateralia, antice acutiuscule rotundata, postice attenuata ac truncatula, 24-55 \approx 7 et primo continua, hyalina, dein 5-10-septata, ad septa leniter constricta, eguttulata, levia. Status conidicus tandem *Helminthosporium pulchellum* sistens.

OTTHIA Nits. — Sacc. Syll. I, p. 735.

1. **Otthia Spirææ** Fuck. — Sacc. Syll. I, p. 735, Winter in Hedwigia 1886, p. 104. — Ascii 175-193 \approx 17-18; sporidia medio parum constricta, opaca, 28-30 \approx 12; paraphyses filiformes.
2. **Otthia Monodiana** Sacc. et Roum. in Rev. Myc. 1883, p. 235, t. 39, 3071 fig. 6. — Peritheciis subcutaneo-erumpentibus, in cæspites minutos collectis, globosis, obtusis, atris, rugulosis, $\frac{1}{2}$ mm. diam.; ascis cylindraceis, 160-170 \approx 18, paraphysatis, breve stipitatis, octosporis; sporidiis oblique monostichis, ellipsoideis, constricto-1-septatis, 23-25 \approx 12-14, fuligineis.

Hab. in ramis Salicis capreæ in Arduennis (LIBERT). — Affinis *O. Alni*.

3. **Otthia Pteleæ** Rab. in Fung. Europ. n. 1846. — Peritheciis 3072 gariis, epidermide tectis, globosis, atris; ascis cylindraceis, octosporis; sporidiis exacte mono-vel distichis, oblongis vel ovoideis, uniseptatis, fuscis, 16-18 \approx 8-9.

Hab. in ramis emortuis Pteleæ trifoliatae, Schlossgarten zu Salem Baden (JACK).

4. **Otthia Amelanchieris** Karst. Frag. XXIV, pag. 3. — Peritheciis 3073 in cæspites erumpentes stipatis, sphæroideis vel mutua pressione valde difformibus, vertice interdum depresso, ostiolo minute papillato v. obsoleto, levibus, glabris, atris, 300-400 μ . diam.; ascis cylindraceo-subclavatis, 200 \approx 20, jodo fulvescentibus; sporidiis monostichis, octonis, ellipsoideis v. oblongatis, utrinque obtusissimis, 1-septatis, ad septum constrictis, fuligineis, opacis v. semipellucidis, 30-36 \approx 12-18; paraphysibus flexuosis.

Hab. ad ramos siccos Amelanchieris vulgaris in horto mustialensi Fenniae.

5. **Otthia alnea** Peck — Syll. I, n. 2804. — Var. **carnosa** Cooke 3074
Grevill. XV, p. 80, *Cucurbitaria carnosa* Cooke (1876). — Sporidiis uniseptatis, demum triseptatis, hyalinis, medio constrictis,
 25 ± 5 .

Hab. in ramis Alni, Portland, Maine Amer. bor. (FULLER).

6. **Otthia Ilicis** H. Fab. Sphér. Vaucl. II, pag. 47, fig. 17. — Peritheciis 3075 in acervulos epidermide rupta cinctos dense congestis, cortici iuxta, globosis, papillulatis, $\frac{1}{2}$ - $\frac{2}{3}$ mm. latis; ascis cylindraceis, breve stipitatis, 8-sporis, parte sporifera 180 ± 18 ; sporidiis oblique monostichis, didymis, medio constrictis utrinque obtusis, fulvo-luteis, dein fuligineis, $24-30 \pm 12-15$.

Hab. ad ramulos dejectos *Quercus Ilicis*, Vaucluse Galliæ. — Differt ab *O. Quercus* ascis cylindricis nec clavatis duploque longioribus etc.

7. **Otthia ambiens** Niessl Hedw. 1881, p. 98. — Peritheciis dense cæ- 3076 spitosis, hemisphaericis v. subglobosis, coriaceo-carbonaceis, atris, obscure umbilicatis et papillatis; ascis clavatis, oblongis, amplis 130-150 \pm 40-50; sporidiis 8, inordinate 2-3-stichis, oblongis vel parum semilanceolatis vel elongatis piriformibus, inæqualiter diblastis, uniseptatis, parte superiori multo majori, inferiori minori, attenuatis, atro-fuscis, 45-52 \pm 15-20; parapysibus numerosis ramosis, guttulatis.

Hab. in ramis siccis *Betulae albæ* in Germania.

8. **Otthia Wistariæ** Pass. Diagn. Fung. nuovi III, n. 16. — Cæspitulis 3077 parvulis, erumpentibus, subrotundis vel rarius oblongis, longitudinaliter subseriatis; peritheciis stromate corticali fusco insidentibus, subglobosis, opacis, ostiolo punctiformi v. obsoleto; ascis cylindricis, paraphysatis, 8-sporis, 105 ± 8 ; sporidiis monostichis, ellipticis, uniseptatis, non v. vix constrictis, fuscis, $12,5 \pm 7,5$.

Hab. in ramis siccis *Wistariae sinensis* in horto Botanico Parmensi Ital. bor. — *O. diminutæ* Karst. affinis, sed ascii breviiores, paraphyses non ramosæ et sporidia latiora.

MELANCONIELLA Sacc. Syll. I, p. 741.

1. **Melanconiella biansata** (Ell. et Ev.) Berl. et Vogl. Add. Syll. p. 128, 3078
Melanconis biansata Ell. et. Ev. in Torr. bot. Club. 1883, p. 118. — Peritheciis in stromate *M. Meschuttii* simili circinatis, globosis, 0,75 mm. diam., mox collapsis, 6-10; ostiolis brevibus, quadrangularibus vel apice imperfecte pertusis, in discum acute ellipsoideum vel orbicularem, sordide brunneum, epidermidem perforantem

et ea rupta cinctum, collectis; ascis 190-200 \approx 30, evanidis (octosporis?); sporidiis oblongo-ellipsoideis, uniseptatis, mox brunneis, sæpius utrinque appendicula brevi, tuberculiformi, obtusa ornatis, 35-45 \approx 15-19 (sine appendiculis).

Hab. in ramis Betulæ, Plainfield Amer. bor.

2. **Melanconiella apocrypta** Ell. Amer. Nat. Febr. 1883, p. 195. — 307⁹
Peritheciis 8-12 subcircinantibus, $\frac{1}{2}$ mm. diam., membranaceis, cortice interiore nidulantibus, stromate obsoleto, epidermide vix pustulatim elevata tectis; ostiolis brevibus, fasciculatis; ascis subclavatis, 114 \approx 22 v. elongatis, 120-150 \approx 12; sporidiis distichis v. monostichis, ellipsoideis, 25-30 \approx 11-13, strato mucoso hyalino obvolutis, dein brunneis.

Hab. in ramis emortuis Populi, Decorah Iowa Americæ borealis (E. W. HOLWAY). — Status conidicus forte refert *Melanconium populinum* Peck.

3. **Melanconiella leucostroma** (Niessl et Rehm) Sacc. *Melanconis leucostroma* Niessl et Rehm Ascom. n. 674, Hedw. 1883, p. 40. — 3080
Stromate pustulato v. subapplanato, vix limitato, superficie albo-pruinoso; peritheciis monostichis, 6-9 circulari-aggregatis v. sub-inordinatis, subglobosis, minutis, ostiolis convergentibus, fasciculatis; ascis elliptico-oblongis, 70 \approx 15, septulato-paraphysatis, octosporis; sporidiis 2-3-stichis, elongato-obtusis, rectis, 1-septatis, 2-nucleatis, constrictis, 18-24 \approx 5-7, fuscescentibus, utrinque appendicula filiformi hyalina 6-8 μ . longa auctis.

Hab. in ramis dejectis Fagi silvaticæ prope Brünn Moraviae (NIESSL).

4. **Melanconiella Meschuttii** (Ell. et Ev.) Berl. et Vogl. Add. Syll. 3081
pag. 129, *Valsa Meschuttii* Ell. et Ev. in Bull. Torr. bot. Club. 1883, pag. 117. — Peritheciis 10-20, circinatis, in stromate lenticulari, nigro, orbiculari v. ellipsoideo, superficie interiore corticis, incidente immersis, ostiolis brevibus, cylindraceis, in discum soridide brunneum, epidermidem transverse erumpentem collectis, apice distincte 4-rimoso; ascis circiter 75 \approx 10 p. sp.; sporidiis distichis, ellipsoideis, leniter curvis, 14-16 \approx 6-8, uniseptatis, vix constrictis, primum subhyalinis, demum brunneis, utrinque appendicula hyalina, corniformi, mox evanida ornatis.

Hab. in ramis emortuis Betulæ, Plainfield Americæ borealis (GEORGE F. MESCHUTT). — An eadem ac var. *major* sequentis?

5. **Melanconiella decorahensis** Ellis Am. Nat. 1883, Febr. p. 195. — 3082
Peritheciis subglobosis, coriaceis, $\frac{1}{2}$ mm. diam., 8-12 circinantibus,

stromate corticali exceptis, epidermideque tectis, ostiolis vix prominentibus, in discum ellipticum, erumpens, sordide griseum conjunctis; ascis cylindraceis, breve stipitatis, 95-115 \approx 9-11 p. sporif.; sporidiis monostichis, ellipsoideis, obtusis, 1-septatis, olivaceis, 15-20 \approx 8-10; conidiis in stromate discreto, obovatis, olivaceo-nigricantibus, ascosporas subæquantibus.

Hab. in ramis emortuis *Betulæ*, Decorah Iowa Americæ borealis (E. W. HOLWAY). — *Var. major* Ellis Journ. Myc. 1887, p. 42: sporidiis longioribus nempe 16-26 \approx 8-10, plerumque biseriatis, cetera eadem. In ramis emortuis *Betulæ*, Plainfield N. J. Amer. bor. (G. F. MESCHUTT).

6. **Melanconiella acrocystis** (Peck) Berl. et Vogl. Add. Syll. I, p. 128, 3083

Valsa acrocystis Peck in 33 Rep. St. Mus. p. 34, t. 2, f. 19-22.

— Pustulis minutis, epidermide tectis et eam leniter elevantibus atque in lacinias acutas transverse rumpentibus; peritheciis saepius 4-12 in quaque pustula circinatis, cortice interiore insidentibus, tomento griseo-brunneo tectis, ostiolis brevibus, nigris, obtusis, seriatim dispositis; ascis oblongis; sporidiis latis, stipatis v. distichis, oblongis, coloratis, uniseptatis, 40-62 \approx 12-20, utrinque appendicula hyalina, subglobosa v. sphæroidea praeditis.

Hab. in ramis emortuis *Betulæ* lentæ, North Greenbush Amer. bor.

MASSARIOVALSA Sacc. Mich. II, p. 569. — Perithecia cortice immersa, circinantia. Asci paraphysati, 4-8-spori. Sporidia uniseptata, ovato-oblonga, fusca, strato hyalino crasso obvoluta. — Est *Valsa* sporidiis *Massariellæ* instructa, unde etymon.

1. **Massariovalsa sudans** (B. et C.) Sacc. *Massariella sudans* (B. et C.) Sacc. Syll. I, p. 717. — Adde: in cortice *Quercus albæ*, Newfield N. J. (ELLIS). — Asci teretes 180-200 \approx 20-25, apice obtusi lumineque coarctato, octospori; paraphyses guttulatæ; sporidia monosticha oblongo-ellipsoidea, constricto-1-septata, olivaceo-fuliginea, 35-40 \approx 15-16, strato crasso hyalino obvoluta.

VALSARIA Ces. et De Not. — Sacc. Syll. I, p. 741.

1. **Valsaria majuscula** Cooke et Harkn. in Grevillea XIII, p. 17. 3084
— Tecta; pustulis sparsis vix prominulis; peritheciis 6-10 compositis, demum cuticulam perforantibus, ostiolis brevibus obtusis; ascis amplis; sporidiis majusculis, ellipticis, medio constrictis, utrinque rotundatis, uniseptatis, fuscis, 50 \approx 25.

Hab. in ramis Salicis in California (HARKNESS).

2. **Valsaria Hookeri** (Cooke) Berl. et Vogl. Add. Syll. I, p. 130, *Me-* 308*logramma Hookeri* Cooke in Grevill. XIII, p. 103. — Acervulis

erumpentibus, discoideis, atris; peritheciis prominentibus, minimis, confluentibus vel subdiscretis; ascis clavatis; sporidiis lanceolatis, uniseptatis, valde constrictis, fuscis, 35-45 \approx 7.

Hab. in corticibus South Marocco (HOOKER).

3. **Valsaria collematoides** (B. et Rav.) Berl. et Vogl. Add. Syll. p. 131, 308*Botryosphaeria collematoides* B. et Rav. in Grevillea XIII, p. 102.

— Stromate effuso, tenui, atro; peritheciis parvulis, ovalibus, congestis, demum apice appланato, opaco, botryoideo-aggregatis, subconfluentibus; ascis cylindraceis, octosporis; sporidiis ellipticis, uniseptatis, vix constrictis, fuscis, 15 \approx 7.

Hab. in corticibus in America boreali.

4. **Valsaria Caproni** (Cooke) Berl. et Vogl. Add. Syll. p. 131, *Pseu-* 308*dovalsa Caproni* Cooke in Grevillea XIV, p. 48. — Pustulis in

ligno immersis; peritheciis subglobosis, atris, circinantibus sursum in colla brevia convergentia productis; ascis cylindraceis, octosporis; sporidiis uniseriatis, ellipticis, medio uniseptatis, nec constrictis fuscis, 15 \approx 5.

Hab. in ligno denudato quercino, Shere Britanniæ (D. CAPRON).

5. **Valsaria nigrispora** (Peck) Berl. et Vogl. Add. Syll. p. 129, *Dia-* 308*trype nigrospora* Peck in 33 Rep. Stat. p. 33. — Stromate minu-

to, subtili, orbiculari, pustulato, nigrescente v. nigro, epidermidem elevante et eam stellatim v. transverse rumpente; peritheciis par-

cis, saepe 6-12, ligno insculptis stromateque sursum tectis; ostio-

lis vix prominulis, discum subcinereum v. nigrescentem perforan-

tibus; ascis subcylindraceis, 8-sporis; sporidiis stipatis vel distichis,

oblongo-ellipsoideis, obtusis, saepe leniter curvulis, uniseptatis, 16 \approx 8, loculis crasse 1-guttatis, primum pallidis, demum coloratis.

Hab. in ramis emortuis Betulæ luteæ, Quaker Street Ameri-

cae borealis.

6. **Valsaria Sarraziniana** Sacc. et Roum. in Rev. Myc. 1883, t. 39, 3089

f. 8. — Stromate late effuso sed vix manifesto, corticali; perithe-

ciis hinc inde 3-4 aggregatis v. subsparsis, globosis, nigris, $\frac{3}{4}$ mm.

diam., omnino abilitis; ostiolis brevibus, punctiformi-perforantibus;

ascis cylindraceis, longiuscule stipitatis, 100-170 \approx 10; paraphy-

sibus filiformibus obvallatis, octosporis; sporidiis oblique 1-stichis,

oblongis, constricto-1-septatis, loculo superiore paullo crassiore,

20-22 \approx 7-7,5, olivaceo-fuligineis.

Hab. in cortice Alni glutinosæ in Arduennis (LIBERT). —
Valsariæ fœdanti videtur affinis.

7. **Valsaria salicina** Ell. et Ev. Phil. Proc. 1890, p. 236. — Stromate 3090 subovato, 2-2,5 mm. diam. cortice immerso, supra intus pallide colorato, tuberculi ad instar brunneo-nigri, subhemisphærici, 1-2 mm. lat., fere 1 mm. alt. prominulo; peritheciis 10-20 in quoque stromate, irregulariter congestis, ovato-globosis, contextu crasso, supra in colla gracilia, 1 mm. l. v. ultra contractis, ostiolis papilliformibus, subprominentibus; ascis gracilibus, 70-80 \approx 5-6 p. sp.; paraphysibus copiosis; ascis longioribus; sporidiis oblique monostichis, oblongis, confertis, cylindraceis, 2-guttatis, brunneis, 1-septatis, 10-12 \approx 3,5.

Hab. in ligno emortuo Salicis, London Canada Amer. bor.
— *V. anthostomoidi* affinis.

8. **Valsaria exasperans** (Ger.) Sacc., *Diatrype exasperans* Ger. Bull. 3091 Torr. Cl. V, pag. 39. — Stromate olivaceo-virente; ascis cylindraceis; sporidiis ovatis, obtusis, 1-septatis, constrictis, 15 \approx 7, fuligineis.

Hab. in cortice Fagi ferrugineæ in Amer. bor.

9. **Valsaria purpurea** Peck in Torr. bot. Club. 1884, p. 28. — Pu- 3092 stulis erumpenti-emergentibus, tomento purpureo tectis; peritheciis 6-20 in quaque pustula, stipatis, subglobosis, nigris; ostiolis tomentum perforantibus, rostratis, cylindraceis vel elongato-conoideis, rugosis, sæpe curvis vel flexuosis, nigris; ascis cylindraceis, paraphysatis, 75-110 \approx 10; sporidiis uniseriatis, coloratis, oblongo-ellipsoideis, uniseptatis, 15-16 \approx 7,5-8.

Hab. in cortice Fraxini, Canada Amer. bor. (MACOUN).

10. **Valsaria stellulata** Romell in Hedwigia 1885, pag. 263. — Stromate 3093 eutypeo; ostiolis emergentibus, astroideis; ascis clavatis, longissime pedicellatis, 8-sporis, 135-150 (parte sporif. 60-90) \approx 10-12; sporidiis botuliformibus, utrinque rotundatis, medio (rarius infra medium) 1-septatis (rarissime inæqualiter 2-septatis), non constrictis, primo olivaceis, continuis et guttulatis, dein fuscis et vix guttulatis, sed distincte septatis, rectis v. curvulis, 16-25 \approx 4-6.

Hab. in ligno truncorum Fraxini excelsioris pr. Upsala in Suecia (ROMELL).

11. **Valsaria Phoradendri** (Berk. et Curt.) Berl. et Vogl. Add. Syll. 3094 p. 130 *Melogramma Phoradendri* B. et Curt., Curt. Cat. 143, Cooke in Grev. XIII, p. 103. — Peritheciis gregariis, tectis, demum cortice fisso-rimoso erumpentibus, atris, obtusis; ascis subcy-

lindricis; sporidiis subellipticis, uniseptatis, leniter constrictis, fuscis, 28-30 \approx 10-12.

Hab. in cortice *Phoradendri* flavescentis in Amer. bor.

12. **Valsaria grandinea** (Berk.) Berl. et Vogl. Add. Syll. p. 130. *Melogramma (Valsaria) grandinea* Berk. nec *Diatrype grandinea* B. et Rav. *Melogramma insidens* Berk. in Grev. IV. 99, nec Schweinitz (ita Cooke in Grevillea XIII, p. 103). — Stromate effuso, subrotundo, fusco-nigro; peritheciis emergentibus, confertis, ovatis, apice depresso subumbilicatoe; ascis clavatis; sporidiis ellipticis, uniseptatis, nec constrictis, 8 \approx 4, fuscis, diu continuis.

Hab. in cortice *Fraxini* in America boreali.

13. **Valsaria Bignoniæ** (Schw.) Cooke in Grevillea XIII, p. 39, *Sphæria Bignoniæ* Schw. Syn. Amer. bor. n, 1310 et Sacc. Syll. II, p. 383. — Sporidia olivaceo-brunnea, uniseptata, 15 \approx 8.

14. **Valsaria Robiniæ** (Schw.) Cooke in Grevillea XIII, p. 37. *Sphæria Robiniæ* Schw. Car. n. 61 et Sacc. Syll. II, p. 369. — Sporidia elliptica, uniseptata, brunnea, 20 μ . longa.

15. **Valsaria actidia** Berk. et Rav. in Herb. Berk. Cooke in Grev. XIV, p. 14. — Stromatibus innato-emergentibus; peritheciis magnis, paucis, 4-6, ostiolis cuneiformibus, stellatim dispositis; ascis amplis (tetrasporis?); sporidiis ellipticis, uniseptatis, constrictis, fuscis, 40-42 \approx 16-18.

Hab. in ramulis *Ostryæ* et *Carpini* in Carolina inf.

16. **Valsaria atrata** Sacc. et Br. Misc. Myc. II, p. 22. n. 97 tab. XI, fig. 29. — Stromatibus gregariis, cortice initio tectis et ligno superficie late nigrificato basi adnatis, globoso-conoideis, aterrimis; peritheciis stromate immersis, paucis, globulosis, ostiolis breviter emergentibus, papillatis; ascis cylindraceis, breve stipitatis, copiose paraphysatis, octosporis; sporidiis monostichis, ellipsoideo-oblongis, constricto-1-septatis, 17-20 \approx 8-11, initio flavicanti-olivaceis, 2-guttulatis, dein opace fuligineis.

Hab. in caulinis emortuis *Brassicæ* *oleraceæ* acervatis, Troyes Galliæ (BRIARD).

17. **Valsaria pustulans** (E. et E.) Sacc. *Diatrype pustulans* E. et E. Journ. Mycol. 1888, p. 80 (ex errore 68). — Stromatibus aplannatis, e matricis substantia vix mutata formatis, cuticula nigrificata, pustulatim elevata, demum ostiolis papilliformibus subprominentibus perforata, tectis, $1\frac{1}{4}$ - $1\frac{1}{2}$ cm. lat., sed usque ad 2-4 cm. v. ultra plus minus confluentibus, area linea nigra limitata; peritheciis membranaceo-coriaceis, subglobosis v. leniter applanatis, ma-

gnitudine media, in quoque stromate 8-12; ascis gracilibus, 75-85 \approx 6-7, subsessilibus; paraphysibus filiformibus; sporidiis monostichis, oblongis, 1-septatis, constrictis, utrinque angustatis, rectis, brunneis, 10-12 \approx 3.

Hab. in caulis emortuis Arundinariæ, St. Martinsville, Louisiana Amer. bor. — Adest socia *Coniothyrii* species, peritheciis minutis immersis sporulisque exiguis, 2 μ . diam. brunneis.

18. **Valsaria atro-fusca** (Schw.) Cooke in Grev. *Sphæria* Schw. Syll. II, n. 4227, Herb. Berk. n. 9925. — Sporidia uniseptata, 10-12 μ . long. — Sec. Cooke ad gen. **Valsarium** pertinere videntur: *Hypoxylon gemmatum* B. et Rav. (Syll. I, n. 1369) (sporid. 1-septatis), *Hypoxylon? Aceris* C. et E. (Syll. I, n. 1423).

*Sect. 6. Phæophragmiæ *).*

MASSARIA De Not. — Sacc. Syll. I, p. 2, Cfr. Berl. Ic. I, p. 23.

1. **Massaria occulta** Romell in Hedwigia 1885, p. 262. — Peritheciis sparsis vel subaggregatis, cortice totis immersis, nigris, intus atris, minutis circ. $\frac{1}{5}$ mm. diam., ostiolis vix emersis, nudo oculo puncta alba in peridermio levi (non elevato) offerentibus; ascis cylindraceis, breviter stipitatis, paraphysatis, 8-sporis, 130-180 (parte sporif.) \approx 10-12; sporidiis monostichis, oblongis, obtusis, saepe nonnihil inæquilateralibus, primo 4-guttulatis, guttulis globosis, deinde guttulis transverse ovalibus v. transverse fusiformibus demum 3-septatis, fuligineo-olivaceis, 20-28 \approx 8-9, vix vel non hyalinis obvolutis.

Hab. ad ramos emortuos Ribis Grossulariæ prope Kumla, Nericia in Suecia (L. ROMELL).

2. **Massaria Winteri** Rehm Hedw. Ascom. Ill. sub n. 698, *M. Corni* 3100 Kunze F. sol. n. 338 nec Auct. — Peritheciis minutis, fuscidulis, sub epidermide nidulantibus eamque infuscantibus et elevantibus; sporidiis fusiformibus subacutis, rectis, medio constrictis, 2-cellularibus, quaque cellula nucleis 2 magnis instructis, hyalinis, 18 \approx 3,5, 8, biseriatis, in ascis clavatis, 60 \approx 18; paraphysibus ramosis.

Hab. in ramis Corni pr. Zürich Helvetiæ (WINTER).

*) Cfr. A. N. Berlese *Icones fungorum ad usum Sylloges Saccardianæ adcommodeatæ*, Asculi Picen. 1890 et seq. opus gravissimi laboris et sane utile, ubi species plures ex autoptica comparatione delentur vel accuratius illustrantur. Hic tantum genera et species novas afferre potui.

3. **Massaria Destreæ** Oud. Contr. Mycol. XIII, p. 29, t. 8, f. 13. — 3101

Peritheciis gregariis, nonnumquam in circulos dispositis cortici immersis, tandem vertice erumpentibus ostioloque punctiformibus dehiscentibus, 1-1,5 mm. diam.; ascis non visis; paraphysibus numerosissimis, filiformibus; sporidiis oblongis, saturate fuscis, 80-84 \approx 18-21, strato gelatinoso, ampio obvolutis, 3-septatis, septo mediano prominente.

Hab. in ramulis Aceris Pseudoplatani, in silva Schéveningue Hollandiæ (Domina DESTRÈE).

4. **Massaria conspurcata** (Wallr.) Sacc. Syll. f. II, p. 782, Bomm. et 3102

Rouss. Contr. Myc. II, p. 16, Berl. Ic. t. 13, f. 3. — Peritheciis globulosis, $\frac{1}{2}$ mm. diam., solitariis v. 2-4-confertis, ligno immersis, epidermide tumidula tectis, nucleo gelatinoso, virente; ostiolo brevissimo, perforante sed vix exerto; ascis tereti-clavatis, sessilibus, 4-sporis, 225-260 \approx 27, p. sporif. 198-204 \approx 27; paraphysibus copiosis longissimis; sporidiis fusoideis, initio hyalinis guttatis, dein 1-3-septatis, 6-8-guttatis, olivaceis tandem brunneis, medio secedentibus, strato mucoso obvolutis, 69-85 \approx 14-18.

Hab. in ramis emortuis Pruni Padi in Germania et Belgio.

— Adest stat. spermogonicus (*Dendrophoma*), sporulis ovoideis, 2-guttatis, 3 \approx 1, basidiis alterne ramosis et st. pycnid. *Hender-*
sonia conspurcata S. B. R.

5. **Massaria fagicola** Allesch. Sudbay. Pilze p. 190, t. II, fig. 1. — 3103

Peritheciis approximatis, majusculis, granuloso-asperatis, nigris, epidermide sœpe lacerata tectis; ascis elongatis, stipitatis, crasse tunicatis, 8-sporis, circ. 160-180 \approx 28-36; sporidiis initio distichis, elongatis, 2-septatis, sœpius inæquilateris, ad septum medium constrictis, primo flavo-brunneolis, dein brunneis, loculis 1-guttatis, muco involutis, 40-50 \approx 16-20; paraphysibus ramulosis.

Hab. in ramulis dejectis Fagi pr. München Bavariæ. — Ad sunt in eadem matrice spermatia cylindracea, utrinque obtusa, sœpe incurvata, hyalina, circ. 10-12 \approx 2-3, nec non stylosporæ fusoideæ, stipitatæ, subincurvæ, pluriseptatæ, articulis 1-guttatis, nigro-brunneis, loculis extimis plus minus hyalinis, circ. 140-160 \approx 12-16.

6. **Massaria cleistotheca** Harkn. New Calif. Fungi p. 24. — Peri- 3104
theciis minutis, tectis; ascis 8-sporis, piriformibus vel obovatis, crasse tunicatis, sessilibus, 48 \approx 30; paraphysibus agglutinatis; sporidiis hyalinis, æqualiter biconoideis, dein triseptatis, strato mucoso obvolutis, 32-40 \approx 8-10.

Hab. in caulis emortuis *Dendromeci rigidii*, Tamalpais in California. — «Si sporidia constanter hyalina, aptius sub *Massarina* locanda».

7. **Massaria pulchra** Harkn. New Calif. Fungi p. 24, Berl. Fc. t. 15. 3105 f. 2. — Peritheciis sparsis, tectis, nucleo albo farctis, $1-1\frac{1}{2}$ mm. diam.; ascis octosporis, late clavatis, 126 ± 36 ; sporidiis fusiformi-navicularibus, irregulariter et inæqualiter biconoideis, muco obvolutis, primum uniseptatis hyalinis, tandem brunneis, inæqualiter 3-5-pseudo-septatis, $58-60 \pm 20-22$.

Hab. in ramulis emortuis *Umbellulariæ californicæ*, Sausalito Californiæ.

8. **Massaria umbrosa** (Niessl) Rehm in litt. *Leptosphaeria umbrosa* 3106 Niessl in Herb. Rabenh. Fungi Europ. n. 1934 et Rehm Ascomyc. fasc. XIV, p. 13, n. 690. — Peritheciis sub epidermide leniter rufescente nidulantibus, minutis, subglobosis, siccitate parum collapsis, atris, coriaceis, ostiolo brevi, papilliformi; ascis oblongis, clavato-oblongis v. subcylindraceis, stipite brevi, abrupto, octosporis, $70-100 \pm 11-16$; sporidiis distichis, fusiformibus, utrinque obtuse rotundatis, rectis curvatisve, 4-septatis et parum constrictis, dilute virescentibus, demum subolivaceis, denique sæpe fuscis, $32-34 \pm 7-9$, muco tenui obvolutis.

Hab. in caulis exsiccatis *Spirææ Arunci*, *Gentianæ asclepiadæ*, *Lysimachia vulgaris* etc. in Germania.

9. **Massaria eryngiana** G. Delacroix Bull. Soc. Myc. 1890, f. IV. — 3107 Peritheciis immersis, atris, basi applanatis, circiter 200μ . diam., collo papillato, subconico, prominente; ascis paucis, ovoideis, leviter pedicellatis, 115 ± 65 , octosporis, mox deliquescentibus; sporidiis primum continuis, hyalino-granulatis, multiguttatis, dein septatis, sed plerumque septo medio uno eoque bene constrictis, interdum fuscidulis, postremo fuscis, ovato-cylindraceis, strato hyalino angustiore circumductis, quinquesepbatis, ad septa non vel vix constrictis, loculis una guttula magna præditis, $90 \pm 25-28$; paraphysis citissime dilabentibus.

Hab. in foliis et caulis exsiccatis *Eryngii campestris*, Saône et Loire Galliæ.

10. **Massaria Holoschœni** Passer. Diagn. F. N. III, n. 17. — Peritheciis insculptis, sparsis, vix papillatis, globoso-depressis, atris, amplis, saccato-oblongis, rectis v. curvis, basi abrupte breviterque oblique stipitatis, paraphysatis, 8-sporis, $150-175 \pm 25-37$; sporidiis bi-tristichis, fusiformibus, medio leniter constrictis, rectis, 9-11

762 Pyrenomyceteæ, Sphæriaceæ, Phæophragmiæ, Leptosphaeria.
septatis, chlorino-hyalinis, tandem fuscis, circulo hyalao circum-
datis, $32-42 \approx 8-10$ (sine circulo).

Hab. in caulis fracidis *Scirpi Holoschoeni*, socio *Lophio-*
tremate pusillo, Vigheffio pr. Parma Ital.

11. **Massaria olivacea** Cooke Grev. XVII, p. 92, *Sphæria olivaceo-*
hirta Schwein. Amer. Bor. n. 1656, Syll. II, 4353. — Sporidia bi-
serialia, lanceolata, 3-5-septata, fusca, $50-60 \approx 12-16$, primitus ocel-
lato-nucleata, medio-constricta.

12. **Massaria distincta** Cooke Grev., XVII, p. 92, *Sphæria distincta*
Schwein. Amer. Bor. n. 1655, Syll. II, n. 4359. — Sporidia bise-
rialia, 5-septata, fusca, $70-80 \approx 16-18$, medio constricta, muco hya-
lino primo obvoluta.

LEPTOSPHÆRIA Ces. et De Not. — Sacc. Syll. II, p. 13.

I. In Dicotyledoneis cauli-ramicolæ.

1. **Leptosphaeria Doliolum** (Pers.) Ces. et De N. — Syll. II, p. 14. — 3109
Var. **dissimilis** Rehm Ascom. Ill. n. 888. — Ascis cylindraceis,
apice rotundatis, 8-sporis, $75-90 \approx 7-8$; sporidiis plerumque ellipti-
cis, rectis, obtusis, 4-loclaribus, loculo utroque apicali minore,
loculo secundo multo majore, ad septa valde constrictis, vix nu-
cleatis, melleis, $12-15 \approx 5-5,5$, 1-stichis; paraphysibus subramosis.

Hab. ad caules putridos Heraclei palmati pr. lacum Ze-
noga alpium Transsylvan. (LOJKA). — Var. **angustispora** Pat. in
Rev. Myc. 1886, p. 180: a typo differt sporulis rufescentibus, tri-
septatis, minoribus rectioribusque, $16-20 \approx 3$. In caulis emortuis
Pedicularis debilis, Yunnan Sinarum (AB. DELAVAY).

2. **Leptosphaeria Longchampsi** (West.) Sacc. in Rev. Myc. 1885, 3110
Sphæria Longchampsi West. in March. Crypt. Belg. — Perithe-
ciis primo tectis, dein epidermide secedente liberis, atris, minutis,
globosis globosove-conoideis, ostiolo conico, longiusculo, acuto præ-
ditis, rugosiusculis $\frac{1}{4}-\frac{1}{3}$ mm. diam.; ascis cylindraceis, sursum ro-
tundatis, deorsum attenuato-stipitatis, paraphysatis, octosporis, $70-$
 $80 \approx 10-12$; sporidiis distichis, cylindraceis, utrinque rotundatis, 4-
septatis, loculo secundo leniter crassiore, luteo-viridulis, $25-30 \approx$
 $4-4,5$.

Hab. in caulis exsiccatis Libanotis vulgaris in Belgio.

3. **Leptosphaeria Eryngii** H. Fab. Sphér. Vaucl. p. 49, f. 19. — Pe. 3111
ritheciis sparsis, tectis, globosis, $\frac{1}{3}-\frac{1}{2}$ mm. latis, epidermidem ma-
tricis nonnihil inflantibus; ostiolo erumpente, punctiformi, vix pro-
minulo; ascis clavatis, longissime et sensim stipitatis, paraphysibus

crassis, articulatis, guttulatis, superantibus circumvallatis, 8-sporis, 150-180 \approx 18-21; sporidiis cylindraceo-elongatis, utrinque conoideis curvulis, 3-septatis, ad septa constrictulis, hyalinis vel pallide luteis, loculis tenue granulosis, 42-50 \approx 8-9.

Hab. ad caules et petiolos exaridos *Eryngii campestris*, Vaucluse Galliæ.

4. **Leptosphæria fœniculacea** H. Fab. Sphér. Vaucl. II, p. 61, fig. 24. 3112
— Peritheciis sparsis, globosis, cortici infossis, $\frac{1}{4}$ mm. latis; ostiolo brevi, crassiusculo, epidermidem vix superante; ascis clavatis, paraphysatis, breve stipitatis, 8-sporis, 130-140 \approx 14; sporidiis subdistichis, fusoideis, hyalinis vel pallidissime flavis, 4 et rarius 5-septatis, loculo secundo in plerisque, tertio in nonnullis crassiore sub eoque constrictulis, 40 \approx 6.

Hab. ad caules emortuos *Fœniculi vulgaris*, Vaucluse Galliæ.

5. **Leptosphæria diaporthoides** Wint. Contr. Flor. Myc. Lus. VI, 3113 p. 11, n. 844. — Peritheciis omnino immersis, semper tectis, serialiter dispositis, depresso globosis, ostiolo brevi, conico, erumpente, membranaceis, atris, glabris, 360-420 μ . latis; ascis exacte clavatis, longe pedicellatis, apicem versus parum attenuatis, 100-120 \approx 17-18, octosporis; sporidiis oblique 2-3-stichis, elongato-fusiformibus, parum curvatis, semper tri septatis, ad septa constrictis, loculo medio parum turgidoire, terminalibus longe attenuatis, pallidissime flavidulis, 36-40 \approx 6-7; paraphysibus filiformibus.

Hab. ad caules sicclos Umbelliferarum, Serra de Montesinho pr. Braganca Lusitanæ (MOLLER). — «An *Metaspheeria*?».

6. **Leptosphæria dolioloides** (Auersw.) Karst. — Sacc. Syll. II, p. 44. 3114
Var. **inops** Karst. in Hedwigia 1884, p. 4. — Peritheciis sparsis sphæroideo-depressis, epidermide tectis, dein denudatis, papilla punctiformi emergente, parte inferiore filamentis tortuosis, fuligineis vel fuscis obsessis, 0,2 mm. diam.; ascis cylindraceo-clavatis, 90-95 \approx 9-10, octosporis; sporidiis tristichis, fusoideo-bacillaribus, utrinque attenuatis, levissime curvulis, spurie 2-septatis, plerumque loculo tertio protuberante, flavescensibus, 32-42 \approx 4-5; paraphysibus gracilibus, subramosis, guttulatis. In caulis *Anthrisci silvestris*, Mustiala Fenniæ. — Var. **Cirsii** Karst. l. c.: peritheciis applanatis, demum subumbilicatis, basi filamentis brevibus, fuscis obsessis; ascis 88-105 \approx 9-12; sporidiis 30 \approx 4.

Hab. in caulis *Cirsii* in Fennia.

7. **Leptosphæria Phytolaccæ** Cav. Mat. Lomb. p. 11, t. I, fig. 3. — 3115

Peritheciis gregariis, sub epidermide initio nidulantibus, dein liberas, globoso-depressis, demum collapsis, patelliformibus, rugosoplicatis, basi filamentis dematiaceis præditis, 300-350 μ . diam.; ostiolo prominulo, 55-65 μ . ampio; ascis cylindraceo-clavatis, apice rotundatis, membrana crassa, stipite nullo, 65-90 \approx 9-11; sporidiis cylindraceis v. fusoideis, rectis aut leniter curvulis, 3-septatis, loculis medianis crassioribus, chlorinis, extimis conicis, obtusis, pallidioribus, omnibus 2-pluriguttulatis, 28-36 \approx 5-6,5; paraphysibus filiformibus, copiosis, ascos æquantibus.

Hab. in caulis vetustis Phytolaccæ decandræ, Torre d'Isola pr. Pavia Italiæ. — *L. platycarpæ* affinis.

8. **Leptosphaeria Lycii** Pass. in Brun. Mat. myc. Saint. p. 17. — Peritheciis sparsis v. approximatis, tectis, pustuliformibus, brunneonigris, ostiolo papillato, brunneo, erumpente, basi hyphis dematioideis, numerosis, cinctis; ascis elongatis, clavatis, basi longe attenuatis, 8-sporis, 75 \approx 10; paraphysibus filiformibus, ramosis, numerosis obvallatis; sporidiis oblique monostichis vel subdistichis, oblongo-subfusoideis; 3-septatis, rectis vel subcurvis, brunneis, 12,5 \approx 5.

Hab. in ramis emortuis Lycii barbari, Saintes Galliæ. — Differt a *L. Coniothyrio* peritheciis basi pilosis, ascis longe stipitatis, paraphysibus ramosis.

9. **Leptosphaeria monotis** Rehm Ascom. Ill. n. 887. — Peritheciis sparsis, immersis, dein erumpentibus et sessilibus, globosis, 300 μ . diam., in papillulam conoideam crassam, vix perspicue pertusam, 30-40 \approx 30, protractis, carbonaceis, nigro-fuscis; ascis clavatis, 8-sporis, 90-100 \approx 1,5; sporidiis fusoideis v. fusoideo-clavatis, dimidia superiore parte latiore, rectis v. curvatis, 6-8-locularibus, vix nucleatis, utrisque apice breviter, hyalino-appendiculatis, flavo-fuscis, 21-24 \approx 5-6, distichis; paraphysibus teneris, subramosis.

Hab. ad cæspites emortuos Saxifragarum juxta glaciem æternam montis Ortler in Tirolia.

10. **Leptosphaeria lasiosphaerioides** Starb. et Grev. Bidr. Sver. Ascom. p. 7, t. I, f. 3. — Peritheciis gregariis, 2-3-4-confertis, maculas indeterminatas, epidermide longe angusteque elevata, plus minus griseas formantibus, globosis v. conoideo-globosis, demum paullum collapsis, totis contextis pilis, ad basin præsertim longis, ramosis, fuscis, obsessis, ostiolo conico, glabro epidermidem perrumpentibus, demum denudatis, 200-300 μ . diam.; ascis cylindraceo-clavatis, vix pedicellatis; paraphysibus gracillimis obvallatis, 72-85 \approx 8-12,5;

sporidiis octonis, elongato-fusoideis, leniter curvulis v. vix inæquilateralibus, 4-septatis, primum vulgo ad septa non constrictis, loculo secundo superiore crassiore, dein saltem plerumque ad septa constrictis et loculo secundo, aliis breviore, inflato, guttulatis, utrinque obtusiusculis, dilutissime lutescentibus, 27-32 \approx 5-7.

Hab. in caulis Aconiti Lycoctoni aridis, Jemtland Sueciæ (GREVILLIUS).

11. **Leptosphaeria Vitalbæ** Niessl Tausch-Verein 1869. — Peritheciis 3119 ligno immersis, ostiolo brevi, recto, erumpente præditis, 1 mm. lat.; ascis clavatis, 8-sporis, 180 \approx 18; sporidiis fusoideis, vix acutis, rectis vel subcurvis, flavo-brunneis, 10-loclaribus, loculis 4, rarius 3 multo majoribus, subinde 1-guttatis, exappendiculatis, 36-40 \approx 7-8, distichis.

Hab. in ramis decorticatis Clematidis Vitalbæ pr. Lofer in Salzburg (NIESSL).

12. **Leptosphaeria Weberi** Oudem. Contr. à la Flore myc. de Nowaja Semlja p. 11, t. I, f. 7. — Peritheciis hypophyllis, minutis 100-150 μ . diam., primitus infra epidermidem absconditis, postremo papilla obtuse conica prominentibus, sparsis, nigris; ascis maturis cylindraceo-clavatis, brevissime stipitatis, parum curvatis, paraphysatis, 60-70 \approx 14-16; sporidiis, ut videtur, tristichis, fusoideis striatis v. plerumque falcatis, utrinque obtusiusculis vel subacutis, tri-septatis, 25-37 \approx 5-6, junioribus dilutius, senioribus saturatiis oliveaceis, loculo secundo in multis ampliore.

Hab. in Ranunculo nivali var. sulphureo, Nowaja Semlja Maris glacialis arctici.

13. **Leptosphaeria Ranunculi** Rost. Fung. Groenl. p. 558. — Peritheciis caulinis, epidermide tectis, dein denudatis, sphæroideo-depressionis; ascis crasse cylindraceo-clavatis, 95-105 \approx 30-33, tunica crassiuscula; sporidiis octonis, distichis, oblongo-elongatis, 3-septatis, loculo secundo leviter tumido, ad septa leniter constrictis, flavis, 30-35 \approx 8-10.

Hab. in caulis Ranunculi affinis in Groenlandia.

14. **Leptosphaeria quadrisepata** Trail Fung. Hardang p. 491. — Peritheciis sparsis, epidermide tectis, globosis, $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam., ostiolo brevi; ascis 8-sporis, subclavatis, pedicellatis, 60-80 \approx 8-9; sporidiis distichis v. tristichis, fusoideis, sæpe inæquilateralibus et incurvis, semper 4-septatis, loculo secundo prominente, 23-32 \approx 4-5, pallide brunneis, totis loculis guttulatis; paraphysibus numerosis, gracilibus, hyalinis, pluriguttulatis; ascis longioribus.

Hab. in caulis emortuis Thalictri minoris, Valerianæ officinalis, Lactucæ muralis et Rumicis Acetosæ, Jordal et Vik Norvegiæ. — *L. fæniculacea* affinis.

15. **Leptosphaeria Resedæ** Passer. Diagn. F. N. III, n. 18. — Peri- 3123 theciis subgregariis, hypophloëis, tandem cortice consumpto nudatis, ligno adnatis, globosis, atris, ostiolo minuto, papillari, interdum depresso; ascis elongato-clavatis, paraphysatis, 8-sporis, 75-100 = 10-12; sporidiis elongato-fusiformibus, distichis, apicibus acuminatis, triseptatis, loculis æqualibus, 30-35 = 5, primo hyalinis, tandem flavidis.

Hab. in caulis fracidis Resedæ luteæ in agro parmensi Italiæ. — *Leptosphaerice Bocconiae* videtur affinis, sed sporidiis nunquam fuscis visis; ceterum hujus descriptio nimis contracta iudicium difficile reddit.

16. **Leptosphaeria Poterii** Passer. Diagn. F. N. IV, n. 13. — Peri- 3124 theciis sparsis, tectis, ostiolo compressiusculo erumpentibus; ascis elongato-clavatis; paraphysibus tenuibus stipatis, basi longe attenuatis, 8-sporis, 50-70 = 6-7; sporidiis fusiformibus, apicibus acutis, 3-4-septatis, lutescentibus, 13 = 2.

Hab. in caulis fracidis Poterii Sanguisorbæ sociâ Rhabdosporâ, Parma Italiæ borealis.

17. **Leptosphaeria subcæspitosa** Cooke et Harkn. in Grevillea XIV, 3125 p. 10. — Erumpens; peritheciis subglobosis, atris, subcæspitosis, 4-8 in cæspitulos congestis vel in lineas elongatas dispositis; ostioli cylindricis, elongatis, emergentibus; ascis clavatis, octosporis; sporidiis lanceolatis, medio constrictis, quinqueseptatis, fuscis 35-38 = 7.

Hab. in caule Geranii in California (HARKNESS).

18. **Leptosphaeria Delavayi** Pat. in Rev. Myc. 1886, p. 82. — Pe- 3126 ritheciis $\frac{1}{4}$ - $\frac{1}{2}$ mm. diam., concavis, nigris, ostiolo non papillato, primum epidermide tectis, demum denudatis; ascis 70-90 = 9-12, 8-sporis; sporidiis elongatis, fusiformibus, 3-4-septatis, pallide bruno-fuligineis, 40-45 = 6; paraphysibus filiformibus, ascorum longitudinem æquantibus.

Hab. in caulis emortuis Primulæ Sikkimensis Likiang Yunnan Sinarum (AB. DELAVAY).

19. **Leptosphaeria multiseptata** Winter. — Sacc. Syll. II, Add. p. LVII, 3127 var. **alpina** Rehm in Hedwigia 1885, p. 235. — Peritheciis gregariis vel sparsis, primitus sub epidermide elevata nidulantibus eamque papillula perforantibus, demum liberis, subglobosis, papillula

minuta, haud perspicue perforatis, in parenchyma fuscum contextis, atris, circ. 0,3 mm. diam., rarissime basi hyphis fuscis, parvulis cinctis; ascis clavatis, 8-sporis, circ. 109-12-14; sporidiis 3-stichis, flavidulis, fusiformibus, subobtusis, rectis, 11-(rarissime 12) cellularibus, cellula quinta semper latiore, minus cellula sexta, quaque plerumque nucleolis nonnullis, parvulis prædita, ad septa subconstrictis, 50-5; paraphysibus filiformibus.

Hab. ad caules sicclos Epilobii Fleischeri juxta glaciem æternam montis Ortler prope Sulden in Tirolia (DR. REHM).

20. **Leptosphaeria Acanthi** Pat. in Rev. Myc. 1886, p. 181. — Peritheciis sparsis, primo epidermide tectis, dein liberis, applanatis, ostiolo vix protuberante; ascis cylindraceis, breve stipitatis, 106-110-9-10; paraphysibus filiformibus, fere gelatinosis, cinctis; sporidiis monostichis, fusiformibus, rectis, demum brunneo-rufis, triseptatis, ad septa parum constrictis, 20-23-6-7.

Hab. in caulis emortuis Adathodæ (an potius Acanthi?) Yunnan Sinarum (AB. DELAVAY).

21. **Leptosphaeria oreophiloides** Sacc. et Penz. Syll. II, p. 20, — Var. 3129 **Scrophulariæ** Karst. in Hedw. 1883, n. 12, pag. 2. Rev. atque Synop. Ascom. fenn. p. 51. — Ascis circ. 75-9; sporidiis octonis, elongato-fusoideis, rectis vel leviter curvulis, subinde loculo secundo superiore levissime crassiore, 30-40-4; paraphysibus eguttulatis.

Hab. in caulis Scrophulariæ nodosæ emortuis ad Mustiala Fenniæ.

22. **Leptosphaeria eutypoides** Peck in 38 Rep. St. Mus. p. 105. — 3130 Peritheciis numerosis, gregariis, 250-280 μ . diam. hemisphæricis vel depressis, primum epidermide tectis, demum denudatis, nigris, ostiolis papilliformibus; ascis clavatis vel subcylindraceis, 100-110-12-16; sporidiis ovoideis vel oblongis, rectis vel leniter curvulis, triseptatis, plerumque ad septa constrictis, luteo-brunneis, 20-23-8-10; paraphysibus filiformibus.

Hab. in caule emortuo Chenopodii albi, Albany Amer bor.

23. **Leptosphaeria Lecanora** H. Fab. Sphér. Vaucl. II, p. 49. f. 20. — 3131 Peritheciis sparsis, initio epidermide tectis, dein superficialibus et vertice collapso patellariformibus, $\frac{1}{4}$ mm. vel minus latis; ostiolo minutissimo; ascis cylindraceo-clavatis, breve stipitatis, 8-sporis, 70-100-10-16; paraphysibus raris vel nullis; sporidiis submonostichis, pallide melleis, oblongo-ovatis, 3-septatis, utrinque obtusis 18-24-8-9.

Hab. ad caules exsiccatos Salsolæ Kali, Vaucluse Galliæ.

24. **Leptosphaeria Oxyriæ** Rost. Fung. Groenl. p. 559. — Peritheciis 3132 gregariis, fuscis, sphæroideo-depressis; ascis crasse cylindraceo-clavatis, pedicellatis, 50-52 \approx 12-14; sporidiis distichis, fusoideo-oblongatis, 3-septatis, ad septa constrictis, flavis.

Hab. ad caules exsiccatos Oxyriæ didymæ in Groenlandia.

25. **Leptosphaeria serbica** Schroet Hedw. 1890, p. 59. — Peritheciis 3133 sparsis, epidermide tectis, globosis, gregariis, circ. 300 μ . diam., ostiolo applanato præditis; ascis numerosis, clavatis, breve stipitatis, 60-70 \approx 10-12; sporidiis octonis, distichis, ovato-fusoideis, uno latere applanatis, 4-locularibus, utrinque rotundatis, exosporio flavo-brunneo, 16-18 \approx 6-6,5.

Hab. in caulis Camphorosmæ monspeliacæ pr. Prokuplje Serbiæ.

26. **Leptosphaeria cucurbitarioides** H. Fab. Sphér. Vaucl. II, p. 50, 3134 fig. 51. — Peritheciis dense connatis, primum tectis, dein nudis, acervulos per epidermidem fissam erumpentes efformantibus, globoso-mammæformibus, $1\frac{1}{2}$ - $2\frac{2}{3}$ mm. latis, ostiolo brevi, conoideo, crassiusculo; ascis clavatis, paraphysatis, longe stipitatis, 8-sporis, 90-100 μ . long., parte sporifera 60 \approx 10; sporidiis subdistichis, oblongis, rectis v. curvulis, utrinque obtusis, 3-septatis, ad septa constrictulis, pallide luteis, 15-18 \approx 6-7.

Hab. ad caules exaridos Dorycnii suffruticosi, Vaucluse Galliæ. — Habitus fere Cucurbitariæ.

27. **Leptosphaeria pratensis** Sacc. et Br. in Rev. Myc. 1885. — Pe. 3135 ritheciis numerosis, innatis, globulosis, parum depresso, corticolis, $1\frac{1}{4}$ - $1\frac{1}{3}$ mm. diam., ostiolo obtuso, epidermidem vix nigrefactam perforante; ascis cylindraceo-clavatis, breviter stipitatis, octosporis, 68-112 \approx 8-12; paraphysibus filiformibus; sporidiis distichis v. subdistichis, fusiformibus, obtusiusculis, primum hyalinis dein luteolis, 3-septatis, septis leniter constrictis, rectis vel curvulis, 28-32 \approx 5-6, granulo-guttulatis.

Hab. in caulis exsiccatis Medicaginis sativæ, Troyes Galliæ.

28. **Leptosphaeria multiseptata** Wint. Hedw. 1872, p. 148. — Peritheciis 3136 sparsis, epidermide tectis demum erumpentibus, globosis, nigris, rugulosis, pilis pallidis, repentibus cinctis, in collum crassum, conicum, apice obtusum, peritheciū dimidium æquans attenuatis, 351 \approx 270; ascis clavato-cylindraceis, sessilibus, 8-sporis, 117 \approx 11; paraphysibus filiformibus, simplicibus, continuis multi-guttulatis, asco dimidio brevioribus; sporidiis longissime fusiformi-

bus, utrinque rotundatis, 8-11-septatis, guttulatis, ad septa constrictis, pallide luteolis, 50-65 \approx 5.

Hab. ad caules aridos *Lathyrus silvestris* prope Oederan, Saxonia (G. WINTER).

29. **Leptosphæria Lathyri** Fautr. Rev. Myc. 1891, p. 8. — Peritheciis sparsis v. approximatis, majusculis, ostiolo elongato erumpentibus; sporidiis olivascentibus, cylindraceis, attenuatis, rectis vel subcurvulis, 8-11-septatis, ad septum secundum v. tertium constrictis, 50-60 \approx 4.

Hab. in *Lathyrus silvestris*, Noidan, Côte-d'Or Galliæ (F. (FAUTREY).

30. **Leptosphæria hæmatites** (Rob.) Niessl. Cfr. Hedw. 1883, p. 10. Sic nominanda *Didymella hæmatites* (Rob.) Sacc. Syll. p. 553 (*Sphærella hæmatites* Cooke Bot. Journ. 1883, pag. 109), nam sporidia matura 24-27 \approx 5, 3-septata, flava.

- 31 **Leptosphæria Sarraziniana** Sacc. et Roum. in Rev. Myc. 1884, 3138 tab. LV, f. 2. — Peritheciis innato-erumpentibus, globoso-papillatis, mox collapso-umbilicatis; ascis clavato-fusoideis, 85-95 \approx 7-10, octosporis; sporidiis distichis, fusoideis, 3-septatis leniter constrictis, obtusiusculis, guttulatis, olivaceis, 18-21 \approx 5-6.

Hab. in nodis Caryophyllaceæ cujusdam, Senlis Galliæ. — Ab affinis *L. pyrenopezizoide* differt sporidiis rectis, crassioribus, loculo secundo vix inflato etc.

32. **Leptosphæria Stellariæ** Rich. Cat. Champ. Marn. n. 1329. — Pe. 3139 ritheciis sphæricis, nigris, subapplanatis, sparsis; ascis clavatis; paraphysibus filiformibus, longissimis; sporidiis fusoideis, elongatis, acutis, 3-septatis, pallide luteolis.

Hab. in caulis vetustis *Stellariæ mediæ*, St. Amand, Marne Galliæ.

33. **Leptosphæria Vahlii** Rost. Fungi Groenl. p. 557. — Hyphis di- 3140 lute fuscis, articulis brevibus inflatis; peritheciis sphæroideo-depressis, vertice umbilicatis, papillatis; ascis numerosis, elongato-clavatis, longe pedicellatis, 90-100 \approx 11-13; paraphysibus copiosis, filiformibus; sporidiis distichis, cymbiformi-fusoideis, fuscis, 5-septatis 20-25 \approx 6-7, loculis 1-2-guttulatis.

Hab. ad caules exsiccatos *Melandrii triflori* in Groenlandia.

34. **Leptosphæria agnita** Sacc. Syll. II, p. 40. — Var. **Chrysanthemi** 3141 Berl. in Soc. Ven. Trent. 1885, tab. XI, fig. 2. — Peritheciis minutis 0,3-0,4 mm. diam., saepe complanatis; ascis crassioribus, 70-90 \approx 12-14; sporidiis tristichis, typice 5-septatis, 45-48 \approx 6.

Hab. in caulis siccis Chrysanthemi corymbosi, Eichenwald prope Sugenheim Franconiae (REHM). — Var. **major** Berl. in Soc. Ven. Trent. 1685, tab XI, fig. 4: a typo differt peritheciis majoribus, $\frac{1}{2}$ mm. diam. et distincte papillatis. In caule emortuo Chærophylli temuli in Gallia (MORTHIER). — Var. **Erigerontis** Berl. in Soc. Ven. Trent. 1885, tab. XI, fig. 3: ascis majoribus 82-90 \approx 15; sporidiis tristichis valde elongatis, 55-50 \approx 4-5, demum 6-septatis, loculo tertio superiore non vel vix turgidiore, utrinque sensim attenuatis. In caule sicco Erigerontis canadensis in Amer. bor. (ELLIS). — Var. **ambigua** Berl. in Soc. Ven. Trent. 1885, tab. XI, fig. 5: peritheciis epipermide velatis, obtuse breviterque papillatis, atris, nitidis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., globo-so-depressis; ascis cylindraceo-clavatis, 80-110 \approx 12; sporidiis irregulariter tristichis vel distichis, leniter biconoideis, loculo tertio superiore parum crassiore, utrinque obtusis, 6-septatis, 35-48 \approx 6, flaveolis. In caule Eupatorii cannabini in Germania. — Var. **Bupleuri** Sacc. Myc. Sib. p. 15, f. 11: peritheciis obtusiusculis, $\frac{1}{3}$ mm. latis; ascis fusoideo-clavatis substipitatis, 75-80 \approx 9-10; sporidiis fusoideis, utrinque acutis 5-septatis, 30-35 \approx 4-5, ad septa levissime constrictis, subolivaceis, loculis subinde senio secedentibus. In caulis emortuis Bupleuri falcati pr. Minussinsk Sibiriae (MARTIANOFF).

35. **Leptosphaeria purpurea** Rehm Ascom. Lojk. n. 69, p. 54. — Pe- 3142
ritheciis gregariis, primitus epidermide tectis, dein ea secedente liberis, in ligni longe lateque purpureo-tincti superficie insidentibus, hemisphaericis, integris, papilla minutissima pertusis, glabris, atris, c. 0,3 mm. diam., hyphis creberrimis, longiusculis, fuscis, basi cinctis; ascis clavatis, 8-sporis, 70 \approx 8; sporidiis fusoideo-elongatis plerumque rectis, 4-cellularibus, cellula secunda multo latiori, cellulis nucleo 1 magno praeditis, melleis, 21 \approx 3-3,5, distichis; paraphysibus ramosis.

Hab. ad caules sicclos Artemisiæ vulgaris in horto Eszterházyano, Kaposvár Hungariae (LOJKA).

36. **Leptosphaeria ophioboloides** Sacc. in Misc. Myc. I, p. 23. — Pe- 3143
ritheciis gregariis, epidermide velatis, globosis, $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam., atris, ostiolo papillato erumpente; contextu parenchymatico densiusculo; ascis clavatis, breve stipitatis, apice obtusiusculis, 120-130 \approx 14, parce paraphysatis, octosporis; sporidiis subtristichis, bacillaribus, utrinque obtuse rotundatis, 8-10-septatis, loculo secundo

vel tertio paullo crassiore, 50-55 \approx 4-4,5, olivaceo-melleis, loculis sæpius minute biguttulatis.

Hab. ad caules *Tragopogonis*, ut videtur, in silvis pr. Neu-châtel Helvetiæ (MORTHIER). — Sporidia subinde bina apicibus connecta, interdum in loculos secedentia, hinc ophioboloidea.

37. **Leptosphaeria derasa** (B. et Br.) Thüm. — Sacc. Syll. II, p. 41. 3144

— Var. **alpestris** Rehm in Hedwigia 1885, pag. 235. — Peritheciis gregariis, primitus sub epidermide caulium nidulantibus eamque papillula protuberante perforantibus, demum liberis, sessilibus globosis, in papillulam rotundam, brevem protractis, haud perspicue perforatis, atris, in parenchyma fusco-nigrum contextis, circ. 0,3-5 mm. diam., hyphis plus minusve longis, circ. 4 μ . cr., undulatis, septatis, fuscis, crebris obsessis; ascis clavatis, 8-sporis, 90-120 \approx 12; sporidiis 2-3-stichis, rectis vel subcurvatis, fusiformibus, subacutis, plerumque 9-cellularibus, itaque cellula quarta latiore, demum 10, rarissime 11-cellularibus et cellula quinta lassisima, quaque nucleolis 2-parvulis et sporidiis 10-cellularibus utraque apice appendicula filiformi, curvata, brevi instructis, flavidulis, 36-45 \approx 4-5; paraphysibus filiformibus, nucleolatis.

Hab. ad caules siccis Compositæ? ad muros viæ jugi Stelvii prope Franzenshöhe in Tirolia (Dr. REHM).

38. **Leptosphaerla carduina** Passer. Diagn. F. N. III, n. 19. — Peritheciis sparsis, pusillis, globosis, epidermide rupta cinctis, atris, basi filis dematiaceis parcis præditis, contextu minute celluloso, fuligineo; ascis paraphysatis, elongato-clavatis, inferne breviter attenuato-stipitatis, 8-sporis, 68-75 \approx 10-12; sporidiis distichis, elongato-ellipticis, utrinque rotundatis, 3-5-septatis, 15-18 \approx 5, loculo uno alterove sæpe longitudinaliter diviso, e flavo fuscidulis.

Hab. in bracteis calathidum dejectarum Cardui nutantis, Vigheffio pr. Parma Italiæ.

39. **Leptosphaeria bella** Passer. Diagn. F. N., I, n. 46. — Peritheciis 3146 subgregariis v. sparsis, punctiformibus, atris; ascis oblongo-clavatis, paraphysatis, 8-sporis, 45-65 \approx 10-12; sporidiis distichis, oblongo-ellipticis, triseptatis, non v. vix constrictis, loculo pænultimo interdum paullo ampliore, flavis, 12-15 \approx 5.

Hab. in ramulis fracidis Chondrillæ juncæ, Vigheffio pr. Parma Italiæ.

40. **Leptosphaeria bicuspidata** (Cooke et Hark.) Berl. et Vogl. Add. 3147 Syll. p. 133, *Sphaeria bicuspidata* Cooke et Hark. in Grevillea XIII, p. 19. — Sparsa, tecta; peritheciis innatis, subprominulis,

globosis, atris, ostiolo brevi; ascis clavatis, octosporis; sporidiis fusiformibus, triseptatis, constrictis, fuscis, utrinque hyalino-cuspidatis, 25-8 (sine cuspidibus).

Hab. in caulis Baccharidis in California (HARNESS).

41. **Leptosphaeria Galiorum** Sacc. Syll. II, p. 22. — Var. **Lampsanæ** Sacc. et Br. in Rev. Myc. 1885. — Peritheciis sparsis, globosis, epidermide tectis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam.; ostiolis papillatis, erumpentibus; ascis clavatis, sursum attenuatis, deorsum longe stipitatis, octosporis, 64-69 = 14-16 p. sp.; paraphysibus filiformibus, guttulatis; sporidiis sursum distichis basi monostichis, ovoido-oblongis, obtusis, 2-septatis, septis constrictis, brunneis, 20-24 = 7, rectis vel leniter curvulis. — In caulis Lampsanæ communis, Troyes Galliæ.

42. **Leptosphaeria corynospora** Berl. et Bres. Micr. Trid. p. 32, t. III, 3148 f. 4. — Peritheciis gregariis, globoso-conicis v. parum depresso-depressis, in ostiolum papillatum, nitidum, distinctum desinentibus, 300-350 μ . diam., atris, epidermide pustulatim inflata tectis; ascis eximie clavatis, in stipitem longum abeuntibus; paraphysibus filiformibus, guttulatis cinctis, 110-120 = 10-12; sporidiis octonis, stipe distichis v. inferne 1-stichis, clavulatis, 3-septatis, parte superiori crassiore, inferiore tenuiori, fere caudiformi, utrinque, sed inferne magis, acutiusculis, pallide virescentibus, loculo secundo inflato, polaribus pallidioribus, 24-26 = 6, guttuligeris.

Hab. in caulis emortuis Cirsii lanceolati pr. Trento.

43. **Leptosphaeria dubia** Sacc. et Paol. in Sacc. Myc. Sibir. pag. 14, 3149 fig. 13. — Peritheciis laxe gregariis, epidermide velatis, globoso-depressis, breve papillatis, dein umbilicatis, nigris, $\frac{1}{4}$ mm. diam.; ascis fusoideo-clavatis, utrinque obtusulis, subsessilibus, 75-90 = 10, obsolete paraphysatis, octosporis; sporidiis 2-3-stichis, recte fusoidis, utrinque obtusulis, triseptatis non vel vix constrictis, pallide fuligineis, 25-30 = 3-4,5.

Hab. in caule Valerianæ dubiæ in lapidosis pr. Minussinsk Sibiræ (MARTIANOFF).

44. **Leptosphaeria Morthieriana** Sacc. Misc. Myc. I, p. 23. — Peritheciis laxe gregariis, parenchymate immersis, globulosis, duruisculis, atris, $\frac{1}{6}$ mm. diam., ostiolo conoideo, acuto, emergente; ascis teretibus, crassis, brevissime abrupte stipitatis, 140-150 = 35-37, paraphysatis, apice subrotundatis, octosporis; sporidiis distichis, oblongo-fusoideois, rectis, utrinque rotundatis, 40-45 = 13-14, 5-septatis, vix constrictis, lutescentibus.

Hab. in foliis Succisæ putrescentibus in palude torfacea les Ponts prope Neuchâtel Helvetiæ (MORTHIER).

45. **Leptosphaeria Molluginis** Passer. Diagn. F. N. IV, p. 5. — Pe- 3151
ritheciis sparsis, tectis, erumpentibus, subglobosis, contextu minute
celluloso, fuligineo, ostiolo pertuso; ascis clavatis, nodulo basilari
sessilibus, paraphysatis, 8-sporis, $60-65 \approx 8-10$; sporidiis distichis
v. oblique monostichis, fusiformibus, utrinque acutis, rectis, trise-
ptatis, rarissime 4-septatis, loculo subultimo vix tumidulo, $15 \approx 5$,
melleis v. flavidulis.

Hab. in caule sicco Galii Molluginis, Vigheffio pr. Parma
Italiæ. — Sporarum parvitate et colore ab aliis congeneribus ga-
liicolis distinguenda.

46. **Leptosphaeria Eranthemi** Pat. in Rev. Myc. 1886, pag. 181. — 3152
Peritheciis sparsis epidermide tectis, applanatis, orbicularibus, ostiolo
conoideo donatis; ascis $63-70 \approx 8-10$, paraphysibus filiformibus ob-
vallatis; sporidiis fusiformibus, rectis vel parum curvatis, fuligi-
neis, $16 \approx 5$, quinquesptatis, raro 3-4-septatis ad septa, præcipue
ad medium, constrictis.

Hab. in pedunculis floralibus Eranthemi, Yunnan Sinarum
(Ab. DELAVAY). — Sporidia sæpe protoplasmate longitudinaliter di-
viso, muriformia apparent.

47. **Leptosphaeria Plumbaginis** Pat. in Rev. Myc. 1886, p. 181. — 3153
Peritheciis nigris, sparsis, globulosis, ostiolo vix emergente prædi-
tis; ascis cylindraceis, breve pedicellatis, $110 \approx 15$, paraphysibus
filiformibus, crebre septatis cinctis; sporidiis primo linearibus, hya-
linis, medio uniseptatis, dein brunneo-luteis, 8-12-septatis, medio
vix constrictis, $50 \approx 6-7$.

Hab. in caulis emortuis Plumbaginis, Yunnan Sinarum
(Ab. DELAVAY).

48. **Leptosphaeria fallaciosa** Berl. Exc. Frioul p. 10, t. 9, fig. 5. — 3154
Peritheciis sparsis, primo epidermide tectis, dein ea secedente sub-
liberis, globoso-depressis, tandem umbilicatis, ostiolo papillato, mi-
nuto præditis, $200-300 \mu$. diam., nigris, nitentibus; ascis oblongo-
clavatis, sæpe stipite curvulo, brevi, noduloso præditis, 8-sporis,
 $70 \approx 14$, paraphysatis; sporidiis subtristichis, cylindraceis, fere asci
longitudinem æquantibus, constanter 7-septatis, loculo quarto su-
periore incrassato, $47-50 \approx 4-5$, rectis v. curvulis, initio guttulatis,
utrinque rotundatis, flavesceni-viridulis.

Hab. in caulis emortuis Satureiæ hortensis Osoppo in agro

- 774 Pyrenomyceteæ, Sphæriaceæ, Phæophragmiæ, Leptosphæria.
 Foro Juliensi Ital. bor. — *L. agnitæ*, *L. acutæ*, *L. petiolicolæ* et
L. derasæ affinis.
49. **Leptosphæria Salviæ** Passer. Diagn. F. N. III, n. 20. — Peri- 3155
 theciis hypophlœis, subgregariis, globosis, parce setulosis, ostiolo cylindrico corticem perforante; ascis paraphysatis, clavatis, inferne longe attenuatis, 8-sporis, 87-125 \approx 12-14; sporidiis fusiformibus, 9-11-septatis, subdistichis, 45 \approx 6, flavidulis, loculo altero ex intermediis subtumido.
- Hab.* in ramis siccis *Salviæ officinalis*, interdum socio *Ophiobolo homosporo* in horto botanico parmensi Italiæ.
50. **Leptosphæria Brunellæ** Ell. et Ev. Proc. Phil. 1890, p. 237. — 3156
 Peritheciis sparsis, minutis, $\frac{1}{5}$ - $\frac{1}{4}$ mm. diam., epidermide subelevata, ostiolo papilliformi pertusa tectis; ascis clavato-cylindraceis, 75-80 \approx 10-12, subsessilibus, filiformi-paraphysatis; sporidiis distichis, fusoideis, subincurvis, pallide flavidio-brunneis, 3-septatis, loculo subultimo inflato, 22-30 \approx 4.
- Hab.* in caulis emortuis *Brunellæ vulgaris*, London Canada Amer. bor. (DEARNESS). — Differt a *L. pyrenopezizoidi* peritheciis non collabentibus, a *L. Parietaricæ* sporidiis pallidioribus. Adsunt socia perithecia spermogonica (*Rhabdospora*), intus pallida, majuscula, sporulis acicularibus 40-55 \approx 2-2,5 fœta, nec non *Phoma* quædam sporulis 4 \approx 1,5.
51. **Leptosphæria Steironematis** Ell. et Ev. Proc. Phil. 1890, p. 237. 3157
 — Peritheciis gregariis, subepidermicis, conico-hemisphæricis, 0,5 mm. diam., epidermidem ostiolo conico pertusam elevantibus; ascis clavato-cylindraceis, 75-100 \approx 15-20, copiosis; sporidiis distichis, oblongis, 3-septatis, ad septa interdum subconstrictis, brunneis, utrinque obtusis, plerumque subincurvis, 15-22 \approx 7-8.
- Hab.* in caulis *Steironematis ciliati*, London, Canada Amer. bor. (J. DEARNESS).
52. **Leptosphæria demissa** Niessl Contr. Fl. Myc. Lus. pag. 13. — 3158
 Peritheciis sparsis, tectis, depresso hemisphæricis seu collapsis, minutis, 0,2 mm. diam., atris, glabris, coriaceis, ostiolo brevi conico, fimbriato, ut in *L. modesta*, erumpente; ascis clavatis, in stipitem brevem attenuatis, 8-sporis; sporidiis distichis, fusiformibus, rectis vel curvatis, utrinque obtusiusculis, 4-cellularibus vel 3-septatis, loculo secundo parum inflato, dilute lutescentibus, subhyalinis, 15-19 \approx 3-3 $\frac{1}{2}$; paraphysibus ascorum longitudine, tenuibus, ramosis.
- Hab.* ad caules aridos Vincæ mediæ, Coimbra Lusitaniæ. — Sporidiorum forma et magnitudine a *L. modesta* distincte diversa.

53. **Leptosphaeria euphorbicola** Brun. Champ. ajout. Saint. p. 244. 3159

— Peritheciis sparsis v. approximatis, globosis, nigris, tectis, dein erumpentibus, ostiolo papillato; ascis cylindraceo-clavatis, 90-100 \approx 10, filiformi-paraphysatis; sporidiis octonis, distichis, oblongo-fusoideis, rectis v. subincurvis, flavis, 5-7-septatis, ad septa subconstrictis, 26-34 \approx 6-7.

Hab. in caulis emortuis Euphorbiæ pilosæ, Saint-Sever pr. Saintes Galliæ.

54. **Leptosphaeria brachysperma** Berl. Exc. Frioul p. 8, t. 9, f. 4. — 3160

Peritheciis sparsis, minutis, 150 μ . diam., epidermide tectis, globoso-conoideis, ostiolo minute papillato, pertuso vix exsertis; ascis rectis v. curvis, in stipitem brevissimum, nodulosum abeuntibus, sursum latiuscule rotundatis, 60-70 \approx 12-14; paraphysibus filiformibus; sporidiis distichis v. basi oblique monostichis, ovoideis v. oblongis, 3-septatis, non constrictis, utrinque rotundatis, loculo secundo superiore parum crassiore, 14-16 \approx 6-8, amœne flavo-olivascentibus.

Hab. in caulis herbaceis emortuis, Osoppo in agro Forojuliensi Ital. bor. — *L. galiicola* var. *brachysporæ* et *L. Sarazianæ* affinis.

55. **Leptosphaeria Winteri** Niessl in Hedw. 1883, p. 1. — 3161

Peritheciis sparsis, globosis, minutis, circa 0,14 mm. diam., submembranaceis, atris, glabris, tectis, vertice erumpentibus, ostiolo crasso, cylindraceo, peritheciæ diametrum æquante; ascis oblongis, demum elongatis, subsessilibus, 90-120 \approx 25-34, 8-sporis; sporidiis farctis 2-3-stichis, plerumque parum curvatis, utrinque late rotundatis, 6-cellularibus seu 5-septatis, loculo secundo inflato, 36-44 \approx 10-12, lutescentibus; paraphysibus paucis, vix superantibus ramulosis.

Hab. ad folia arida Plantaginis alpinæ, Albulapass, prope Weissenstein Rhætiæ.

56. **Leptosphaeria Cynops** H. Fab. Sphér. Vaucl. II, p. 50, fig. 23. — 3162

Peritheciis parce gregariis aliis corticicolis et infixis, tectis, aliis lignicolis et superficialibus, nudis, conideo-mastoideis, $\frac{1}{3}$ mm. latis; ostiolo brevi, papillato, nitido; ascis clavatis, abrupte et breve stipitatis, paraphysatis, 8-sporis, 100-120 \approx 15; sporidiis subdistichis, fusiformibus, 5-septatis, medio vix vel non constrictis, pallide flavis; loculis 1-guttulatis, 50 \approx 6.

Hab. ad caules emortuos Plantaginis Cynopis, Vaucluse Galliæ.

57. **Leptosphaeria vagabunda** Sacc. — Syll. II, p. 31. — ***alvariensis** 3163

Sterb. Ascom. Oeland. p. 13, f. 4. — Peritheciis sparsis, conico-sphæroideis, ostiolo papillato, paullum prominulo v. conoideo-cylindraceo, interdum lateraliter compresso, altitudine peritheciorum dimidio breviore præeditis, 200-250 μ . diam.; ascis clavatis, vix pedicellatis, 80-100 \approx 20-26; sporidiis distichis, primitus hyalinis, simplicibus, multiguttulatis, dein æque hyalinis, medio constricto-uniseptatis, sensim olivascentibus, 4-magniguttulatis, demum 3-septatis, fuligineo-fuscis, loculis externis dilutioribus, 20-30 \approx 10-12; paraphysibus numerosissimis, gracilibus, agglutinatis.

Hab. ad rhizomata, terrâ non tecta, Plantaginis maritimæ var. gentilis in Allvaren pr. Næsby Oelandiæ. — Var. **Dulcamaræ** Karst. Symb. Fenn. XXIX, pag. 34: sporidiis 3-septatis, luteolo-fuscescentibus, 14-19 \approx 5-6. In ramis emortuis Solani Dulcamaræ prope oppidum Jakobstad Fenniæ.

58. **Leptosphaeria obesula** Sacc. Syll. II, p. 15, Mout. Ascom. Liég. II, p. 8, t. I, f. 12. — In diagnosi Syll. I. c. corrigi: ascis obclavatis, apice attenuato-rotundatis, basi brevissime crasse stipitatis, mox omnino sessilibus et late rotundatis, 45-60 \approx 20-30; paraphysibus obsoletis. Nascitur quoque in Angelica, Spiræa, Acere Pseudoplatano, Beaufays, Forêt Belgii (MOUTON).

59. **Leptosphaeria brasiliensis** Speg. F. Puigg. n. 280. — Maculis 3164 amphigenis, subdeterminatis, 5-15 mm. diam., fusco- vel cinere-scenti-arescentibus; peritheciis innato-erumpentibus, confertiusculis, epiphyllis, globoso-conoideis, 150-180 μ . diam.. atris, levibus; ascis mox diffluentibus; sporidiis triseptatis, subfusiformibus, olivaceis, 25-30 \approx 6-8.

Hab. ad folia viva Xanthoxyleæ vel Anacardiaceæ cù-jusdam in dumetis pr. Apiah. — An *Hendersonia*?

60. **Leptosphaeria odora** (Cooke et Hark.) Berl. et Vogl. Add. Syll. 3165 p. 137, *Sphæria odora* Cooke et Hark. in Grev. XIII, p. 19. — Tecta, subsparsa; peritheciis mediis, globosis, atris, in cortice nidulantibus, epidermide convexa tectis, ostiolo punctiformi; ascis clavatis, octosporis; sporidiis lanceolatis, biseriatis, primum uniseptatis, demum triseptatis, hyalino-luteolis, 22-25 \approx 5.

Hab. in ramis Umbellulariæ in California (HARKNESS).

61. **Leptosphaeria Ceanothi** (Cooke et Hark.) Berl. et Vogl. Add. Syll. 3166 p. 136, *Sphæria Ceanothi* Cooke et Hark. in Grev. XIII, p. 19. — Gregaria, tecta; peritheciis minutis, subglobosis, epidermide velatis, ostiolo brevi, punctiformi; ascis breviter clavatis, octosporis; sporidiis sublanceolatis, triseptatis, hyalino-luteolis, 16-18 \approx 4,5.

Hab. in ramulis *Ceanothi* in California (HARKNESS).

62. **Leptosphæria rubro-tincta** E. et E. in Journ. of Myc. I, pag. 150. 3167

— Peritheciis sparsis, membranaceis, depresso-globosis, 200-250 μ . diam., apice perforatis, primo tectis dein leniter prominentibus et epidermidem, pallide rufo-sanguineo-maculatam, vix perforantibus; ascis clavato-cylindraceis, sessilibus, 100-110 \approx 10-13, paraphysibus cinctis; sporidiis biseriatis, cylindraceis, rectis vel leniter curvatis, luteolis, 8-10-septatis, loculo medio leniter inflato, 25-40 \approx 4-5, subobtusis.

Hab. in ramis *Staphyleæ trifoliæ* in America boreali.

63. **Leptosphæria cassiæcola** E. et E. in Journ. of Myc. 1886, p. 41. 3168

— Peritheciis gregariis, minutis, $1/8$ - $1/6$ mm. diam., epidermide nigrescente, minute pustulatim inflata, demum ostiolo papilliformi perforata, tectis; ascis 70-75 \approx 8-10; sporidiis distichis, fusiformibus, leniter curvulis, 5-septatis, flavo-brunneis, non constrictis.

Hab. in pedunculis emortuis *Cassiæ*, Houston Texas Americae borealis.

64. **Leptosphæria consimilis** E. et E. in Journ. of Myc. 1886, p. 41. 3169

— Peritheciis sparsis, carbonaceo-coriaceis, $1/3$ mm. diam., primo tectis fibrasque ligni vel corticis minute pustulatim elevantibus, ostiolo papilliformi demum erumpentibus; ascis subcylindraceis, 80-100 \approx 15-20, copiose paraphysatis, octosporis; sporidiis subdistichis, cylindraceo-fusiformibus, 3-septatis, primum subhyalinis, dein luteis, demum obscure brunneis, 7-septatis, non vel vix ad septa constrictis, 28-35 \approx 8-10.

Hab. in ramulis emortuis salicinis prope Huron Dakota Amer. (NELLIE E. CROUCH).

65. **Leptosphæria Gillotiana** Sacc. et Roum. in Rev. Myc. 1883, p. 236, 3170

tab. 40, fig. 9. — Peritheciis inæqualiter gregariis, globoso-depressis, $1/2$ mm. diam., et ultra, atris, cortice innatis; ascis fusoides, breve stipitatis, 70-75 \approx 10, octosporis, paraphysatis; sporidiis distichis, oblongis, utrinque rotundatis, 3-septatis, non v. vix constrictis, 18 \approx 7, rufo-fuligineis.

Hab. in ramis corticatis Salicis in Gallia. — Affinis *Lept. platycarpæ*, a qua peritheciis minus depressis, sporidiis obtusioribus, minus constrictis, brevioribus differt.

66. **Leptosphæria californica** (Cooke et Hark.) Berl. et Vogl. Add. Syll. 3171

pag. 137, *Sphæria californica* Cooke et Hark. in Grevillea XIII, pag. 20. — Densissime gregaria, tecta; peritheciis hemisphærico-prominulis, obtusis, atris, diu epidermide velatis, demum apice

- 778 Pyrenomyceteæ, Sphæriaceæ, Phæophragmiæ, Leptosphaeria.
subnudis; ascis clavatis; sporidiis biseriatis, sublanceolatis, utrinque obtusis, 4-septatis, cellula pænultima subincrassata, pallide fuscis, 25-30 = 8.
- Hab.* in ramis et foliis Araucariæ imbricatæ, Sarothamni, Rhododendri, Evonymi in California.
67. **Leptosphaeria cirricola** Passer. Diagn. F. N. I, n. 44. — Peritheciis sparsis, superficialibus, atris, ostiolo papilliformi; ascis clavatis, basi longe attenuato-stipitatis, 8-sporis; paraphysibus tenuibus obvallatis, 65-75 = 7,5-8; sporidiis distichis vel oblique monostichis, fusiformibus, leniter arcuatis, 5-septatis, septis extimis tenuioribus et interdum obscuris, haud constrictis, loculo altero ex intermediis paullo crassiore, olivaceo-fuscis, 17-18 = 5.
- Hab.* in cirris aridis denudatis Vitis viniferæ, Vigheffio pr. Parma Italiae. — A *Leptosphaeria vineali* Passer. differt peritheciis discretis non maculicolis, crassioribus et sporidiis 5-septatis brevioribus etc.
68. **Leptosphaeria dichroa** Passer. Diagn. F. N. I, n. 45. — Peritheciis sparsis, subcutaneis, atris, sphæroideis, ostiolo minuto erumpente; ascis tubulosis vel vix clavatis, 8-sporis; sporidiis oblique monostichis v. subdistichis, oblongis, utrinque obtusis, triseptatis, non constrictis, rectis, loculis intermediis majoribus, castaneo-fuscis, interdum guttulatis, extimis multo minoribus, hyalinis, 15-17 = 5-6; paraphysibus non visis.
- Hab.* in ramis Deutziae scabré in horto botanico Parmensi Italiae bor. — *L. vagabundæ* proxima, sed sporidiis plane diversis. An novi generis typus?
69. **Leptosphaeria Spirææ** Karst. Fragm. Myc. Fenn. XXIV, p. 3. — Peritheciis solitariis v. saepius cæspitosis, subinde connatis, per peridermum erumpentibus, globulosis v. inæqualibus, solito plus minus depresso, nigris, glabris, saepe papillatis, 0,4-0,5 mm. diam.; ascis cylindraceis, 100-125 = 14-16; sporidiis octonis, distichis vel oblique monostichis, oblongatis, leviter curvulis v. rectis 3- raro 4-6-septatis, loculo secundo paullo crassiore, ad septa constrictis, dilutissime melleis v. hyalino-chlorinis, 22-30 = 8-10; paraphysibus numerosis, haud bene discretis, ascos superantibus.
- Hab.* ad ramulos deciduos aridos Spirææ sorbifoliæ in horto mustialensi Fenniae.
70. **Leptosphaeria aculeorum** Passer. Diagn. F. N. IV, p. 4. — Peritheciis hypodermeis, erumpentibus, atris, nitidulis, globosis, ostiolo brevi, acuto, contextu minute celluloso fuligineo; ascis paraphysa-

tis, clavatis, attenuato-stipitatis, 8-sporis, 70-75 \approx 10-12; sporidiis tristichis, elongato-fusiformibus, 7-9-septatis, flavis, 30 \approx 2,5-3.

Hab. in aculeis Rosæ caninæ, Vighèffio pr. Parma Italiæ.

71. **Leptosphæria Thomasiana** Sacc. et Roum. in Rev. Myc. 1883, 3176

tab. 40, fig. 10. — Peritheciis dense gregariis, globoso-conicis, obtususcule papillatis nigris, solidiusculis, initio epidermide velatis, mox liberatis (melanomoideis), vix $\frac{1}{2}$ mm. diam.; ascis teretiusculis, breviter stipitatis, parce paraphysatis, octosporis, 80-90 \approx 8-10; sporidiis oblique monostichis vel distichis, oblongo-fusoideis, 3-septatis, leniter constrictis, rectis vel curvulis, 20-22 \approx 6, fuscidulis.

Hab. in sarmenis Rubi emortuis in Gallia.

72. **Leptosphæria Ribis** Karsten in Rev. Myc. 1885, p. 106. — Pe- 3177

ritheciis sparsis, cortice nidulantibus, epidermide demum rimose fissa tectis, pustulas parvas efficientibus, sphæroideo-depressis, atris, glabris, contextu tenui, minute celluloso, fuligineo, ostiolo obsoleto, circiter 150 μ . latis; ascis cylindraceis 80-90 \approx 8-9, octosporis; sporidiis monostichis, ovoideo-vel fusoideo-oblongatis, rectis, triseptatis, ad septa non vel vix constrictis, flavidо-hyalinis, demum obscurioribus subfuscidule flavescentibus, 15-18 \approx 6; paraphysibus haud bene discretis.

Hab. in ramis exsiccatis Ribis alpini pr. Mustiala Fenniæ.

73. **Leptosphæria Harknessiana** Ell. et Ev. in Journ. of Myc. 1885, 3178

— Peritheciis sparsis vel gregariis, primum epidermide tectis, demum liberis atque superficialibus, atris, hemisphericis $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam.; ostiolo brevi, cylindraceo late pertuso; ascis cylindraceis, 100-110 \approx 10-12, octosporis, paraphysibus filiformibus obvallatis; sporidiis monostichis, ellipsoideis, flavo-brunneis, 3-septato-constrictis, 18-22 \approx 7-9, sursum obtusis, basi rotundatis.

Hab. in trunco emortuo Anamirtæ Cocculi, Emery Co. Utah Amer. bor.

74. **Leptosphæria fallax** Berl. Fung. Mor. fasc. IV. — Peritheciis 3179

minutissimis, atris, contextu parenchymatico, globoso-conoideis, ostiolo papilliformi, pertuso donatis, tectis, dein emergentibus $\frac{1}{8}$ - $\frac{1}{7}$ mm. diam., ascis cylindraceis vel clavato-cylindraceis, subsessilibus, paraphysatis, 60-70 \approx 12, octosporis; sporidiis distichis, fusoideis, 5-septatis, loculo secundo crassiore, rectis vel curvulis, pallide viridulis, 28 \approx 4-5, utrinque rotundatis.

Hab. in ramulis exsiccatis Mori albæ, Fiumicello Italiæ bor.

— Fructificatione ad *L. Yulan* accedit.

780 Pyrenomyceteæ, Sphaeriaceæ, Phæophragmiæ, Leptosphaeria.

75. **Leptosphaeria platanicola** (Howe) Sacc., *Sphaeria platanicola* Howe 3180

Bull. Torr. Bot. Cl. V, p. 42. — Peritheciis globoso-conoideis, erumpentibus; ascis cylindraceis vel subclavatis; sporidiis monostichis vivide fuligineis, 3-septatis.

Hab. in ramulis Platani in Amer. bor.

76. **Leptosphaeria Cisti** Celott. Mic. Montp. p. 32, f. 1. — Peritheciis 3181 sparsis, innato-erumpentibus, globoso-oblongis, nigris, $500 \approx 200$; ascis cylindraceis, $80-90 \approx 12-15$, paraphysatis; sporidiis octonis, 5-septatis, semilunatis, utrinque acutis, medio vix constrictis, $25-30 \approx 6-8$, fuligineis.

Hab. in ramis Cisti albidi, Montpellier Galliæ.

77. **Leptosphaeria Berberidis** Rich. Cat. Champ. Marn. n. 1330. — 3182
Peritheciis sparsis, subcutaneis, lenticularibus; ascis clavatis, paraphysibus obvallatis; sporidiis fusoideo-oblongis, 3-septatis, pallide olivaceis.

Hab. in ramulis vivis junioribus Berberidis vulgaris pr. spinarum fasciculos, St. Amand, Marne Galliæ.

78. **Leptosphaeria Caprifolii** Brun. Champ. Saint. 1887, pag. 428. — 3183
Peritheciis sparsis v. approximatis, globoso-conicis v. subglobosis, nigris, cortice tectis lignoque adnatis, vix erumpentibus, cortice secedente superficialibus, ostiolo papillato; ascis oblongo-clavatis, $60-80 \approx 12-14$, 8-sporis, paraphysatis; sporidiis distichis, oblongis, curvis, 5-septatis, constrictis, $18-22 \approx 3-3,5$, flavo-brunneis.

Hab. in sarmentis emortuis Lonicerae Caprifolii, Saintes Galliæ.

79. **Leptosphaeria Periclymeni** Oud. Contr. Myc. Pays-bas XII, p. 27. 3184
— Peritheciis sparsis, $\frac{1}{2}$ mm. diam., inter fibras immersis, peridermum tandem perforantibus, glabris, carbonaceis, subnitidis, ostiolo obtuse conico-prominulo; ascis cylindraceis, pedicellatis, paraphysibus filiformibus, immersis; sporidiis monostichis oblongis, fusoideis vel subpiriformibus, rectis vel subcurvis, 3-5-septatis, leniter constrictis, $15-25 \approx 7-8,5$, loculis submediis subinde paullo crassioribus septoque verticali vel oblique partitis, hyalinis dein olivaceo-subfuscis.

Hab. ad ramulos Lonicerae Periclymeni Schéveningue Nederlandiæ (Dom. DESTRÉE).

80. **Leptosphaeria Ericæ** (Fr.) Malbr. in litt. *Sphaeria Ericæ* Fr. 3185
S. M. Elench. 2, pag. 101, Sacc. Syll. II, pag. 422. — Perithecia sparsa vel aggregata, fusca, primo tecta, dein erumpentia; asci clavati, longe pedicellati, $85-95 \approx 10-11$ (p. sp. 60-65), paraphysa-

ti; sporidia disticha, elliptica vel oblonga, utrinque obtusa, 3-septata, loculis mediis latioribus, medio non v. vix constricta, sæpissime recta, fusca, 18-19 \approx 6-7, eguttulata.

Hab. ad truncos Callunæ vulgaris, Brionne Galliæ.

81. **Leptosphæria Kalmiæ** Peck 39 Rep. pag. 53, t. II, f. 10-14. — 3186

Peritheciis subcæspitosis, erumpentibus, 350-450 μ . diam., subhemisphæricis, crassis, nigris, ostiolis pertusis vel leviter papillatis; ascis cylindraceis, 100-120 \approx 7-9; sporidiis uniseriatis, oblongis vel subfusoideis, 3-septatis, subinde medio leviter constrictis, 16-20 \approx 6-7, coloratis; paraphysibus filiformibus.

Hab. in ramulis Kalmiæ angustifoliæ Adirondack Mount.

Amer. bor. — *Lept. vagabundæ* affinis.

82. **Leptosphæria Ailanthi** Karst. et Har. Journ. Bot. 1889. p. 206. — 3187

Peritheciis sparsis, sphæroideo-applanatis, epidermide tectis, ostiolo papilliformi, rotundato, perforante donatis, atris, circ. 200 μ . lat.; ascis cylindraceo-clavatis, 75-95 \approx 8-10; sporidiis octonis, distichis, fusoideo-bacillaribus, utrinque obtusis, rectis vel leniter curvulis, 3-septatis, ad septum medium subconstrictis, loculo pænultimo levissime protuberante, luteolis, 30 \approx 3-4; paraphysibus non visis.

Hab. ad ramulos Ailanthi glandulosæ in Gallia.

83. **Leptosphæria endophæna** Bomm. Rouss. Sacc. sp. nov. — Pe. 3188

ritheciis sparsis, corticolis, globulosis, punctiformibus, vix $\frac{1}{4}$ mm. diam., externe haud conspicuis; ascis anguste clavatis, sessilibus, 105-126 \approx 15-18; paraphysibus flexuosis, crebris; sporidiis octonis, distichis, fusoideis, obtusis, 3-septatis, medio constrictis, hyalinis, nubilosis, 4-guttatis, demum olivaceis, 22 \approx 6.

Hab. in ramis emortuis Ailanthi glandulosæ, Watermael Belgii. — Affinis *L. vagabundæ*.

II. In Dicotyledoneis folicolæ.

84. **Leptosphæria Camelliæ** Cooke et Mass. Grev. XVI, p. 5. — Ma. 3189

culis epiphyllis, arescendo fuscescentibus, vagis; peritheciis sparsis, punctiformibus, vix papillatis, pertusis; ascis subclavatis, plerumque 4-sporis, breviter stipitatis; sporidiis distichis, fusoideis, curvulis, triseptatis, vix constrictis, virescentibus, 25-27 \approx 5-6.

Hab. in foliis vivis Camelliæ in Victoria Australiæ (F. CAMPBELL).

85. **Leptosphæria Fraxini** Ell. et Ev. Journ. Myc. 1807, p. 44. — 3190

Maculis amphigenis, subtus ærugineis, supra sordide albis, margine ærugineis, parvis, orbicularibus, 1 mm. diam. v. elongatis, angustis,

2-3 mm. long.; peritheciis nigris, epiphyllis, paucis, pusillis, 75 μ . diam., erumpentibus, supra indistincte pertusis, basi rudimento mycelii præditis; ascis clavato-cylindraceis, 55-70 \times 8-10; sporidiis fusoideis, subincurvis, flavis, 3-5-sæpius 4-septatis, maturitate medio constrictis, interdum ad tota septa leniter constrictis, 20-25 \times 3,5-4.

Hab. in foliis vivis Fraxini americanæ, Columbia, Montana Amer. bor. (B. T. GALLOWAY).

86. **Leptosphæria Tini** Ell. et Ev. Journ. Myc. 1888, p. 64. — Ma. 3191 culicola, amphigena, maculis magnis cinereis, anguste rubenti-purpureo-marginatis, 2-3 cm. diam., insidens; peritheciis punctiformibus, minutis, innatis, apice subacuto extus tantum perspicuo; ascis oblongo-cylindraceis, subsessilibus; paraphysibus filiformibus, 40-50 \times 7; sporidiis distichis, fusoideis, flavidis, subincurvis, vix 3-septatis, 14-16 \times 3.

Hab. in foliis Viburni Tini, Lafayette Louisiana Amer. bor. (Rev. A. B. LANGLOIS).

87. **Leptosphæria hyperborea** (Fuck.) Berl. et Vogl. Add. Syll. p. 140, 3192 *Pleospora* Fuck. 2^e Deutsche Nordpolfahr. II, p. 92 et Oudemans Contr. Flor. Myc. Nov. Seml. p. 153, tab. II, fig. 9. — Peritheciis in foliorum aridorum pagina superiore nidulantibus, sparsis, punctiformibus, semiimmersis et basi globosa a foliorum substantia grisea, elevata cinctis, parte libera obtuse conica, siccis aplana-tis, aterrimis, perforatis; ascis oblongo-ovatis, utrinque attenuatis, 8-sporis, 60 \times 24; sporidiis in ascis media parte stipatis, oblongis, utrinque obtusis, rectis, 3-septatis, ad septum medium constrictis, 24 \times 8, flavo-fuscis.

Hab. in pagina superiore foliorum aridorum Andromedæ tetragonæ, Shannon insulæ Groenlandiæ et Spitsbergen. — «An a *L. Andromedæ* satis distincta?»

88. **Leptosphæria Physalidis** Ellis et Ev. in Torr. bot. Club. 1884, 3193 p. 42. — Peritheciis minutis, globoso-depressis, contextu celluloso, apice poro latiusculo pertusis; ascis 35-40 \times 7-8, a paraphysatis; sporidiis distichis, fusiformibus, leniter curvulis, luteolis, 3-septatis, 15-19 \times 3.

Hab. in maculis sordide albis, rotundatis foliorum *Physalis* pubescenti, prope Lexington Amer. bor. (KELLERMANN).

89. **Leptosphæria hysteroides** Ellis et Ev. in Bull. Wash. Lab. Nat. 3194 Nat. Hist. 1884, p. 4. — Peritheciis globosis, $\frac{1}{6}$ mm. diam., epidermide atrata, demum ostiolo obtuse conico perforata, tectis, sæpe

confluentibus et *Hysterium* simulantibus; ascis oblongis vel oblongo-cylindraceis, 85-112 \approx 12-15; paraphysibus apice intertextis; sporidiis biseriatis, oblongo-fusoideis, hyalinis, 4-nucleatis, saepe 3-septatis, brunneis, 20-24 \approx 7-9, ad septa constrictis, utrinque obtusis.

Hab. in foliis emortuis *Xerophylli* tenacis in Monte Paddo Amer. bor.

90. **Leptosphæria Helichrysi** Cooke in Grev. XIX, pag. 7. — Hypo- 3195 phylla, superficialis; peritheciis subglobosis, opacis, atris, in tomento nidulantibus, ostiolo impresso, pertuso; ascis clavatis; sporidiis congestis, sublanceolatis triseptatis, rectis curvulisve, fuligineis, 50-55 \approx 8.

Hab. in foliis *Helichrysi*, Natal Africæ (Wood). — «An hujus generis?»

91. **Leptosphæria salebricola** Bonn. Rouss. Sacc. sp. nov. — Peri- 3196 theciis sparsis, punctiformibus, subepidermicis, leviter prominulis, 135-150 μ . diam., ostiolo brevi conoideo; ascis clavatis, 84-90 \approx 15-18, sessilibus, octosporis, paraphysatis; sporidiis 2-3-stichis, fusoideis, granulosis (non guttulatis), 3-septatis, 35-39 \approx 6-7, obtusis, rectis, coloratis, loculo uno submedio leviter inflato.

Hab. in foliis morientibus *Stellariæ gramineæ*, Comblain Belgii. — A *L. uliginosa* macula atra deficiente, ascis sporidiisque majoribus etc. diversa.

92. **Leptosphæria Stellariæ** Rostr. Fung. Groenl. p. 557. — A proxima 3197 *Leptosphæria Silenes* differt ascis 45 \approx 10, sporidiis 20 \approx 6.

Hab. in foliis *Stellariæ humifusæ* in Groenlandia.

93. **Leptosphæria Rehmiana** Voss Myc. Carn. III, pag. 167. — Peri- 3198 theciis serialibus, sparsis, globosis, atris, glabris, sessilibus, minutissimis, membranaceis; ascis cylindraceis, sessilibus, 8-sporis. 64-66 \approx 11-13; paraphysibus filiformibus; sporidiis oblique monostichis v. subdistichis, oblongis v. late fusoideis, utrinque rotundatis, rectis v. leniter curvulis, 3-septatis, ad septa constrictis, loculo secundo protuberante, fuscis, 15-17 \approx 6-8.

Hab. in foliis emortuis *Drypidis spinosæ*, Stranje pr. Stein in Carniola super.

94. **Leptosphæria antarctica** Speg. F. Fueg. n. 228. — Maculis nulis v. fuscescentibus, indeterminatis, parvulis; peritheciis saepius gregariis v. subseriatis, immersis, non prominulis, globulosis, 200-250 μ . diam., atris, glabris, tenui-membranaceis, contextu parenchymatico fuligineo, ostiolo vix papillulato, epidermidem perforante

sed non exerto ornatis; ascis cylindraceis, antice obtuse rotundatis, crassiuscule tunicatis, late 1-foveolatis, postice attenuato-cuneatis, pedicello brevissimo, crasso, noduloso suffultis, 75-80 \approx 10, 8-sporis; paraphysibus filiformibus obvallatis; sporidiis elongato-fusoideis, utrinque obtusiuscule rotundatis, 45 \approx 3, non v. vix inaequilateralibus, 7-septatis, ad septum medium tantum constrictis, loculis granuloso-farctis, pallide olivascentibus.

Hab. verisimiliter in foliis in Fuegia Amer. austr. (Matrix loc. cit. non indicata).

III. In Monocotyledoneis.

95. **Leptosphaeria Musarum** Sacc. et Berl. Myc. aliq. Guineens. p. 3. — 3200

Amphigena, plerumque hypophylla; peritheciis gregariis, innatis, globulosis, $\frac{1}{6}$ mm. diam., ostiolo obtuse papillato, erumpente; ascis fusoideo-elongatis brevissime noduloso-stipitatis, apice obtusiusculis 60 \approx 10-12, obsolete paraphysatis; sporidiis distichis, fusoideis, rectis, rarius curvulis, utrinque obtusiusculis, 15-18 \approx 5-6, triseptatis, ad septa vix constrictis, olivaceo-fuscis.

Hab. in foliis emortuis Musæ S. Thomæ Nova Moka Afr. Occid. (Ad. F. MOLLER). — Ab affini *Lept. crastophila* differt ascis multo angustioribus sporidiisque paullo longioribus.

96. **Leptosphaeria Xiphii** Passer. Diagn F. N. I, n. 48. — Peritheciis 3201 sparsis, minutis, hypodermiis, ostiolo punctiformi atro, tandem emerso; ascis oblongo-clavatis, paraphysibus tenuibus obvallatis, 8-sporis, 62-70 \approx 9-10; sporidiis, distichis, oblongo-cylindricis, apicibus obtusis, 5-7-septatis, loculo nullo inflato, 17,5-20 \approx 5, castaneis.

Hab. in foliis languidis et subaridis Iridis fætidissimæ, Tabiano in agro parmensi Italæ. — A *Leptosphaeria Vecti* Ces. et De Not. sporidiis brevioribus, loculis æqualibus præcipue differt. An varietas?

97. **Leptosphaeria Filamentosæ** Ell. et Ev. Journ. Myc. 1888, pag. 64. 3202

— Peritheciis immersis, sparsis, parvulis, 200-225 μ . diam., depresso-globosis, supra cuticulam subelevantibus et erumpentibus, ostiolo non prominente; ascis subcylindraceis, 75-80 \approx 7-8, copiose paraphysatis; sporidiis distichis, oblongo-cylindraceis, 3-septatis, flavis, ad septa constrictis, rectis 12-15 \approx 4-5, utrinque obtusis.

Hab. in areis emortuis foliorum viventium Yuccæ filamentosæ, Newfield, New Jersey Amer. bor. — Fungus spermogonicus (*Coniothyrium concentricum*?) sporulis minutis, 4 μ . diam., subglobosis, brunneis.

98. **Leptosphæria præclara** Karst. in Hedw. 1884, n. I, p. 2 et Rev. 3203

atque Synop. Ascom. fenn. p. 54. — Peritheciis sparsis vel subgregariis, depresso sphæroideis, subcutaneo-erumpentibus, contextu parenchymatico, tenui, dilute olivaceo-fuscescente, glabris, ostiolo vix prominulo pertusis, circ. 150 μ . lat.; ascis elongato-vel oblongato-subclavatis, sessilibus, curvatis, 56-100 \approx 13-26; sporidiis octonis, distichis vel subtristichis, oblongato-fusoideis, curvulis vel inæquilateralibus, 5-, raro 3-septatis, loculo tertio subinde crassiore, ad septa leviter constrictis, lète luteolis, 20-32 \approx 8-9; paraphysibus haud bene discretis.

Hab. in caulis emortuis *Asparagi officinalis* ad Mustiala Fenniæ.

99. **Leptosphæria ruscicola** Karst. et Har. Rev. Myc. 1890 n. 47. — 3204

Peritheciis gregariis, nonnumquam conglomeratis, mox superficialibus, sphæroideis vel rotundatis, vertice applanatis vel depresso, ostiolo umbilicato, atris, circiter 0,2 mm. latis; ascis cylindraceo-clavatis, 120-130 \approx 18; sporidiis octonis, distichis, fusoideo-elongatis, 3-septatis, loculo secundo inflatulo, flavescenti vel chlorino-hyalinis, 33 \approx 6; paraphysibus filiformibus, numerosis.

Hab. in caulis putrescentibus *Rusci* in Gallia (P. HARIOT). — A *Leptosphæria Rusci* valde diversa.

100. **Leptosphæria asparagina** Karst. in Hedw. 1884, n. I, pag. I et 3205

Rev. atque Synop. Ascom. fenn. p. 55. — Peritheciis subsparsis vel gregariis, hinc inde in series dispositis, sphæroideis vel saepius sphæroideo-applanatis, hyphis repentibus, fuligineis obsessis, epidermide velatis, nigris, papilla conoidea, saepè perithecium dimidium æquante, prominula præditis, circ. 0,2 mm. lat.; ascis cylindraceo-clavatis, 95-105 \approx 9-10; sporidiis octonis, distichis, elongato-fusoideis, ut plurimum curvulis, 7-septatis, ad septa leviter vel vix constrictis, eguttulatis, dilute flavis, 24-30 \approx 4,5-6; paraphysibus filiformibus, bene discretis, ascos superantibus.

Hab. ad caules *Asparagi officinalis* emortuos in Mustiala Fenniæ. — Status spermogonicus est *Phoma asparagina*.

101. **Leptosphæria Passeriniana** Sacc. L. *Asparagi Passer.* Diagn. 3206

F. N. I, n. 47, nec Peck. — Peritheciis laxe gregariis, epidermide tectis dein nudatis, globosis, atris, ostiolo vix perspicuo; ascis oblongo-clavatis, paraphysatis, 8-sporis, 60-90 \approx 10-12,5; sporidiis fusiformi-lanceolatis, distichis, 6-9-septatis, 20-22,5 \approx 5, flavis.

Hab. in ramis siccis *Asparagi officinalis*, socia *Phoma Asparagi*, Parma Italiæ bor.

102. **Leptosphaeria Asparagi** Peck 40 Rep. p. 70. — Peritheciis late 3207 conicis, 250-350 μ . diam., epidermide perforata tectis, dein nudatis, atris; ascis clavatis v. cylindraceis, breve pedicellatis, 70-100 \approx 12-15; sporidiis oblongis v. fusoideis, confertis, 20-30 \approx 7, initio hyalinis 3-septatis, dein dilute coloratis, 5-septatis, ad septa constrictis.

Hab. in caulis emortuis Asparagi, Menands Amer. bor.

103. **Leptosphaeria translucens** Winter Contr. Fl. Myc. Lus. V, p. 15. 3208 — Peritheciis subsparsis, tectis, translucentibus, subglobosis, ostiolo punctiformi epidermidem perforantibus, fere coriaceis, atris, 0,2-0-24 mm. diam.; ascis cylindraceis, deorsum parum attenuatis, sessilibus, 75-95 \approx 10-11, octosporis; sporidiis distichis, cylindraceis, utrinque rotundatis, interdum parum curvatis, 6-cellularibus, cellula quarta (ab ascis apice numerata) perparum incrassata, mel leis, demum fuscis, 21-24 \approx 5-6; paraphysibus filiformibus, ascos superantibus.

Hab. ad folia arida Fourcroyæ, Coimbra in Lusitania (MOLLER).

104. **Leptosphaeria punctoidea** Karst. in Hedw. 1884, n. I, pag. 2, et 3209 Rev. atque Synop. Ascom. fenn. p. 53. — Peritheciis sparsis, sphæroideo-applanatis, innato-erumpentibus, subcutaneis, contextu parenchymatico, tenuissimo, olivaceo-pallescente, glabris, atris poro latiusculo pertusis, circ. 1 mm. lat.; ascis cylindraceo-clavatis, 45-75 \approx 7-11, pseudoparaphysatis; sporidiis octonis, distichis, elongato-fusoideis, curvulis, 3-septatis, eguttulatis, loculo secundo subinde leviter crassiore, hyalinis, demum dilute flavescentibus, 15-26 \approx 3-5.

Hab. in caulis emortuis Asparagi officinalis ad Mustiala Fenniae.

105. **Leptosphaeria infernalis** Niessl Contr. Fl. Myc. Lus. p. 13. — 3210 Amphigena; peritheciis dense gregariis, in maculis fuscis, sub epidermide dense nidulantibus, globosis, majusculis, 0,3 et ultra mm. diam., atris, glabris, rigidis, subcarbonaceis, ostiolo brevi, cylindrico, vix protuberante; ascis clavatis, stipite valde elongato, 8-sporis, 200-230 μ . longis, parte sp. 90-100 \approx 18-22; sporidiis distichis, oblongis vel parum lanceolatis seu medio valde constrictis, utrinque acutiusculis, rectis vel inaequilateralibus, 3-septatis, loculis mediis biguttulatis, atro-fuscis, demum subopacis, 19-24 \approx 9-12; paraphysibus ascorum longitudinem vel paullum superantibus, crassis, guttulatis et laxe ramosis.

Hab. ad folia emorta Fourcroyæ longævæ pr. Coimbra

Lusitaniæ. — Differt a proxima *L. clivensi* præcipue peritheciis minoribus, ascis elongatis, sporidiis duplo latioribus, sublanceolatis.

106. **Leptosphaeria rivularis** Bomm. Rouss. Sacc. sp. nov. — Peritheciis 3211 innato-erumpentibus, subcutaneis, sphæricis, atris, 250-270 μ . d., pertusis, hyphis, brunneis, repentibus basi cinctis; ascis clavato-oblongis, subsessilibus, $69 \approx 15-18$ v. clavatis, $90 \approx 12-15$, octosporis, paraphysatis; sporidiis distichis, chlorino-flavidis, fusoideis curvulis, utrinque obtusis 3-septatis, medio leniter constrictis, 24-25 $\approx 7-8$, loculo submedio leviter inflato.

Hab. in caulis Alismatis Plantaginis, Remouchamps Belgii.

107. **Leptosphaeria sabalicola** Ellis et Martin Amer. Nat. 1882. p. 810. 3212 — Gregaria vel sparsa; peritheciis coriaceis, solidiusculis, $\frac{1}{3}$ mm. diam., epidermide conico-tumidula et circum circa infuscata tectis; ascis clavato-cylindraceis, $57 \approx 8$; sporidiis distichis oblongis, 3-septatis, $11 \approx 3,5$, fuligineis.

Hab. in petiolis Sabal serrulatæ, Green Cove Springs, Florida Americæ bor. (Doct. MARTIN). An *Clypeosphaeria*?

108. **Leptosphaeria Magnusiana** Berl. et Sacc. Add. Syll. p. 414. — Peritheciis 3213 sparsis, minutis, 100-130 μ . diam., globosis, vix papillatis, epidermide dein ostiolo perforata tectis; ascis cylindraceis, basi abrupte breviterque stipitatis, sæpe curvis, crassiuscule tunicatis, apice subrotundatis, paraphysibus filiformibus, subinde pseudo-septulatis, longioribus cinctis, $70 \approx 18$, octosporis; sporidiis irregulariter distichis vel sursum basique monostichis, oblongis vel subcylindraceis, utrinque rotundatis, demum triseptatis, loculo secundo crassiore. fusco-fuligineis, 18-20 ≈ 6 .

Hab. in foliis exsiccatis Chamæropis humilis in insula Sicilia, socia *Sphaeropsida dothideoide*. — Affinis *L. Tritici*, a qua loculo secundo paullo inflato et ascis multo majoribus differt. et *L. elæosporæ*, a qua sporidiis conspicue crassioribus recedit, ab utroque insuper matrice.

109. **Leptosphaeria microscopica** Karst. — Sacc. Syll. II, pag. 59, 3214 var. **Calami** Karst. in Hedw. 1883, n. 12, p. 2, Rev. atque Synop. Ascom. fenn. p. 52. — Peritheciis laxe gregariis, subcutaneis, sphæroideis, raro oblongatis, subastomis, glabris, atris, 1 mm. lat.; ascis elongato-clavatis, obtusis, $66-80 \approx 12-15$; sporidiis octonis, di-vel subtristichis, elongato-fusoideis, rectis vel leviter curvulis 3-septatis, eguttulatis, loculo subultimo subinde leviter crassiore, dilute flavis, $20-24 \approx 6$; paraphysibus obsoletis.

Hab. in foliis putrescentibus Acori Calami in insula Run-sala Fenniæ.

110. **Leptosphaeria Corallorrhizæ** Peck 38 Rep. St. Mus. p. 105, t. 2, 3215
fig. 20-23. — Peritheciis numerosis, minutis, 100-125 μ , d., erum-pentibus nigris, minute ostiolatis; ascis cylindraceis, sessilibus, 50-75 \times 7-8; sporidiis stipatis vel distichis, subfusiformibus, trisepta-tis, medio leniter constrictis, brunneolis, 20-26 \times 4-5.

Hab. in caulis emortuis Corallorrhizæ multifloræ, Caroga Amer. bor.

111. **Leptosphaeria licatensis** Sacc. Syll. II, pag. 70, Var. **rupeforten-sis** Pass. in litt. Brun. in Rev. Myc. 1886, p. 206. — Ascis pa-raphysatis, 75 \times 12,5; sporidiis fusiformibus, 5-septatis, castaneo-brunneis, 25 \times 5-6. In foliis emortuis Typhæ latifoliæ in Horto botanico Rochefort Galliæ.

112. **Leptosphaeria typhiseda** Sacc. et Berl. in Rev. Myc. 1886, p. 33. 3216
— Peritheciis sparsis, minutis, primum tectis, dein epidermide per-forata emergentibus, globosis globosove-conoideis, ostiolo minuto, papillato instructis, $\frac{1}{8}$ mm. diam., atris; ascis cylindraceis, sur-sum rotundatis, basi breviter stipitatis, paraphysibus filiformibus, simplicibus obvallatis, 70-80 \times 14, octosporis; sporidiis oblique vel recte monostichis, cylindraceo-clavulatis, apice rotundatis, deorsum acutioribus, 24-28 \times 7, olivaceis, 5-6-septatis, ad septum medium vix constrictis,

Hab. in foliis emortuis putrescentibus Typhæ angustifoliæ Algeria. — *Lept. helminthosporæ* affinis, a qua præ ceteris ma-trice distinguenda.

113. **Leptosphaeria striolata** Passer. Diagn. F. N. IV, p. 5, n. 15. — 3217
Peritheciis sparsis, primo epidermide velatis, dein nudatis, minutis-simis, atris, contextu minute celluloso-fuligineo; ascis subclavatis, amplis, apice rotundatis, basi vix attenuatis, breviter noduloso-sti-pitatis, 62-75 \times 22-25; paraphysibus nullis; sporidiis octonis, sub-tristichis, clavulatis, initio pluriguttulatis, olivaceis, integris, dein fuligineis, obscure 5-7-septatis, episporio longitudinaliter subtilis-sime striato, loculo ultimo paullo crassiore, 20-37,5 \times 6-8.

Hab. in culmis aridis Scirpi Holoschoeni, Vigheffio, Par-ma Italiæ.

114. **Leptosphaeria Nardi** (Fr.) Ces. De Not. Syll. II, p. 72. — **dubiosa** 3218
Mout. Ascom. Lièg. II, p. 9. — Sparsa; peritheciis globosis, $\frac{1}{5}$ mm. d., innatis, nonnullis hyphis obductis, papillis minimis epidermi-dem perforante instructis; ascis teretibus, sessilibus, 70-80 \times 11-14;

sporidiis subtristichis, fusoideis, acutatis, luteolis, $35 \approx 4$, non constrictis, 5-septatis.

Hab. in foliis *Scirpi silvatici* pr. Liège Belgii. — A typo præcipue sporidiis longioribus diversa.

115. **Leptosphæria hemicypta** Oud. Contr. Flor. Myc. Pays-Bas. XI, 3219 p. 21. — Peritheciis epiphyllis, membranaceis, nigris, $\frac{1}{6}$ mm. latis, depresse globosis, semiimmersis, pro dimidia fere parte occultatis sub epidermidis portiuncta antice hiante, postice vero a contextu non soluta, ostiolo simplici vel breviter papilloso præditis; ascis numerosis, cylindricis, $50-70 \approx 10-12$, versus basim angustioribus, 8-sporis, partim paraphysibus filiformibus, subtilissimis, partim pseudo-paraphysibus piriformibus articulatis, longe pedicellatis, sursum in appendicem mucroniformem abeuntibus, stipatis; sporidiis distichis, dilutissime fuscescentibus, fusiformibus, paulum curvatis, 5-guttulatis, ob maturitatis stadium imperfectum septis hucusque destitutis.

Hab. in foliis *Caricis* cujusdam, Putten Hollandiæ.

116. **Leptosphæria folliculata** Ell. et Ev. Proc. Phil. 1890, pag. 237. 3220 — Maculis pallide albis, ellipsoideis, $2-4 \approx 1-1,5$; peritheciis immersis, apice sub prominulo, 1-6 in quaque macula, $60-75 \mu$. diam., ascis clavato-cylindraceis, $50 \approx 10-12$; sporidiis distichis, oblongis v. clavato-oblongis, 2-septatis, ad septa subconstrictis, $12-15 \approx 3$, flavidobrunneis, utrinque obtusis.

Hab. in foliis *Caricis folliculatae* in Amer. bor. — Ab aliis speciebus ad *Caricem* viventibus differt maculis distinctis sporidiisque minoribus.

117. **Leptosphæria marina** Ellis et Ev. in Journ. of Myc. 1885, p. 43. 3221 — Peritheciis irregulariter sparsis, subellipticis, $\frac{1}{3}-\frac{3}{4}$ mm. diam., sursum conoideis, epidermide integra, nigrescente tectis, apice subtruncatis, ostiolo papilliformi, mox deciduo præditis, demum late perforatis; ascis $112-150 \approx 25-35$, basi constrictis, substipitatis; sporidiis bi-tristichis, fusiformibus vel clavato-fusiformibus, luteolis, 1-3-septatis, (sæpe 1-septatis), medio leniter constrictis, $50-60 \approx 10-12$, utrinque subobtusis.

Hab. in culmis emortuis *Spartinæ*, Cape May Amer. bor. (CAROLINA TREAT).

118. **Leptosphæria sticta** E. et E. in Journ. of Myc. I, pag. 43. — 3222 Peritheciis sparsis, suboyatis, membranaceis, epidermide nigrefacta, ostiolo punctiformi perforata, tectis dein prominentibus; ascis $100-125 \approx 15-20$; paraphysibus numerosis; sporidiis 1-2-seriatis,

fusiformibus, pallide luteis, 5-septatis, ad septa leniter constrictis, utrinque obtusis, 30-40 \approx 7-9.

Hab. in culmis *Spartinæ*, Cape May Amer. bor. (CAROLINA TREAT).

119. **Leptosphaeria Spartinæ** Ell. et Ev. in Journ. of Myc. 1885, 3223 p. 43. — Peritheciis gregariis, $\frac{1}{4}$ mm. diam., epidermide nigrescente, ostiolo papilliformi, late pertuso rupta, elevata tectis; ascis 115-120 \approx 12-15; paraphysibus, filiformibus, obvallatis, 8-sporis; sporidiis distichis, late fusoideis, pallide luteis, 5-septatis, 35-45 \approx 8-10.

Hab. in culmis emortuis *Spartinæ*, Cape May Amer. bor. (CAROLINA TREAT).

120. **Leptosphaeria lineolaris** Niessl in Herb., Linhart fung. hung. 3224 n. 466. — Peritheciis minutis, circiter 0,15-0,20 mm. diam., seriatim dispositis, saepe in lineas breves confluentibus, submembranaceis, depresso globosis vel e latere compressis elongatisque, atrofuscis, glabris, ostiolo punctiformi; ascis elongato-oblongis, stipite brevi, 43-58 \approx 10-12, octosporis, paraphysibus longioribus obvallatis; sporidiis 2-3-stichis, fusiformibus, inaequilateralibus v. parum curvatis, obtusiusculis, 5-7-loocularibus, vix nodulosis, dilute lutescentibus, 19-24 \approx 3-4.

Hab. in culmis exsiccatis Airæ cæspitosæ prope Petrozsény Hungariæ.

121. **Leptosphaeria Ammophilæ** Rehm Ascomyc. fasc. XIV, n. 691. 3225 — Peritheciis immersis dein prominulis, late ellipticis, ostiolo crasso, prominente donatis, nigris, levibus, 260-340 \approx 210-260; ascis oblongis, basi subinde leniter incrassatis, octosporis, 115-123 \approx 35; paraphysibus filiformibus obvallatis; sporidiis inordinate distichis, oblongis, 7-8-loocularibus, loculis quarto vel quinto subinde crassioribus, 38-45 \approx 12-14, luteis.

Hab. in caulinis siccis Psammæ arenariæ in Germania. — « Verisimiliter eadem ac *Lept. littoralis* ».

122. **Leptosphaeria Sporoboli** Ell. et Gall. Journ. Myc. 1889 p. 66. 3226 — Peritheciis sparsis, erumpenti-superficialibus subhemisphaericis, sublevibus, nigris, $\frac{1}{5}$ - $\frac{1}{4}$ mm. diam.; ostiolo brevi, crasso, mammiformi; ascis clavato-cylindraceis, 75-80 \approx 16-18, copiose paraphysatis; sporidiis conferte distichis, imbricatis, oblongo-fusoideis, utrinque subobtusis, rectis, 6-septatis, ad septa non v. deum leniter constrictis, circ. 22 \approx 7.

Hab. in culmis Sporoboli depauperati, Sand Coulee, Ca.

scade County, Montana, Amer. bor. (F. W. ANDERSON). — A *L. culmifraga* sporidiis brevioribus, constanter 6-septatis, a *L. culmicola* peritheciis subsuperficialibus differt.

123. **Leptosphæria conimbricensis** Berl. et Fr. Sacc. Contr. Myc. Lus. 3227 p. 5, t. 83, f. 5. — Peritheciis sparsis, minutis, epidermide tectis, globoso-conoideis, $\frac{1}{4}$ mm. diam., ostiolo papillato, minuto; ascis clavatis, sursum late rotundatis, subsessilibus, 80-90 \approx 8-9; sporidiis 8-sporis, subdistichis, fusoideis, 7-septatis, loculo quarto crassiore, 22-24 \approx 5-6, pallide flavo-virentibus.

Hab. in culmis Piptatheri miliacei, Coimbra Lusitaniæ.

124. **Leptosphæria culmifraga** (Fr.) Ces. De Not. — Syll. II, p. 75 — 3228 Var. **bromicola** Bres. in Rev. Myc. 1890 pag. 184. — Peritheciis sparsis v. in series congestis, globoso-conoideis, in ostium papillatum desinentibus, 300-320 μ . diam., atris, subsuperficialibus, basi ut plurimum filamentis fuscidulis interdum dichotomis, 2 μ . cr. præditis; ascis clavatis, stipitatis, paraphysibus filiformibus, guttulatis cinctis, 40-80 \approx 10-12; sporidiis octonis, superne 3-stichis, fusoideo-elongatis, curvulis, 6-8-septatis, loculo tertio interdum leviter crassiore, ad septum subconstrictis, flavis, 22-25 \approx 3,5-4.

Hab. in culmis Bromi asperi, pr. Leutschau Hungariæ. — Differt a typo peritheciis perfecte conoideis, subsuperficialibus, ascis sporidiisque minoribus.

125. **Leptosphæria phormicola** Cooke et Harkn. in Grevillea XIV, 3229 p. 10. — Gregaria; peritheciis punctiformibus, erumpentibus, epidermide tectis dein rimoso-emergentibus, atris, convexulis, papillatis; ascis clavatis, octosporis; sporidiis sublanceolatis, utrinque rotundatis, biserialibus, triseptatis, vix constrictis, pallide fuscis, 22 \approx 6.

Hab. in Phormio in California (HARKNESS).

126. **Leptosphæria Cattanei** Thuem. Pilze d. Reispf. p. 5. — Peritheciis sparsis, innato-erumpentibus, atris, globulosis, papillatis, parvulis, in macula subexarida griseola; ascis elongato-clavatis, 8-sporis, breve stipitatis, 100-110 \approx 10-13; sporidiis plerumque distichis, viridi-flaventibus, fusoideis, 3-septatis, ad septa vix constrictis, 22-28 \approx 4; paraphysibus nullis (?)

Hab. in culmis foliisque languidis Oryzæ sativæ, Aquileia Ital. boreal.

127. **Leptosphæria algida** Rost. Fung. Groenl. p. 560. — Peritheciis globulosis, 160 μ . diam.; ascis cylindraceo-clavatis, curvulis, 50-

- 792 Pyrenomyceteæ, Sphæriaceæ, Phæophragmiæ, Leptosphaeria.
54 \approx 12, aparaphysatis; sporidiis oblique monostichis, oblongis, saepe curvulis, flavis, 3-septatis, 16-20 \approx 6-7.
Hab. in foliis Catabrosæ algidæ in Groenlandia.
128. **Leptosphaeria Typharum** (Desm.) Karst. — Sacc. Syll. II, p. 64, 3232
var. **phragmitina** Karst. in Hedw. 1884, n. 1, pag. 5, et Rev. atque Synops. Ascom. fenn. pag. 54. — Peritheciis sparsis, innato-erumpentibus, glabris, atris, exiguis; ascis oblongatis, subsessilibus, 75-88 \approx 20-22; sporidiis distichis, oblongatis, utrinque obtusis, subrectis, 3-septatis, ad septa leviter constrictis, dilute luteolis, 24-29 \approx 9-11; paraphysibus parum notabilibus.
Hab. in foliis Phragmitis communis putrescentibus ad Mustiala Fenniæ.
129. **Leptosphaeria clavicarpa** E. et E. in Jour. of Myc. I, p. 43, — 3233
Peritheciis plerumque seriatis, in lineas $\frac{1}{4}$ -1 cm. longas dispositis, epidermide nigrefacta tectis dein prominentibus et epidermidem, ostiolis vix conspicuis, perforantibus; ascis latis, oblongo-cylindraceis, 75-100 \approx 22-25, obtusis, basi constrictis; paraphysibus filiformibus; sporidiis stipatis, polystichis, clavato-oblongis, leniter curvatis, apice obtusis vel rotundatis, basi abrupte constrictis, 7-9-septatis, maturis luteo-brunneis, 25-40 \approx 8-10, ad septa leniter constrictis; spermatiis in peritheciis similibus, oblongo-cylindraceis, leniter curvatis, hyalinis, 3-5 \approx 1 $\frac{1}{4}$.
Hab. in culmis emortuis Phragmitis communis, Spirit Lake, Iowa Amer. (PROF. S. C. ARTHUR).
130. **Leptosphaeria pachycarpa** Sacc. et March. in Rev. Myc. 1885. 3234
— Peritheciis gregariis, erumpenti-superficialibus, globoso-papillatis, $\frac{1}{3}$ mm. diam., duriusculis, atris; ascis tereti-clavulatis, brevè stipitatis, paraphysatis, 140 \approx 18, octosporis; sporidiis distichis, fusoides, utrinque acutiusculis, rectis curvulisve, 5-septatis, ad septum medium leniter constrictis, 35-40 \approx 6,5-7,5, fuligineis, loculis extimis subhyalinis.
Hab. in culmis Graminum majorum in Belgio. — Peritheciis crassiusculis sporidiisque utrinque subhyalinis statim dignoscitur.
131. **Leptosphaeria rhodophæa** Bizz. Fung. Ven. nov. v. crit. pag. 2, 3235
t. II, f. 3, a, b, c, d. — Peritheciis sparsis vel 2-3-aggregatis, subimmersis, globoso-depressiusculis, ostiolo papillulato, vix exerto, pertusis, 0,5 mm. diam; ascis cylindraceo-clavatis, 8-sporis, 80-100, parte sporif. 70-75 \approx 10, paraphysatis; sporidiis distichis, fuligineis, fusiformibus, 5-septatis, ad septa intermedia constrictis, loculo medio crassiore et rotundato, 28-30 \approx 7-7,5.

Hab. in culmis emortuis Arundinis Donacis in horto botanico patavino.

132. **Leptosphæria recessa** Passer. Diagn. F. N. IV, pag. 5. — Pe- 3236
ritheciis in gemmarum fovea aggregatis, erumpentibus, tandem nudatis, subglobosis, atris, ostiolo minuto papillari; ascis paraphysatis, elongato-clavatis, basi breviter attenuatis, 8-sporis, 65-90 \approx 10-12; sporidiis distichis v. oblique monostichis, fusiformibus, 5-septatis, loculo nullo inflato, flavidis, 25-27,5 \approx 6-5.

Hab. in culmis fracidis Arundinis Donacis, Vigheffio pr. Parma Ital.

133. **Leptosphæria grisea** Passer. Diagn. F. N. IV, p. 5. — Prithe- 3237
ciis in matrice dealbata maculam longitudinalem griseam sistentibus, globulosis, atris, erumpentibus, ostiolo acutiusculo; ascis tubulosis v. clavatis, sessilibus a paraphysatis?, 8-sporis, 75 \approx 10; sporidiis distichis v. oblique monostichis, fusiformibus, curvulis, 5-septatis, loculo altero ex intermediis tumido, melleis, 25-27,5 \approx 4,5-5.

Hab. in culmo fracio Sorgi vulgaris, Vigheffio pr. Parma Italiae.

134. **Leptosphæria fuscidula** Passer. Diagn. F. N. I, n. 50. — Perithe- 3238
ciis pusillis, sparsis v. subgregariis, in maculis linearibus albican-tibus, subsuperficialibus atris, ostiolo acuto; ascis cylindricis, a para-physatis, 8-sporis, 55-80 \approx 10-12; sporidiis fusiformibus, biseri-ibus, rectis v. curvulis, triseptatis, non constrictis, loculo altero subultimo non v. vix crassiore, olivaceo-fuscis, 17-20 \approx 5-6.

Hab. in foliis aridilis Melicæ altissimæ in horto botanico parmensi Italiae bor.

135. **Leptosphæria Hierochloæ** Oudem. Cont. à la Flore myc. de No- 3239
waja Semlja p. 10, t. I, f. 3. — Peritheciis foliicolis, sparsis, pri-mitus occultatis, postremo erumpentibus, subglobosis, atris, struc-turæ parenchymatosæ flaccidæ, simpliciter perforatis (i. e. ostiolo nullo vel vix prominente), 180-250 μ . diam.; ascis brevissime pe-dicellatis, parum curvatis, paraphysatis, 90-95 \approx 30, membrana ubique æquali, 8-sporis; sporidiis tristichis, oblongo-fusoideis, re-ctis, utrinque obtusatis, mellei coloris, 5-septatis, ad septa levis-sime constrictis, 30 \approx 9, loculo quarto omnium maximo.

Hab. in foliis Hierochloæ alpinæ, Nowaja Semlja.

136. **Leptosphæria fuegiana** Speg. Fueg. n. 231. — Maculis nullis; 3240
peritheciis sparsis, hypophyllis, immersis, non v. vix prominulis, 150-200 μ . diam., atris, glabris, membranaceis, contextu pellucido, grosse parenchymatico-fuligineo, ostiolo subcarbonaceo, papillulato,

epidermidem perforante sed non exerto donatis; ascis cylindraceis, antice rotundatis, postice breviter cuneato-rotundatis, stipite brevissimo crassoque suffultis, 40-45 \approx 10, 8-sporis, apophysatis; sporidiis di-tri-stichis, elliptico-fusoideis, utrinque acutiuscule rotundatis, 18-20 \approx 5, triseptatis, ad septum medium non v. leniter constrictis, loculis eguttulatis, secundo supero crassiore, flavescentibus.

Hab. in Hierochloæ antarcticæ foliis culmisque putrescentibus in pratis pr. Penguin rookery, Staten island. Amer. austr.

137. **Leptosphaeria Puiggarii** Speg. F. Puigg. n. 281. — Maculis ³²⁴¹ fuscis, minutis indeterminatis; peritheciis globosis, infossis, seriatis, atris, glabris, 200-250 μ . diam.; ascis fusoideis, 8-sporis, 90-100 \approx 20, paraphysatis; sporidiis distichis fusoideis, 5-septatis, melleis, 28-30 \approx 8-9.

Hab. ad folia emortua Andropogonis saccharoidis in pratis pr. Apiah Brasiliæ.

138. **Leptosphaeria Brizæ** Passer. Diagn. F. N. I, n. 51. — Peritheciis ³²⁴² superficialibus, sparsis, pusillis, subglobosis, atro-nitidis; ascis oblongis, saccato-ventricosis, basi breviter angustatis, 8-sporis, 55 \approx 17-18; sporidiis tristichis, elongato-fusiformibus rectis, vel curvulis, triseptatis, flavidis, ad septum medium vix constrictis, interdum scissilibus, loculis quandoque guttulatis, 35-40 \approx 5,5-6; paraphysibus non visis.

Hab. in culmis et pedunculis siccis Brizæ mediæ, Monte Prinzena iu agro parmensi Italiæ.

139. **Leptosphaeria dactylina** Passer. Diagn. F. N. I, n. 52. — Peritheciis ³²⁴³ remote sparsis, tectis, epidermidem sublevantibus, globosis, atris; ascis clavatis, paraphysatis, 8-sporis, 62-85 \approx 10; sporidiis subtristichis, multiseptatis, fusiformibus, loculis guttulatis, æqualibus, flavis, 20-25 \approx 3,-5-4.

Hab. in culmis aridis Dactylidis glomeratæ, Vigheffio pr. Parma Italiæ. — *L. seriatam* fructificatione æmulatur, sed peritheciis multo grandioribus, diversissime dispositis, primo obtutu distinguenda.

140. **Leptosphaeria seriata** Passer. Diagn. F. N. I, n. 49. — Peritheciis ³²⁴⁴ exiguis, atris, per epidermidem rimosam parallele seriatis, ostiolo minuto, acuto; ascis oblongo-clavatis, paraphysatis, 8-sporis, 50-60 \approx 9-10; sporidiis distichis, fusiformibus, rectis v. curvulis, 9-11-septatis, loculis æqualibus, flavidis, 22-25 \approx 2,5-3.

Hab. in culmo fracido Zeæ Maydis, socio *Epicocco neglecto*, Vigheffio pr. Parma Italiæ.

141. **Leptosphaeria patellæformis** Passer. Diagn. F. N. III, n. 21. — 3245

Peritheciis subgregariis v. sparsis, subglobosis, ostiolo minutissimo vix papillari, atris, siccitate collapsis, scutellatis; ascis elongato-subclavatis, paraphysibus filiformibus superatis, 8-sporis, 80-100 \approx 8-10; sporidiis fusiformibus, 3-septatis, distichis v. oblique monostichis, dilute flavis, senio fuscouscentibus, 15-17,5 \approx 5.

Hab. in culmo fracido Zeæ Maydis, Vigheffio prope Parma Italiæ.

142. **Leptosphaeria rhizomatum** Passer. Diagn. F. N. III, n. 22. — 3246

Peritheciis parce gregariis, tectis, globosis, atris, ostiolo epidermidem perforante v. subinde pontiformi sublevante, tandem nudatis, glabris; ascis paraphysatis, plus minus late clavatis, basi sensim attenuatis, noduloso-stipitatis, 8-sporis, 62-87 \approx 6-10; sporidiis distichis v. oblique monostichis, in ascis latioribus superne tristichis, fusiformibus, 7-9-septatis, loculis æqualibus v. altero ex interme- diis vix tumidulo, e luteo fuscis, ad apices acuminatis et quandoque apiculatis, 22-25 \approx 2,5-3,5.

Hab. in rhizomatibus Cynodontis Dactyli et stolonibus Agrostidis vulgaris, Vigheffio pr. Parma Italiæ. — A *L. culmifraga* (Fr.) non longe distare videtur, sed peritheciis glabris et sporidiorum longitudine, satis diversa.

143. **Leptosphaeria pleurospora** Niessl in Rehm Ascom. Ill. n. 937. 3247

— Peritheciis sparsis, sub epidermide adusta v. expallente et de-albata nidulantibus, minutis, vix 150 μ . diam., depresso-globosis, membranaceis, atris, ostiolo punctiformi; ascis paucis in singulis peritheciis, oblongis demum, per extensionem sæpe clavatis, stipitatis, 80-135 \approx 24-36, 8-sporis; sporidiis oblique 2-stichis, fusoido-cylindraceis, inferne parum angustatis, utrinque obtusis, plurimque rectis, sed inæquilateralibus, 7-cellularibus, loculo tertio paullo protuberante, obscure fuliginosis, demum atris, subopacis, episporio longitudinaliter elegantissime striato, 33-38 \approx 7-9.

Hab. in culmis siccis Airæ cæspitosæ pr. Lofer Salisburgi (NIESSL).

144. **Leptosphaeria pachytheca** Hariot et Briard Rev. Myc. 1890 Oct. 3248

— Peritheciis parvis, subepidermicis, nigris, 100-120 μ . diam., in series longitudinales plus minus adproximatis; ascis ovoideis, sessilibus, 8-sporis, 40-44 \approx 14-16, aparaphysatis; sporidiis oblongis, ellipsoideis, subfusiformibus, 1-3-septatis, obtusiusculis, olivaceo-pallidis, non v. parum constrictis, 8-12 \approx 3-4.

Hab. ad culmos et folia sicca Nardi strictæ in planicie Foolz,

Aube Galliæ (P. HARIOT). — **Lept. Fuckelii** Niessl form. **filamentifera** Karst. Fragm. Myc. XXIV, p. 3. In vaginis Phalaridis arundinaceæ prope Mustiala est obvia. — **Lept. littoralis** Sacc. Huc, teste Berl. Alcun. Lept. p. 15, t. II, f. 13, spectant *Sphæria sabuletorum* Berk. pro parte et *Lept. Ammophilæ* Rehm, sed non *Sph. Ammophilæ* Lasch. — **Lept. culmorum** Auersw. var. **hungarica** Rehm Ascom. n. 941: peritheciis plerumque gregariis in maculis subatris. In calamis Luzulæ albidae, Eperjes Hungariæ.

IV. In Acotyledoneis.

145. **Leptosphaeria lycopodicola** Peck 38 St. Rep. Mus. pag. 105, 3249 tab. 2, fig. 16-19. — Peritheciis minutis, 120-150 μ . diam., sphæroideis vel ellipsoideis, erumpentibus, nigris, ascis subcylindraceis, subsessilibus, 65-75 \approx 8-10; sporidiis oblongis vel subfusiformibus, pallide coloratis, 3-5-septatis, 20-25 \approx 4-5; paraphysibus filiformibus.

Hab. in pedunculis emortuis Lycopodii clavati in montibus Adirondack Amer. bor. — **Lept. helvetica** Sacc. et Sp. — Syll. II. p. 80. — Var. **major** Rehm Ascom. Ill. n. 884: a typo differt sporidiis multo majoribus, 18-24 \approx 6, hyalinis, 3-septatis; ascis 90-18. An potius ætatis gradus?

146. **Leptosphaeria apocalypta** (Rehm) Winter Die Pilze II, p. 442, 3250 *Xenosphaeria apocalypta* Rehm Ascomyc. n. 346, — Peritheciis sparsis vel subgregariis, superficialibus, majusculis, fere globosis, atris, minute papillatis, guttulas oleosas seriatim dispositas continentibus; ascis clavatis, tunica apice incrassata, 4-6-sporis, 100-120 \approx 12; sporidiis monostichis, late ellipsoideis vel oblongis, saepe curvis vel inæquilateralibus, 4-8-cellularibus, obscure brunneis, 24-31 \approx 9-10,5; paraphysibus? crassis, guttulatis longe filiformibus.

Hab. in superficie thalli Stereocauli alpini in Germania.

147. **Leptosphaeria Sphyridiana** (Lahm) Winter Die Pilze II, p. 442, 3251 *Xenosphaeria Sphyridiana* Lahm in litt. et in Arnold Lichen. Fragm. XVI, pag. 29. — Peritheciis punctiformibus, nigris, thallo immersis; ascis cylindraceis, elongatis, 8-sporis, paraphysibus distichis, validis obvallatis; sporidiis 4-cellularibus, saepe 4-guttulatis, medio leniter constrictis, utrinque obtusis, brunneis, 24-27 \approx 6-9.

Hab. in pagina superiore thalli Sphyridii fungiformis in Germania.

148. **Leptosphaeria stereicola** Ellis Amer. Nat. 1883, p. 317. — Pe. 3252
ritheciis erumpentibus, hemisphæricis, nigris, rugosis, $\frac{1}{9}$ mm. diam.,
ostiole acute papillato; ascis subcylindricis, $57 \approx 7,5-8$, paraphysa-
tis (?); sporidiis biseriatis, oblongo-fusoideis, 3-septatis, fuligineis,
 $11-13 \approx 3-3,5$.

Hab. in hymenio *Sterei bicoloris*, Decorah. Iowa Amer.
bor. (E. W. HOLWAY).

149. **Leptosphaeria fungicola** Winter. in Hedwigia 1886, p. 101. — 3253
Peritheciis sparsis, superficialibus (an semper?), minutissimis, glo-
bosis, poro simplici pertusis, membranaceis, atris, glabris, 100μ .
diam.; ascis obclavatis, e basi lata sursum longe attenuatis, sessili-
bus, 8-sporis, $30-44 \approx 9-10,5$; sporidiis conglobatis, fusiformibus,
inæquilateralibus vel parum curvatis, 5-septatis, cellula tertia la-
tissima, dilutissime viridulis, $12-16 \approx 3,5$.

Hab. in hymenio *Sterei subpileati* vetusto in insula S.
Thomæ Afr. occ.

150. **Leptosphaeria Coræ** Pat. Journ. Bot. 1888, p. 150, f. 5. — Pe. 3254
ritheciis numerosis, sparsis, nigris, subglobosis, $\frac{1}{3}-\frac{1}{2}$ mm. diam.,
innatis dein emergentibus, hyphis mycelicis brunneis affixis, ostiole
dein late aperto; ascis ventricosis, subsessilibus, $80 \approx 25$, apice in-
crassato, tenuato-rotundato; sporidiis octonis, ovato-oblongis, 5-
septatis, non constrictis, pallide fuligineis, $20-24 \approx 6-7$.

Hab. in thallo *Coræ Pavoniæ* in Amer. centr. — Asci jodo
cœrulescentes v. partim roseo-violascentes.

151. **Leptosphaeria marina** Rostr. Journ. Bot. Copenhag. XVII, p. 234. 3255
— Peritheciis in matrice atrata gregariis, globosis, tectis; ascis
fusoideis paraphysatis, 8-sporis, $50 \approx 10$; sporidiis fusoideis, pallide
flavescentibus, 1-3-septatis, $16 \approx 4$.

Hab. in thallo *Chondri crispi* ad littora marina Daniæ. —
Maculæ in matrice 1-2 mm. diam.

CHITONOSPORA Bomm. Rouss. Sacc. nov. gen. (Etym. *chiton* tunica
et *spora*). — Perithecia subglobosa, papillata, atra, subcutanea,
submembranacea. Asci elongati, facile evanidi, paraphysati. Spor-
idia ellipsoideo-oblonga, subtriseptata, episporio crassissimo fuligi-
neo, endosporio subhyalino prædita.

1. **Chitonospora ammophila** Bomm. Rouss. Sacc. sp. n. — Peritheciis 3256
glabris, 300μ . diam., epidermide leniter atrata tectis, ostiole brevi
conoideo epidermidem perforante; ascis crassis, cylindraceis, sessili-
bus, $75-90 \approx 15-16$; paraphysibus ramulosis, flexuosis, guttulatis,

parum distinctis; sporidiis compressis, tereti-ellipsoideis, subinde utrinque obtuse mucronulatis, fuligineis, 3-septatis, 24-27 \approx 12-15, episporio denique inæqualiter lacerato et endosporium didymum, 3-septatum 21-22 \approx 9 subhyalinum relinquente.

Hab. in foliis emortuis *Psammæ arenariæ*, Ostende Belgii.
— Sporidia sæpe medio opacissima, septa celant et quum utrinque sint pallidiora, *samaram* imitantur.

HEPTAMERIA Rehm et Thuem. — Sacc. Syll. II, p. 88.

1. **Heptameria uncinata** (Niessl) Rehm Ascomyc. fasc. XV, n. 735. 3257

Leptosphaeria uncinata Niessl in litt. *Lept. obesa* Rehm Ascomyc. Lojk. pag. 54. n. 70. — Peritheciis parenchymaticis, brunneis, ascis clavatis, octosporis, 150 \approx 18; sporidiis fusoideis, rectis vel subcurvatis, acutatis, 9-cellularibus, cellula media magna, 9 μ . longa, fusca nucleo magno, centrali prædita, ceteris cellulis latiore 42 \approx 5, distichis; cellulis lateralibus fuscidulis; paraphysibus ramosis.

Hab. ad caules sicclos Artemisiæ campestris in horto Eszterházyano Kaposvar (Somogy) (Lojka) et prope Brünn in Moravia (NISSL). — Species quoad sporidia egregia!

CLYPEOSPHÆRIA Fuck. — Sacc. Syll, II, p. 90.

1. **Clypeosphaeria Morreni** (West.) Sacc. in Rev. Myc. 1884. A *Clypeosphaeria Notarisii*, cui peraffinis, ascis 98 \approx 8-10; sporidiis cylindraceis, curvulis, utrinque rotundatis, 20-21 \approx 4,5, 3-septatis, non constrictis, fuligineis dignoscitur. 3258

Hab. in caulis Epilobii spicati, Louette-Saint Pierre Gallæ (G. AUBERT).

2. **Clypeosphaeria Asparagi** Fuck. Wint. Hedw. 1887, p. 62. — Pe- 3259 ritheciis plerumque gregariis, rarius sparsis 2-pluribus confertis, semper peridermio aterrimo tectis, globosis, ostiolo papilliformi prominulus, membranaceis, atris, 250-300 μ . diam.; ascis anguste clavatis, deorsum longe attenuatis, 8-sporis, 66-77 \approx 8-9; sporidiis indistincte distichis, fusoideis, plerumque curvatis, 3-4-septatis, loculo secundo valde incrassatis, 21-28 \approx 4, luteolis; paraphysibus filiformibus.

Hab. in caulis Asparagi officinalis in Rhenogovia. — Sec. Winter l. c. eadem ac *Diap. Asparagi* Fuck.

CHÆTOSPHÆRIA Tul. — Sacc. Syll. II, p. 92.

1. **Chætosphæria ornata** Harkn. New Calif. Fungi p. 25. — Pe- 3260
ritheciis superficialibus, orbicularibus, nigris, non rugosis, 360 μ .
diam., setis hyphoideis, $\frac{1}{2}$ mm. vel amplius longis, nigris, non acu-
minatis, quandoque septatis; ascis octosporis, clavatis, in pedicellum
longum, guttulis oleosis vel granulis farctum, desinentibus 100 \approx
8-10; paraphysibus vix visilibus; sporidiis fusiformibus, triseptatis,
loculo medio olivaceo-brunneo, extimis hyalinis, 16 \approx 6.

Hab. in corticibus putridis *Eucalypti*, San Francisco Cali-
forniæ.

2. **Chætosphæria Saccardiana** Schulz. in Rev. Myc. 1884 et III. Fung. 3261
Slav. n. 74. — Peritheciis densiuscule gregariis, globoso-depressis,
subsuperficialibus, $\frac{1}{3}$ mm. diam., nigris rugulosis, hyphulis laxe
vestitis; ostiolo cylindraceo, truncato, perithecio plus quam dimidio
breviore; ascis tereti-clavatis, breve stipitatis, paraphysatis, octo-
sporis; sporidiis subdistichis oblongo-fusoideis, medio constrictis,
1- dein, 3-septatis 4-guttulatis, saepius curvulis, utrinque obtusiu-
sculis, dilute fuscis.

Hab. in ligno putrescente *Carpini* in Slavonia, socia forma
conidiophora *Helminthosporioidea*.

3. **Chætosphæria pezizæformis** Schulz. in Rev. Myc. 1884 et Ill. 3262
Fung. Slav. n. 62. — Peritheciis laxe v. dense gregariis, globosis,
superficialibus, $\frac{1}{6}$ - $\frac{1}{4}$ mm. diam., e lutescenti nigris, asperulis, in-
ter hyphas demetiaceas (*Helminthosporii obclavati*) interspersis,
apice convexis, tandem collabascendo umbilicato-subpezizæformibus;
ascis crasse clavatis, deorsum acuminatis, brevissime stipitatis.
indistincte paraphysatis, octosporis; sporidiis subtristichis, fusoideis,
incurvatis utrinque obtuse attenuatis, 7-septatis, ad septa non con-
strictis, 44-50 \approx 8-10, fusco-lutescentibus, loculis extimis hyalinis.

Hab. in ramis decorticatis *Carpini Betuli*, Vinkovce Sla-
voniæ.

4. **Chætosphæria rubiginosa** (Cooke) Sacc. *Byssosphæria rubiginosa* 3263
Cooke in Grev. XV, p. 80, *Gibbera rubiginosa* Cooke in Herb.
— Peritheciis gregariis, superficialibus, subiculo tenui, filamentoso
fusco insidentibus, globosis, atro-rubiginosis, subrugulosis, ostiolo
impresso; ascis clavato-cylindricis, 8-sporis; sporidiis biserialibus,
lanceolatis, triseptatis, medio constrictis, hyalinis, demum pallido
fuscis, 20-24 \approx 4-6.

Hab. in ligno putri, Poughkeepsie, New York Amer. bor.

5. **Chætosphaeria longipila** Peck 42 Rep. p. 35. — Peritheciis minutiissimis, gregariis, subiculo ex hyphis cinereis vel griseolo-brunneis, longissimis gracilibus, arachnoideis insidentibus v. involutis; ascis oblanceolatis, p. sp. $40 \pm 12,5$; sporidiis confertis v. distichis, rectis, 3-septatis, loculis mediis coloratis, apicalibus hyalinis, $10-12,5 \pm 5-6,5$.
- Hab.* in doliis vetustis Flatbush Amer. bor. (ZABRISKIE). — *C. phæostromati* et *C. phæostromoidi* affinis, sed subiculo pallidore sporidiisque brevioribus et rectis.
6. **Chætosphaeria Crepini** Sacc. Bomm. et Rouss. Fl. myc. Belg. II, p. 17. — Peritheciis subsphæricis, $\frac{2}{5}$ mm. diam., ostiolo punctiformi dehiscentibus, tomento brunneo, præcipue basi, obductis, subiculo repenti, paucoso, brunneo insertis; ascis tereti-clavatis, 8-sporis; paraphysibus filiformibus copiosis; sporidiis monostichis fusoideo-naviculäribus, hyalinis, 1-septatis, dein olivaceis, 3-septatis, 4-guttatis, medio contractis, utrinque attenuatis, muticis, $24 \pm 7-7,5$.
- Hab.* in caulinis siccis Epilobii hirsuti, Groenendael Belgii. — Ab *Herpotrichia Rubi* subiculo copioso, a *Chæt. callimorpha* ostiolo haud cupulato differt.
7. **Chætosphaeria holophæa** (B. et C.) Sacc. *Byssosphaeria holophæa* B. et C. in Cooke Grev. XV, pag. 82. — Peritheciis subglobosis, pilosis, gregariis, bysso atro-fusca, rigida, nidulantibus, $\frac{1}{2}$ mm. diam.; ascis clavatis, octosporis; sporidiis biseriatis, elongato-ellipticis, medio constrictis, triseptatis, fuscis, $22-24 \pm 8$.
- Hab.* ad ramos in Pennsylvania Amer. bor. — In *Chætosphaeria albo-ferruginea* (Crouan) Sacc. Syll. II, n. 3216, sec. Grev. XVI, p. 47, sunt sporidia lanceolata, uniseptata, 4-6-nucleata, demum forte 3-5-septata, hyalina $45-50 \pm 8$. Nascitur quoque prope Carlisle Britanniæ in Calluna.
8. **Chætosphaeria exilis** (Schw.) Sacc. *Sphæria exilis* Schw. sec. Cooke Grev. XV, p. 82. — Peritheciis minutis globosis, $\frac{1}{8}$ mm. diam.; sporidiis fusoideis triseptatis, brunneis, $16-18 \pm 5$.
- Hab.* in lignis putridis, Nazareth Amer. bor.
9. **Chætosphaeria squamulata** (Schw.) Sacc. *Sphæria* Schw. sec. Cooke Grev. XV, p. 83, Sacc. Syll. II, n. 4290. — Sporidia lanceolata, 3-5-septata, fusca, $20-40 \mu$. longa.
- OHLERIA** Fuck. — Syll. II, p. 96, Berl. Ic. I, p. 28.
1. **Ohleria adjecta** Passer. Diagn. F. N. I, n. 53. — Peritheciis in ligno exarido et nigrificato sparsis v. subgregariis, parvulis, sphæ-